

Aquachloridobis[2-(1,3-thiazol-4-yl- κ N)-1H-benzimidazole- κ N³]nickel(II) nitrate

Peng Liang, Wei Wu, Wei-Man Tian and Xian-Hong Yin*

 Department of Chemistry and Chemical Engineering, Guangxi University for Nationalities, Nanning 530006, People's Republic of China
 Correspondence e-mail: 6628yxh@163.com


Received 19 May 2012; accepted 29 June 2012

 Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.007$ Å; R factor = 0.054; wR factor = 0.174; data-to-parameter ratio = 12.8.

In the title compound, $[\text{NiCl}(\text{C}_{10}\text{H}_7\text{N}_3\text{S})_2(\text{H}_2\text{O})]\text{NO}_3$, the Ni^{II} ion is coordinated by four N atoms from two chelating 2-(1,3-thiazol-4-yl)-1H-benzimidazole ligands, one Cl atom and one water molecule in a distorted octahedral geometry. In the crystal, $\text{O}-\text{H}\cdots\text{O}$, $\text{N}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{Cl}$ hydrogen bonds link the complex cations and nitrate anions into a three-dimensional network. $\pi-\pi$ interactions between the thiazole and imidazole rings and between the thiazole and benzene rings are observed [centroid-centroid distances = 3.592 (3) and 3.735 (3) Å].

Related literature

For background to the synthesis and properties of benzimidazole derivatives, see: Agh-Atabay *et al.* (2003); Devereux *et al.* (2004); Inoue *et al.* (2002). For a related structure, see: Mothilal *et al.* (2004).



Experimental

Crystal data

 $[\text{NiCl}(\text{C}_{10}\text{H}_7\text{N}_3\text{S})_2(\text{H}_2\text{O})]\text{NO}_3$ $M_r = 576.68$

 Monoclinic, $P2_1/c$
 $a = 16.091$ (5) Å
 $b = 11.189$ (3) Å
 $c = 13.931$ (3) Å
 $\beta = 113.275$ (3)°
 $V = 2304.1$ (11) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 1.18$ mm⁻¹
 $T = 296$ K
 $0.26 \times 0.25 \times 0.24$ mm

Data collection

 Bruker SMART APEX CCD diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\text{min}} = 0.748$, $T_{\text{max}} = 0.764$

 11654 measured reflections
 4050 independent reflections
 3129 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.072$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.054$
 $wR(F^2) = 0.174$
 $S = 1.07$
 4050 reflections

 316 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.80$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.53$ e Å⁻³
Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1}-\text{H1A}\cdots\text{O3}^{\text{i}}$	0.84	2.00	2.751 (5)	149
$\text{O1}-\text{H1B}\cdots\text{O2}^{\text{ii}}$	0.79	2.53	3.293 (6)	160
$\text{O1}-\text{H1B}\cdots\text{O4}^{\text{ii}}$	0.79	2.38	3.025 (5)	139
$\text{N4}-\text{H4}\cdots\text{O3}^{\text{iii}}$	0.86	2.35	2.962 (5)	129
$\text{N4}-\text{H4}\cdots\text{O4}^{\text{iii}}$	0.86	2.14	2.996 (5)	173
$\text{N7}-\text{H7}\cdots\text{Cl1}^{\text{iv}}$	0.86	2.35	3.159 (4)	157

 Symmetry codes: (i) $x - 1, y, z$; (ii) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$; (iii) $x - 1, -y + \frac{3}{2}, z - \frac{1}{2}$; (iv) $-x + 1, -y + 1, -z + 1$.

Data collection: SMART (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: DIAMOND (Brandenburg, 1999); software used to prepare material for publication: SHELXTL (Sheldrick, 2008).

This work was supported by the Key Scientific Research Projects of Guangxi University for Nationalities (2009MDZD42) and the Innovation Project of Guangxi University for Nationalities (gxun-chx2011092).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2553).

References

- Agh-Atabay, N. M., Dulger, B. & Gucin, F. (2003). *Eur. J. Med. Chem.* **38**, 875–881.
- Brandenburg, K. (1999). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2007). *SMART* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Devereux, M., McCann, M., Shea, D. O., Kelly, R., Egan, D., Deegan, C., Kavanagh, K., McKee, V. & Finn, G. (2004). *J. Inorg. Biochem.* **98**, 1023–1031.
- Inoue, Y., Hoshino, M., Takahashi, H., Noguchi, T., Murata, T., Kanzaki, Y., Hamashima, H. & Sasatru, M. (2002). *J. Inorg. Biochem.* **92**, 37–42.
- Mothilal, K. K., Karunakaran, C., Rajendran, A. & Murugesan, R. (2004). *J. Inorg. Biochem.* **98**, 322–332.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2012). E68, m1029 [https://doi.org/10.1107/S1600536812029728]

Aquachloridobis[2-(1,3-thiazol-4-yl- κ N)-1H-benzimidazole- κ N³]nickel(II) nitrate

Peng Liang, Wei Wu, Wei-Man Tian and Xian-Hong Yin

S1. Comment

Thiabendazole (TBZ) aroused considerable interest in biology and medicine due to its antiproliferative activities. It is an antimicrobial drug belonging to the benzimidazole derivatives and exhibits wide applications in human and veterinary medicine (Agh-Atabay *et al.*, 2003; Devereux *et al.*, 2004; Inoue *et al.*, 2002). As part of our studies of the synthesis and characterization of these compounds, we report here the synthesis and crystal structure of the title compound (Fig. 1).

In the crystal, intermolecular O—H \cdots O, N—H \cdots O and N—H \cdots Cl hydrogen bonds link the complex cations and nitrate anions into a three-dimensional network (Table 1, Fig. 2) (Mothilal *et al.*, 2004). π – π interactions between the thiazole and imidazole rings and between the thiazole and benzene rings are observed [centroid–centroid distances = 3.592 (3) and 3.735 (3) Å]. In the complex cation, the two ligand planes are nearly perpendicular to each other, with a dihedral angle of 89.14 (7)°.

S2. Experimental

A solution of TBZ (0.201 g, 1 mmol) in 3 ml DMF was added dropwise with stirring at room temperature to a solution of Ni(NO₃)₂·6H₂O (0.290 g, 1 mmol) and NiCl₂·6H₂O (0.237 g, 1 mmol) in a mixture of 10 ml water and 5 ml ethanol. Then an aqueous solution of sodium hydroxide was added dropwise with stirring to adjust the pH value of the solution being 6. The resulting mixture was sealed in a 23 ml Teflon-lined stainless reactor, kept under autogenous pressure at 403 K for 72 h, and then slowly cooled to room temperature at a rate of 5 K per hour. Dark-green block crystals suitable for X-ray diffraction were isolated, washed with ethanol and dried in air (yield: 65% based on Ni).

S3. Refinement

H atoms on C and N atoms were positioned geometrically and refined as riding atoms, with C—H = 0.93 and N—H = 0.86 Å and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$. The water H atoms were located from a difference Fourier map and refined as riding with $U_{\text{iso}}(\text{H}) = 0.079 \text{ \AA}^2$.




Figure 1

The Molecular structure of the title compound. Displacement ellipsoids are drawn at the 50% probability level. H atoms have been omitted for clarity.




Figure 2

Crystal packing of the title compound. Dashed lines denote hydrogen bonds.

Aquachloridobis[2-(1,3-thiazol-4-yl- κ N)-1H-benzimidazole- κ N³]nickel(II) nitrate

Crystal data

$[\text{NiCl}(\text{C}_{10}\text{H}_7\text{N}_3\text{S})_2(\text{H}_2\text{O})]\text{NO}_3$

$M_r = 576.68$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 16.091(5) \text{ \AA}$

$b = 11.189(3) \text{ \AA}$

$c = 13.931(3) \text{ \AA}$

$\beta = 113.275(3)^\circ$

$V = 2304.1(11) \text{ \AA}^3$

$Z = 4$

$F(000) = 1176$

$D_x = 1.662 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 2584 reflections

$\theta = 2.3\text{--}26.6^\circ$

$\mu = 1.18 \text{ mm}^{-1}$

$T = 296$ K $0.26 \times 0.25 \times 0.24$ mm
 Block, green

Data collection

Bruker SMART APEX CCD diffractometer Radiation source: fine-focus sealed tube Graphite monochromator φ and ω scans Absorption correction: multi-scan (SADABS; Sheldrick, 1996) $T_{\min} = 0.748$, $T_{\max} = 0.764$	11654 measured reflections 4050 independent reflections 3129 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.072$ $\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 1.4^\circ$ $h = -18 \rightarrow 19$ $k = -13 \rightarrow 13$ $l = -14 \rightarrow 16$
--	--

Refinement

Refinement on F^2 Least-squares matrix: full $R[F^2 > 2\sigma(F^2)] = 0.054$ $wR(F^2) = 0.174$ $S = 1.07$ 4050 reflections 316 parameters 0 restraints Primary atom site location: structure-invariant direct methods	Secondary atom site location: difference Fourier map Hydrogen site location: inferred from neighbouring sites H-atom parameters constrained $w = 1/[\sigma^2(F_o^2) + (0.083P)^2 + 1.4424P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\text{max}} = 0.045$ $\Delta\rho_{\text{max}} = 0.80 \text{ e } \text{\AA}^{-3}$ $\Delta\rho_{\text{min}} = -0.53 \text{ e } \text{\AA}^{-3}$
--	--

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0.26114 (3)	0.47368 (5)	0.32612 (4)	0.0352 (2)
Cl1	0.31740 (8)	0.27662 (10)	0.40665 (10)	0.0490 (4)
S1	0.36620 (9)	0.71229 (11)	0.62788 (10)	0.0507 (4)
S2	0.13043 (9)	0.27528 (12)	0.00588 (11)	0.0552 (4)
O1	0.13949 (19)	0.4517 (3)	0.3478 (3)	0.0412 (7)
O3	1.0666 (2)	0.6635 (3)	0.3784 (3)	0.0606 (10)
O4	1.0018 (2)	0.8339 (3)	0.3421 (3)	0.0693 (11)
O2	0.9808 (3)	0.7045 (5)	0.2204 (3)	0.0964 (18)
N3	0.2001 (2)	0.3971 (3)	0.1739 (3)	0.0344 (8)
N5	0.3884 (2)	0.5240 (3)	0.3241 (3)	0.0313 (8)
N6	0.3205 (2)	0.5777 (3)	0.4687 (3)	0.0324 (8)
N2	0.2024 (2)	0.6272 (3)	0.2292 (3)	0.0343 (8)
N7	0.5158 (2)	0.6289 (3)	0.3975 (3)	0.0370 (9)
H7	0.5562	0.6742	0.4414	0.044*

N4	0.1133 (2)	0.6929 (3)	0.0713 (3)	0.0430 (9)
H4	0.0801	0.6916	0.0054	0.052*
N1	1.0157 (2)	0.7334 (4)	0.3117 (3)	0.0447 (10)
C8	0.4375 (3)	0.5129 (4)	0.2623 (3)	0.0343 (10)
C15	0.1532 (3)	0.4763 (4)	0.0953 (4)	0.0368 (10)
C20	0.5192 (3)	0.5792 (4)	0.3096 (4)	0.0371 (10)
C7	0.1553 (3)	0.5995 (4)	0.1300 (3)	0.0339 (9)
C19	0.4372 (2)	0.5941 (3)	0.4029 (3)	0.0321 (9)
C17	0.4038 (2)	0.6273 (3)	0.4827 (3)	0.0327 (9)
C6	0.1899 (3)	0.7514 (4)	0.2356 (4)	0.0388 (10)
C13	0.1927 (3)	0.2884 (4)	0.1364 (4)	0.0419 (11)
H13	0.2196	0.2229	0.1783	0.050*
C5	0.1327 (3)	0.7920 (4)	0.1353 (4)	0.0441 (11)
C18	0.4377 (3)	0.7027 (4)	0.5649 (4)	0.0421 (11)
H18	0.4923	0.7435	0.5842	0.051*
C11	0.5629 (4)	0.5228 (5)	0.1731 (5)	0.0571 (14)
H11	0.6040	0.5248	0.1412	0.069*
C1	0.2222 (3)	0.8290 (4)	0.3202 (4)	0.0495 (12)
H1	0.2602	0.8024	0.3862	0.059*
C12	0.5820 (3)	0.5849 (5)	0.2636 (5)	0.0540 (14)
H12	0.6349	0.6294	0.2935	0.065*
C14	0.1106 (3)	0.4261 (5)	-0.0009 (4)	0.0467 (12)
H14	0.0759	0.4676	-0.0613	0.056*
C10	0.4830 (4)	0.4559 (5)	0.1265 (4)	0.0507 (12)
H10	0.4723	0.4144	0.0649	0.061*
C9	0.4209 (3)	0.4511 (4)	0.1705 (4)	0.0443 (11)
H9	0.3681	0.4069	0.1392	0.053*
C16	0.2942 (3)	0.6140 (4)	0.5385 (4)	0.0412 (11)
H16	0.2400	0.5892	0.5412	0.049*
C4	0.1057 (4)	0.9106 (5)	0.1180 (5)	0.0562 (14)
H4A	0.0679	0.9375	0.0520	0.067*
C2	0.1958 (4)	0.9471 (4)	0.3023 (5)	0.0612 (15)
H2	0.2169	1.0019	0.3569	0.073*
C3	0.1367 (4)	0.9856 (4)	0.2012 (6)	0.0661 (18)
H3	0.1186	1.0652	0.1916	0.079*
H1A	0.1016	0.5069	0.3358	0.079*
H1B	0.1000	0.4040	0.3241	0.079*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1	0.0293 (3)	0.0361 (3)	0.0344 (4)	-0.0022 (2)	0.0065 (3)	-0.0048 (2)
Cl1	0.0429 (6)	0.0411 (6)	0.0506 (8)	0.0038 (5)	0.0053 (5)	0.0024 (5)
S1	0.0636 (8)	0.0451 (6)	0.0402 (8)	-0.0015 (5)	0.0170 (6)	-0.0142 (5)
S2	0.0516 (7)	0.0592 (8)	0.0476 (8)	-0.0047 (6)	0.0119 (6)	-0.0238 (6)
O1	0.0344 (15)	0.0363 (15)	0.051 (2)	-0.0037 (12)	0.0148 (14)	-0.0003 (14)
O3	0.072 (2)	0.054 (2)	0.046 (2)	0.0172 (18)	0.0134 (19)	0.0004 (17)
O4	0.059 (2)	0.057 (2)	0.093 (3)	0.0218 (18)	0.032 (2)	0.011 (2)

O2	0.090 (3)	0.159 (5)	0.025 (2)	-0.052 (3)	0.006 (2)	-0.010 (2)
N3	0.0294 (16)	0.0339 (17)	0.034 (2)	-0.0017 (14)	0.0065 (15)	-0.0072 (15)
N5	0.0255 (16)	0.0344 (17)	0.029 (2)	-0.0029 (13)	0.0058 (15)	-0.0020 (15)
N6	0.0306 (17)	0.0380 (18)	0.025 (2)	0.0000 (14)	0.0077 (15)	-0.0059 (15)
N2	0.0346 (17)	0.0303 (17)	0.035 (2)	-0.0006 (14)	0.0104 (16)	-0.0017 (15)
N7	0.0291 (16)	0.0394 (18)	0.036 (2)	-0.0067 (14)	0.0055 (16)	0.0033 (16)
N4	0.042 (2)	0.045 (2)	0.036 (2)	0.0034 (17)	0.0087 (17)	0.0068 (18)
N1	0.0357 (19)	0.061 (3)	0.038 (2)	-0.0123 (18)	0.0146 (18)	0.001 (2)
C8	0.032 (2)	0.035 (2)	0.033 (2)	0.0016 (17)	0.0102 (19)	0.0018 (18)
C15	0.032 (2)	0.045 (2)	0.032 (3)	-0.0053 (17)	0.0113 (19)	-0.0058 (19)
C20	0.034 (2)	0.034 (2)	0.044 (3)	0.0039 (17)	0.0153 (19)	0.010 (2)
C7	0.0296 (19)	0.040 (2)	0.033 (2)	0.0007 (17)	0.0126 (18)	0.0016 (18)
C19	0.0265 (18)	0.033 (2)	0.033 (2)	0.0024 (16)	0.0076 (18)	0.0031 (18)
C17	0.0296 (19)	0.0275 (18)	0.032 (2)	-0.0016 (16)	0.0032 (17)	0.0002 (17)
C6	0.033 (2)	0.030 (2)	0.054 (3)	-0.0005 (17)	0.017 (2)	0.004 (2)
C13	0.031 (2)	0.044 (2)	0.046 (3)	-0.0004 (18)	0.010 (2)	-0.009 (2)
C5	0.039 (2)	0.045 (3)	0.054 (3)	0.0063 (19)	0.024 (2)	0.018 (2)
C18	0.042 (2)	0.037 (2)	0.040 (3)	-0.0067 (18)	0.007 (2)	-0.0074 (19)
C11	0.069 (3)	0.057 (3)	0.062 (4)	0.013 (3)	0.043 (3)	0.017 (3)
C1	0.048 (3)	0.035 (2)	0.065 (3)	0.001 (2)	0.021 (2)	-0.003 (2)
C12	0.040 (2)	0.053 (3)	0.071 (4)	0.003 (2)	0.024 (3)	0.023 (3)
C14	0.045 (2)	0.057 (3)	0.033 (3)	-0.002 (2)	0.010 (2)	0.001 (2)
C10	0.064 (3)	0.055 (3)	0.040 (3)	0.004 (2)	0.028 (3)	0.006 (2)
C9	0.046 (3)	0.043 (2)	0.043 (3)	0.001 (2)	0.018 (2)	-0.001 (2)
C16	0.036 (2)	0.046 (2)	0.038 (3)	-0.0030 (19)	0.012 (2)	-0.004 (2)
C4	0.060 (3)	0.045 (3)	0.071 (4)	0.016 (2)	0.035 (3)	0.022 (3)
C2	0.069 (3)	0.038 (2)	0.092 (5)	-0.002 (2)	0.048 (3)	-0.005 (3)
C3	0.067 (3)	0.034 (2)	0.116 (6)	0.011 (2)	0.057 (4)	0.013 (3)

Geometric parameters (Å, °)

Ni1—O1	2.109 (3)	C8—C9	1.383 (6)
Ni1—N3	2.133 (3)	C8—C20	1.424 (6)
Ni1—N5	2.134 (3)	C15—C14	1.361 (6)
Ni1—N2	2.159 (3)	C15—C7	1.456 (6)
Ni1—N6	2.171 (3)	C20—C12	1.396 (7)
Ni1—C11	2.4772 (13)	C19—C17	1.461 (6)
S1—C18	1.704 (5)	C17—C18	1.351 (6)
S1—C16	1.721 (4)	C6—C1	1.389 (7)
S2—C13	1.700 (5)	C6—C5	1.410 (6)
S2—C14	1.713 (5)	C13—H13	0.9300
O1—H1A	0.8372	C5—C4	1.387 (7)
O1—H1B	0.7947	C18—H18	0.9300
O3—N1	1.244 (5)	C11—C12	1.364 (8)
O4—N1	1.252 (5)	C11—C10	1.406 (8)
O2—N1	1.214 (5)	C11—H11	0.9300
N3—C13	1.310 (5)	C1—C2	1.381 (7)
N3—C15	1.380 (5)	C1—H1	0.9300

N5—C19	1.326 (5)	C12—H12	0.9300
N5—C8	1.386 (6)	C14—H14	0.9300
N6—C16	1.271 (6)	C10—C9	1.364 (7)
N6—C17	1.391 (5)	C10—H10	0.9300
N2—C7	1.324 (5)	C9—H9	0.9300
N2—C6	1.412 (5)	C16—H16	0.9300
N7—C19	1.354 (5)	C4—C3	1.356 (8)
N7—C20	1.367 (6)	C4—H4A	0.9300
N7—H7	0.8600	C2—C3	1.420 (8)
N4—C7	1.335 (5)	C2—H2	0.9300
N4—C5	1.379 (6)	C3—H3	0.9300
N4—H4	0.8600		
O1—Ni1—N3	90.37 (13)	N2—C7—C15	119.8 (4)
O1—Ni1—N5	169.10 (12)	N4—C7—C15	126.7 (4)
N3—Ni1—N5	99.19 (13)	N5—C19—N7	112.6 (4)
O1—Ni1—N2	88.88 (12)	N5—C19—C17	120.3 (3)
N3—Ni1—N2	77.42 (13)	N7—C19—C17	127.0 (4)
N5—Ni1—N2	88.12 (13)	C18—C17—N6	114.4 (4)
O1—Ni1—N6	91.69 (12)	C18—C17—C19	131.0 (4)
N3—Ni1—N6	171.22 (14)	N6—C17—C19	114.5 (3)
N5—Ni1—N6	78.08 (13)	C1—C6—C5	121.2 (4)
N2—Ni1—N6	94.09 (13)	C1—C6—N2	130.8 (4)
O1—Ni1—C11	91.44 (9)	C5—C6—N2	107.9 (4)
N3—Ni1—C11	91.99 (10)	N3—C13—S2	115.4 (4)
N5—Ni1—C11	93.43 (9)	N3—C13—H13	122.3
N2—Ni1—C11	169.41 (10)	S2—C13—H13	122.3
N6—Ni1—C11	96.48 (10)	N4—C5—C4	132.9 (5)
C18—S1—C16	89.2 (2)	N4—C5—C6	105.9 (4)
C13—S2—C14	89.5 (2)	C4—C5—C6	121.1 (5)
Ni1—O1—H1A	122.0	C17—C18—S1	110.3 (3)
Ni1—O1—H1B	130.8	C17—C18—H18	124.9
H1A—O1—H1B	90.7	S1—C18—H18	124.9
C13—N3—C15	110.2 (4)	C12—C11—C10	121.9 (5)
C13—N3—Ni1	134.7 (3)	C12—C11—H11	119.0
C15—N3—Ni1	115.1 (3)	C10—C11—H11	119.0
C19—N5—C8	105.7 (3)	C2—C1—C6	117.4 (5)
C19—N5—Ni1	113.7 (3)	C2—C1—H1	121.3
C8—N5—Ni1	140.4 (3)	C6—C1—H1	121.3
C16—N6—C17	111.0 (4)	C11—C12—C20	117.5 (5)
C16—N6—Ni1	135.5 (3)	C11—C12—H12	121.3
C17—N6—Ni1	113.2 (3)	C20—C12—H12	121.3
C7—N2—C6	105.0 (3)	C15—C14—S2	110.0 (4)
C7—N2—Ni1	113.3 (3)	C15—C14—H14	125.0
C6—N2—Ni1	141.4 (3)	S2—C14—H14	125.0
C19—N7—C20	107.6 (3)	C9—C10—C11	120.8 (5)
C19—N7—H7	126.2	C9—C10—H10	119.6
C20—N7—H7	126.2	C11—C10—H10	119.6

C7—N4—C5	107.6 (4)	C10—C9—C8	119.2 (5)
C7—N4—H4	126.2	C10—C9—H9	120.4
C5—N4—H4	126.2	C8—C9—H9	120.4
O2—N1—O3	120.7 (5)	N6—C16—S1	115.2 (3)
O2—N1—O4	121.6 (5)	N6—C16—H16	122.4
O3—N1—O4	117.7 (4)	S1—C16—H16	122.4
C9—C8—N5	132.1 (4)	C3—C4—C5	117.4 (5)
C9—C8—C20	119.5 (4)	C3—C4—H4A	121.3
N5—C8—C20	108.4 (4)	C5—C4—H4A	121.3
C14—C15—N3	114.9 (4)	C1—C2—C3	120.5 (5)
C14—C15—C7	130.8 (4)	C1—C2—H2	119.8
N3—C15—C7	114.3 (4)	C3—C2—H2	119.8
N7—C20—C12	133.3 (4)	C4—C3—C2	122.4 (5)
N7—C20—C8	105.7 (4)	C4—C3—H3	118.8
C12—C20—C8	121.0 (5)	C2—C3—H3	118.8
N2—C7—N4	113.5 (4)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1—H1A \cdots O3 ⁱ	0.84	2.00	2.751 (5)	149
O1—H1B \cdots O2 ⁱⁱ	0.79	2.53	3.293 (6)	160
O1—H1B \cdots O4 ⁱⁱ	0.79	2.38	3.025 (5)	139
N4—H4 \cdots O3 ⁱⁱⁱ	0.86	2.35	2.962 (5)	129
N4—H4 \cdots O4 ⁱⁱⁱ	0.86	2.14	2.996 (5)	173
N7—H7 \cdots Cl1 ^{iv}	0.86	2.35	3.159 (4)	157

Symmetry codes: (i) $x-1, y, z$; (ii) $-x+1, y-1/2, -z+1/2$; (iii) $x-1, -y+3/2, z-1/2$; (iv) $-x+1, -y+1, -z+1$.