

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 2,2'-[(4-Ethoxyphenyl)methylene]bis(3-hydroxy-5,5-dimethylcyclohex-2-en-1-one)

N. Sureshbabu and V. Sughanya\*

Department of Chemistry, Annamalai University, Annamalai Nagar 608 002, Tamil Nadu, India

Correspondence e-mail: saisukanyashri@gmail.com


Received 24 July 2012; accepted 29 July 2012

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.042;  $wR$  factor = 0.122; data-to-parameter ratio = 14.6.

In the title compound,  $\text{C}_{25}\text{H}_{32}\text{O}_5$ , the two cyclohexenone rings have envelope conformations with the C atom bearing two methyl groups as the flap atom in each ring. Relatively strong intramolecular  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bonds are observed.

### Related literature

For the synthesis of bisdimedones, see: Vang & Stankevich (1960); Hilderbrand & Weissleder (2007). For their pharmaceutical properties, see: Lambert *et al.* (1997); Poupelin *et al.* (1978); Hideo (1981); Selvanayagam *et al.* (1996); Jonathan *et al.* (1988). For the crystal structures of related xanthene derivatives, see: Odabaşoğlu *et al.* (2008); Mehdi *et al.* (2011); Ravikumar *et al.* (2012). For the assignment of ring conformations, see: Cremer & Pople (1975).



### Experimental

#### Crystal data

$\text{C}_{25}\text{H}_{32}\text{O}_5$   
 $M_r = 412.51$   
 Monoclinic,  $P2_1/n$   
 $a = 9.774$  (5) Å  
 $b = 10.698$  (5) Å

$c = 21.578$  (5) Å  
 $\beta = 93.735$  (5)°  
 $V = 2251.5$  (16) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation

$\mu = 0.08$  mm<sup>-1</sup>  
 $T = 293$  K

0.30 × 0.20 × 0.20 mm

#### Data collection

Bruker Kappa APEXII CCD  
 diffractometer  
 Absorption correction: multi-scan  
 (SADABS; Bruker, 2004)  
 $T_{\min} = 0.961$ ,  $T_{\max} = 0.989$

17055 measured reflections  
 3962 independent reflections  
 3065 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.028$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$   
 $wR(F^2) = 0.122$   
 $S = 1.03$   
 3962 reflections

272 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.24$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.23$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O3}-\text{H3A}\cdots\text{O5}$	0.82	1.83	2.631 (2)	164
$\text{O4}-\text{H4A}\cdots\text{O2}$	0.82	1.78	2.5864 (19)	167

Data collection: APEX2 (Bruker, 2004); cell refinement: APEX2 and SAINT (Bruker, 2004); data reduction: SAINT and XPREP (Bruker, 2004); program(s) used to solve structure: SIR92 (Altomare *et al.*, 1993); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and Mercury (Macrae *et al.*, 2008); software used to prepare material for publication: SHELXL97.

The authors thank Dr Babu Varghese and the SAIF, IIT Madras, for the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WN2485).

### References

- Altomare, A., Casciarano, G., Giacovazzo, C. & Guagliardi, A. (1993). *J. Appl. Cryst.* **26**, 343–350.
- Bruker (2004). *SADABS, APEX2, SAINT and XPREP*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Hideo, T. (1981). Jpn Kokai Tokkyo Koho JP 56 005480.
- Hilderbrand, S. A. & Weissleder, R. (2007). *Tetrahedron Lett.* **48**, 4383–4385.
- Jonathan, R. D., Srinivas, K. R. & Glen, E. B. (1988). *Eur. J. Med. Chem.* **23**, 111–117.
- Lambert, R. W., Martin, J. A., Merrett, J. H., Parkes, K. E. B. & Thomas, G. J. (1997). PCT Int. Appl. WO 9706178.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Mehdi, S. H., Hashim, R., Ghalib, R. M., Yeap, C. S. & Fun, H.-K. (2011). *Acta Cryst.* **E67**, o1449.
- Odabaşoğlu, M., Kaya, M., Yıldırım, Y. & Büyükgüngör, O. (2008). *Acta Cryst.* **E64**, o681.
- Poupelin, J. P., Rut, G. S., Blanpin, O. F., Narcisse, G., Ernouf, G. U. & Lacroise, R. (1978). *Eur. J. Med. Chem.* **13**, 67–71.
- Ravikumar, N., Gopikrishna, G. & Solomon, K. A. (2012). *Acta Cryst.* **E68**, o265.
- Selvanayagam, Z. E., Gnanavendhan, S. G., Balakrishnan, K., Rao, R. B., Sivaraman, R. E. & Subramanian, K. (1996). *J. Nat. Prod.* **59**, 664–667.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Vang, G. Y. & Stankevich, E. L. (1960). *Zh. Obshch. Khim.* **30**, 3287.

## supporting information

*Acta Cryst.* (2012). E68, o2638 [doi:10.1107/S1600536812033934]

## 2,2'-[(4-Ethoxyphenyl)methylene]bis(3-hydroxy-5,5-dimethylcyclohex-2-en-1-one)

N. Sureshbabu and V. Sughanya

### S1. Comment

Several methods have been reported in the literature (Vang & Stankevich, 1960; Hilderbrand & Weissleder, 2007) for the synthesis of the title compound. Xanthene derivatives possess biological properties such as antibacterial, antiviral and anti-inflammatory activities (Jonathan *et al.*, 1988) and are therefore used in medicine.


In the title compound, the cyclohexenone rings C10–C15 and C18–C23 both adopt envelope conformations, with flap atoms C13 and C21, respectively. The dihedral angle between the two cyclohexenone planes Q(C10/C11/C12/C14/C15) and R(C18/C19/C20/C22/C23) is 58.42 (3)°. The dihedral angle between the benzene ring P(C3–C8) and cyclohexenone planes Q and R are 66.37 (3)° and 61.41 (4)°, respectively. The hydroxy and carbonyl oxygen atoms face each other and are oriented to allow for the formation of two intramolecular O—H···O hydrogen bonds (Table 1, Fig. 2), typical for xanthene derivatives.

### S2. Experimental


The title compound was prepared in a single stage. A mixture of 4-Ethoxybenzaldehyde (1.2g, 8 mmol), 5,5-dimethylcyclohexane-1,3-dione (2.24g, 16 mmol) and 20ml of ethanol was heated to 70 °C for about 10 minutes. The reaction mixture was allowed to cool to room temperature and the resulting title compound, 2,2'-((4-ethoxyphenyl)methylene) bis-(3-hydroxy-5,5-dimethylcyclohex-2-enone) was filtered and dried (M.p. 411 K; Yield 78% ).

### S3. Refinement

All hydrogen atoms were identified from difference electron density peaks and subsequently treated as riding atoms with  $d(\text{Csp}^2\text{—H}) = 0.93 \text{ \AA}$ ,  $d(\text{Cmethyl—H}) = 0.96 \text{ \AA}$ ,  $d(\text{Cmethylene—H}) = 0.97 \text{ \AA}$ ,  $d(\text{Cmethine—H}) = 0.98 \text{ \AA}$ ;  $d(\text{O—H}) = 0.82 \text{ \AA}$ ;  $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C}, \text{O})$ , where  $x = 1.5$  for methyl H and 1.2 for all other H atoms.

**Figure 1**

A view of the structure of the title compound. Displacement ellipsoids are drawn at the 50% probability level. Hydrogen atoms are shown as spheres of arbitrary radius.

**Figure 2**

A view of the packing in the crystal structure, showing intramolecular O—H...O hydrogen bonds as dotted lines.

### 2,2'-[(4-Ethoxyphenyl)methylene]bis(3-hydroxy-5,5-dimethylcyclohex-2-en-1-one)

#### Crystal data

$C_{25}H_{32}O_5$

$M_r = 412.51$

Monoclinic,  $P2_1/n$

$a = 9.774 (5) \text{ \AA}$

$b = 10.698 (5) \text{ \AA}$

$c = 21.578 (5) \text{ \AA}$

$\beta = 93.735 (5)^\circ$   
 $V = 2251.5 (16) \text{ \AA}^3$   
 $Z = 4$   
 $F(000) = 888$   
 $D_x = 1.217 \text{ Mg m}^{-3}$   
 Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 6149 reflections  
 $\theta = 2.1\text{--}27.1^\circ$   
 $\mu = 0.08 \text{ mm}^{-1}$   
 $T = 293 \text{ K}$   
 Block, colourless  
 $0.30 \times 0.20 \times 0.20 \text{ mm}$

*Data collection*

Bruker Kappa APEXII CCD  
 diffractometer  
 Radiation source: fine-focus sealed tube  
 Graphite monochromator  
 $\omega$  and  $\varphi$  scan  
 Absorption correction: multi-scan  
 (SADABS; Bruker, 2005)  
 $T_{\min} = 0.961$ ,  $T_{\max} = 0.989$

17055 measured reflections  
 3962 independent reflections  
 3065 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.028$   
 $\theta_{\max} = 25.0^\circ$ ,  $\theta_{\min} = 2.1^\circ$   
 $h = -11 \rightarrow 7$   
 $k = -12 \rightarrow 12$   
 $l = -24 \rightarrow 25$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.042$   
 $wR(F^2) = 0.122$   
 $S = 1.03$   
 3962 reflections  
 272 parameters  
 0 restraints  
 Primary atom site location: structure-invariant  
 direct methods  
 Secondary atom site location: difference Fourier  
 map

Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0573P)^2 + 0.7618P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.24 \text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.23 \text{ e \AA}^{-3}$   
 Extinction correction: SHELXL97 (Sheldrick,  
 2008),  $F_c^* = kFc[1 + 0.001x\lambda^3/\sin(2\theta)]^{-1/4}$   
 Extinction coefficient: 0.0143 (14)

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.1136 (3)	0.5154 (3)	0.33826 (11)	0.0776 (7)
H1A	0.1550	0.5226	0.2993	0.116*
H1B	0.0836	0.5964	0.3510	0.116*
H1C	0.0363	0.4600	0.3337	0.116*
C2	0.2151 (2)	0.4651 (3)	0.38572 (10)	0.0707 (7)
H2A	0.2457	0.3831	0.3732	0.085*
H2B	0.2943	0.5198	0.3900	0.085*
C3	0.22806 (19)	0.40529 (19)	0.49280 (8)	0.0495 (5)
C4	0.3642 (2)	0.3757 (2)	0.49343 (9)	0.0567 (6)

---

H4	0.4129	0.3911	0.4586	0.068*
C5	0.42951 (19)	0.32300 (19)	0.54571 (9)	0.0518 (5)
H5	0.5221	0.3032	0.5452	0.062*
C6	0.36275 (17)	0.29861 (16)	0.59866 (7)	0.0368 (4)
C7	0.22621 (18)	0.33226 (18)	0.59738 (8)	0.0474 (5)
H7	0.1781	0.3191	0.6326	0.057*
C8	0.15904 (19)	0.3846 (2)	0.54565 (9)	0.0547 (5)
H8	0.0669	0.4061	0.5463	0.066*
C9	0.44542 (16)	0.24953 (15)	0.65658 (7)	0.0350 (4)
H9	0.4980	0.3224	0.6720	0.042*
C10	0.55496 (16)	0.15470 (16)	0.64172 (7)	0.0359 (4)
C11	0.52716 (17)	0.05495 (16)	0.60035 (7)	0.0373 (4)
C12	0.63981 (18)	-0.02547 (18)	0.57850 (8)	0.0474 (5)
H12A	0.6192	-0.0451	0.5350	0.057*
H12B	0.6408	-0.1035	0.6014	0.057*
C13	0.78274 (18)	0.0323 (2)	0.58573 (8)	0.0497 (5)
C14	0.80122 (19)	0.0814 (2)	0.65212 (9)	0.0564 (5)
H14A	0.8101	0.0108	0.6803	0.068*
H14B	0.8858	0.1288	0.6566	0.068*
C15	0.68583 (18)	0.16245 (18)	0.67054 (8)	0.0433 (4)
C16	0.7976 (2)	0.1388 (2)	0.53921 (10)	0.0651 (6)
H16A	0.8883	0.1733	0.5446	0.098*
H16B	0.7314	0.2027	0.5462	0.098*
H16C	0.7825	0.1071	0.4977	0.098*
C17	0.8898 (2)	-0.0678 (2)	0.57416 (11)	0.0714 (7)
H17A	0.8820	-0.1349	0.6033	0.107*
H17B	0.9798	-0.0319	0.5793	0.107*
H17C	0.8746	-0.0994	0.5326	0.107*
C18	0.36132 (16)	0.21294 (15)	0.71075 (7)	0.0341 (4)
C19	0.25324 (17)	0.12859 (16)	0.70621 (8)	0.0373 (4)
C20	0.1721 (2)	0.09839 (17)	0.76079 (9)	0.0472 (5)
H20A	0.2115	0.0251	0.7815	0.057*
H20B	0.0790	0.0777	0.7459	0.057*
C21	0.16771 (18)	0.20449 (17)	0.80774 (8)	0.0434 (4)
C22	0.31497 (19)	0.24452 (19)	0.82458 (8)	0.0457 (5)
H22A	0.3135	0.3233	0.8468	0.055*
H22B	0.3576	0.1829	0.8526	0.055*
C23	0.40218 (18)	0.25987 (16)	0.77023 (8)	0.0394 (4)
C24	0.08261 (19)	0.31254 (19)	0.78021 (9)	0.0516 (5)
H24A	0.0811	0.3791	0.8100	0.077*
H24B	-0.0093	0.2845	0.7698	0.077*
H24C	0.1224	0.3422	0.7434	0.077*
C25	0.1019 (2)	0.1593 (2)	0.86609 (10)	0.0667 (6)
H25A	0.1000	0.2266	0.8954	0.100*
H25B	0.1543	0.0912	0.8842	0.100*
H25C	0.0099	0.1318	0.8552	0.100*
O1	0.15317 (15)	0.45642 (16)	0.44304 (6)	0.0703 (5)
O2	0.40539 (12)	0.02575 (12)	0.58061 (5)	0.0467 (3)

O3	0.71819 (13)	0.23817 (14)	0.71624 (6)	0.0597 (4)
H3A	0.6481	0.2663	0.7299	0.072*
O4	0.21555 (12)	0.06855 (12)	0.65595 (6)	0.0477 (3)
H4A	0.2793	0.0663	0.6331	0.057*
O5	0.51622 (13)	0.31629 (13)	0.78119 (6)	0.0527 (4)

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0780 (16)	0.0951 (18)	0.0589 (14)	−0.0061 (14)	−0.0029 (12)	0.0242 (13)
C2	0.0646 (14)	0.0954 (18)	0.0529 (13)	0.0138 (12)	0.0107 (11)	0.0272 (13)
C3	0.0478 (11)	0.0603 (12)	0.0411 (10)	0.0145 (9)	0.0076 (8)	0.0138 (9)
C4	0.0508 (12)	0.0773 (14)	0.0440 (11)	0.0176 (10)	0.0187 (9)	0.0229 (10)
C5	0.0379 (10)	0.0693 (13)	0.0497 (11)	0.0141 (9)	0.0138 (8)	0.0178 (10)
C6	0.0370 (9)	0.0376 (9)	0.0365 (9)	0.0018 (7)	0.0066 (7)	0.0024 (7)
C7	0.0406 (10)	0.0626 (12)	0.0405 (10)	0.0109 (9)	0.0132 (8)	0.0102 (9)
C8	0.0390 (10)	0.0751 (14)	0.0510 (11)	0.0181 (9)	0.0117 (8)	0.0158 (10)
C9	0.0330 (8)	0.0373 (9)	0.0349 (9)	−0.0032 (7)	0.0035 (7)	−0.0011 (7)
C10	0.0345 (9)	0.0440 (10)	0.0294 (8)	0.0006 (7)	0.0034 (7)	0.0028 (7)
C11	0.0385 (9)	0.0458 (10)	0.0278 (8)	0.0013 (7)	0.0036 (7)	0.0048 (7)
C12	0.0498 (11)	0.0536 (11)	0.0393 (10)	0.0099 (9)	0.0070 (8)	−0.0028 (9)
C13	0.0400 (10)	0.0690 (13)	0.0404 (10)	0.0151 (9)	0.0056 (8)	0.0012 (9)
C14	0.0379 (10)	0.0842 (15)	0.0464 (11)	0.0132 (10)	−0.0034 (8)	−0.0067 (11)
C15	0.0369 (9)	0.0597 (11)	0.0334 (9)	−0.0015 (8)	0.0019 (7)	−0.0024 (9)
C16	0.0528 (13)	0.0821 (16)	0.0623 (13)	0.0027 (11)	0.0171 (10)	0.0066 (12)
C17	0.0564 (13)	0.0975 (18)	0.0607 (13)	0.0304 (12)	0.0077 (10)	−0.0055 (13)
C18	0.0346 (9)	0.0354 (9)	0.0325 (9)	0.0014 (7)	0.0043 (7)	−0.0005 (7)
C19	0.0379 (9)	0.0359 (9)	0.0384 (9)	0.0019 (7)	0.0056 (7)	−0.0020 (8)
C20	0.0492 (11)	0.0432 (10)	0.0509 (11)	−0.0054 (8)	0.0168 (8)	0.0013 (9)
C21	0.0460 (10)	0.0459 (10)	0.0397 (10)	0.0004 (8)	0.0142 (8)	0.0002 (8)
C22	0.0494 (11)	0.0536 (11)	0.0347 (10)	0.0085 (9)	0.0067 (8)	−0.0011 (8)
C23	0.0386 (9)	0.0394 (9)	0.0405 (10)	0.0030 (8)	0.0048 (7)	−0.0019 (8)
C24	0.0437 (11)	0.0551 (12)	0.0569 (12)	0.0061 (9)	0.0096 (9)	−0.0031 (10)
C25	0.0746 (15)	0.0732 (15)	0.0559 (13)	−0.0010 (12)	0.0305 (11)	0.0050 (11)
O1	0.0564 (9)	0.1055 (13)	0.0500 (8)	0.0307 (8)	0.0115 (7)	0.0312 (8)
O2	0.0411 (7)	0.0590 (8)	0.0398 (7)	−0.0045 (6)	0.0008 (5)	−0.0075 (6)
O3	0.0373 (7)	0.0860 (10)	0.0556 (8)	−0.0051 (7)	0.0016 (6)	−0.0234 (8)
O4	0.0441 (7)	0.0527 (8)	0.0472 (7)	−0.0104 (6)	0.0099 (6)	−0.0117 (6)
O5	0.0459 (8)	0.0660 (9)	0.0460 (7)	−0.0096 (6)	0.0029 (6)	−0.0164 (7)

*Geometric parameters (Å, °)*

C1—C2	1.481 (3)	C14—C15	1.497 (3)
C1—H1A	0.9600	C14—H14A	0.9700
C1—H1B	0.9600	C14—H14B	0.9700
C1—H1C	0.9600	C15—O3	1.299 (2)
C2—O1	1.415 (2)	C16—H16A	0.9600
C2—H2A	0.9700	C16—H16B	0.9600

C2—H2B	0.9700	C16—H16C	0.9600
C3—C4	1.367 (3)	C17—H17A	0.9600
C3—O1	1.373 (2)	C17—H17B	0.9600
C3—C8	1.380 (3)	C17—H17C	0.9600
C4—C5	1.380 (3)	C18—C19	1.388 (2)
C4—H4	0.9300	C18—C23	1.411 (2)
C5—C6	1.377 (2)	C19—O4	1.293 (2)
C5—H5	0.9300	C19—C20	1.497 (2)
C6—C7	1.381 (2)	C20—C21	1.524 (3)
C6—C9	1.536 (2)	C20—H20A	0.9700
C7—C8	1.377 (3)	C20—H20B	0.9700
C7—H7	0.9300	C21—C24	1.522 (3)
C8—H8	0.9300	C21—C22	1.523 (3)
C9—C18	1.523 (2)	C21—C25	1.529 (3)
C9—C10	1.524 (2)	C22—C23	1.503 (2)
C9—H9	0.9800	C22—H22A	0.9700
C10—C15	1.388 (2)	C22—H22B	0.9700
C10—C11	1.406 (2)	C23—O5	1.276 (2)
C11—O2	1.276 (2)	C24—H24A	0.9600
C11—C12	1.497 (2)	C24—H24B	0.9600
C12—C13	1.526 (3)	C24—H24C	0.9600
C12—H12A	0.9700	C25—H25A	0.9600
C12—H12B	0.9700	C25—H25B	0.9600
C13—C14	1.525 (3)	C25—H25C	0.9600
C13—C17	1.529 (3)	O3—H3A	0.8200
C13—C16	1.531 (3)	O4—H4A	0.8200
C2—C1—H1A	109.5	H14A—C14—H14B	107.7
C2—C1—H1B	109.5	O3—C15—C10	123.30 (17)
H1A—C1—H1B	109.5	O3—C15—C14	114.36 (15)
C2—C1—H1C	109.5	C10—C15—C14	122.33 (16)
H1A—C1—H1C	109.5	C13—C16—H16A	109.5
H1B—C1—H1C	109.5	C13—C16—H16B	109.5
O1—C2—C1	108.82 (19)	H16A—C16—H16B	109.5
O1—C2—H2A	109.9	C13—C16—H16C	109.5
C1—C2—H2A	109.9	H16A—C16—H16C	109.5
O1—C2—H2B	109.9	H16B—C16—H16C	109.5
C1—C2—H2B	109.9	C13—C17—H17A	109.5
H2A—C2—H2B	108.3	C13—C17—H17B	109.5
C4—C3—O1	124.68 (16)	H17A—C17—H17B	109.5
C4—C3—C8	118.85 (17)	C13—C17—H17C	109.5
O1—C3—C8	116.46 (16)	H17A—C17—H17C	109.5
C3—C4—C5	120.04 (17)	H17B—C17—H17C	109.5
C3—C4—H4	120.0	C19—C18—C23	117.68 (15)
C5—C4—H4	120.0	C19—C18—C9	124.00 (15)
C6—C5—C4	122.47 (17)	C23—C18—C9	118.13 (14)
C6—C5—H5	118.8	O4—C19—C18	123.79 (15)
C4—C5—H5	118.8	O4—C19—C20	114.57 (15)

C5—C6—C7	116.33 (16)	C18—C19—C20	121.63 (15)
C5—C6—C9	119.16 (15)	C19—C20—C21	113.67 (15)
C7—C6—C9	124.19 (15)	C19—C20—H20A	108.8
C8—C7—C6	122.12 (16)	C21—C20—H20A	108.8
C8—C7—H7	118.9	C19—C20—H20B	108.8
C6—C7—H7	118.9	C21—C20—H20B	108.8
C7—C8—C3	120.15 (17)	H20A—C20—H20B	107.7
C7—C8—H8	119.9	C24—C21—C22	111.33 (16)
C3—C8—H8	119.9	C24—C21—C20	110.15 (16)
C18—C9—C10	114.33 (14)	C22—C21—C20	107.52 (15)
C18—C9—C6	115.51 (13)	C24—C21—C25	108.22 (16)
C10—C9—C6	113.32 (13)	C22—C21—C25	109.52 (16)
C18—C9—H9	103.9	C20—C21—C25	110.09 (16)
C10—C9—H9	103.9	C23—C22—C21	114.81 (14)
C6—C9—H9	103.9	C23—C22—H22A	108.6
C15—C10—C11	117.65 (16)	C21—C22—H22A	108.6
C15—C10—C9	120.37 (15)	C23—C22—H22B	108.6
C11—C10—C9	121.95 (14)	C21—C22—H22B	108.6
O2—C11—C10	122.34 (15)	H22A—C22—H22B	107.5
O2—C11—C12	116.30 (16)	O5—C23—C18	121.93 (16)
C10—C11—C12	121.34 (15)	O5—C23—C22	116.10 (15)
C11—C12—C13	114.95 (16)	C18—C23—C22	121.96 (16)
C11—C12—H12A	108.5	C21—C24—H24A	109.5
C13—C12—H12A	108.5	C21—C24—H24B	109.5
C11—C12—H12B	108.5	H24A—C24—H24B	109.5
C13—C12—H12B	108.5	C21—C24—H24C	109.5
H12A—C12—H12B	107.5	H24A—C24—H24C	109.5
C14—C13—C12	106.67 (15)	H24B—C24—H24C	109.5
C14—C13—C17	110.72 (16)	C21—C25—H25A	109.5
C12—C13—C17	109.32 (18)	C21—C25—H25B	109.5
C14—C13—C16	110.46 (19)	H25A—C25—H25B	109.5
C12—C13—C16	111.08 (16)	C21—C25—H25C	109.5
C17—C13—C16	108.59 (17)	H25A—C25—H25C	109.5
C15—C14—C13	113.79 (15)	H25B—C25—H25C	109.5
C15—C14—H14A	108.8	C3—O1—C2	118.21 (15)
C13—C14—H14A	108.8	C15—O3—H3A	109.5
C15—C14—H14B	108.8	C19—O4—H4A	109.5
C13—C14—H14B	108.8		
O1—C3—C4—C5	-179.1 (2)	C9—C10—C15—O3	8.8 (3)
C8—C3—C4—C5	1.7 (3)	C11—C10—C15—C14	9.3 (3)
C3—C4—C5—C6	-0.3 (3)	C9—C10—C15—C14	-172.57 (17)
C4—C5—C6—C7	-1.3 (3)	C13—C14—C15—O3	-157.30 (18)
C4—C5—C6—C9	-175.17 (19)	C13—C14—C15—C10	24.0 (3)
C5—C6—C7—C8	1.5 (3)	C10—C9—C18—C19	-79.6 (2)
C9—C6—C7—C8	175.01 (18)	C6—C9—C18—C19	54.6 (2)
C6—C7—C8—C3	-0.1 (3)	C10—C9—C18—C23	95.19 (18)
C4—C3—C8—C7	-1.6 (3)	C6—C9—C18—C23	-130.60 (16)



O1—C3—C8—C7	179.2 (2)	C23—C18—C19—O4	-172.22 (15)
C5—C6—C9—C18	-174.17 (16)	C9—C18—C19—O4	2.6 (3)
C7—C6—C9—C18	12.5 (2)	C23—C18—C19—C20	6.6 (2)
C5—C6—C9—C10	-39.5 (2)	C9—C18—C19—C20	-178.58 (15)
C7—C6—C9—C10	147.15 (18)	O4—C19—C20—C21	-152.94 (16)
C18—C9—C10—C15	-89.34 (19)	C18—C19—C20—C21	28.2 (2)
C6—C9—C10—C15	135.45 (16)	C19—C20—C21—C24	69.2 (2)
C18—C9—C10—C11	88.71 (18)	C19—C20—C21—C22	-52.3 (2)
C6—C9—C10—C11	-46.5 (2)	C19—C20—C21—C25	-171.51 (17)
C15—C10—C11—O2	166.58 (16)	C24—C21—C22—C23	-75.1 (2)
C9—C10—C11—O2	-11.5 (2)	C20—C21—C22—C23	45.6 (2)
C15—C10—C11—C12	-11.6 (2)	C25—C21—C22—C23	165.22 (16)
C9—C10—C11—C12	170.32 (15)	C19—C18—C23—O5	165.19 (16)
O2—C11—C12—C13	162.12 (15)	C9—C18—C23—O5	-10.0 (2)
C10—C11—C12—C13	-19.6 (2)	C19—C18—C23—C22	-13.8 (2)
C11—C12—C13—C14	49.0 (2)	C9—C18—C23—C22	171.03 (15)
C11—C12—C13—C17	168.76 (16)	C21—C22—C23—O5	166.69 (16)
C11—C12—C13—C16	-71.4 (2)	C21—C22—C23—C18	-14.3 (2)
C12—C13—C14—C15	-50.8 (2)	C4—C3—O1—C2	8.4 (3)
C17—C13—C14—C15	-169.63 (19)	C8—C3—O1—C2	-172.5 (2)
C16—C13—C14—C15	70.0 (2)	C1—C2—O1—C3	176.6 (2)
C11—C10—C15—O3	-169.30 (16)		

*Hydrogen-bond geometry (Å, °)*

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
O3—H3A...O5	0.82	1.83	2.631 (2)	164
O4—H4A...O2	0.82	1.78	2.5864 (19)	167