

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N-(2-{[5-Bromo-2-(piperidin-1-yl)-pyrimidin-4-yl]sulfanyl}-4-methoxyphenyl)-4-methylbenzenesulfonamide

 Mohan Kumar,^a L. Mallesha,^b M. A. Sridhar,^{a*} Kamini Kapoor,^c Vivek K. Gupta^c and Rajni Kant^c

^aDepartment of Studies in Physics, Manasagangotri, University of Mysore, Mysore 570 006, India, ^bPG Department of Studies in Chemistry, JSS College of Arts, Commerce and Science, Ooty Road, Mysore 570 025, India, and ^cX-ray Crystallography Laboratory, Post-Graduate Department of Physics & Electronics, University of Jammu, Jammu Tawi 180 006, India
Correspondence e-mail: mas@physics.uni-mysore.ac.in


Received 28 August 2012; accepted 29 August 2012

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.037; wR factor = 0.093; data-to-parameter ratio = 14.6.

In the title compound, $\text{C}_{23}\text{H}_{25}\text{BrN}_4\text{O}_3\text{S}_2$, the benzene rings bridged by the sulfonamide group are tilted relative to each other by $69.7(1)^\circ$ and the dihedral angle between the sulfur-bridged pyrimidine and benzene rings is $70.4(1)^\circ$. The molecular conformation is stabilized by a weak intramolecular π - π stacking interaction between the pyrimidine and the 4-methyl benzene rings [centroid-centroid distance = $3.633(2)$ Å]. The piperidine ring adopts a chair conformation. In the crystal, molecules are linked into inversion dimers by pairs of $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

For a related structure and background to sulfonamides, see: Kant *et al.* (2012).



Experimental

Crystal data

$\text{C}_{23}\text{H}_{25}\text{BrN}_4\text{O}_3\text{S}_2$
 $M_r = 549.50$
 Triclinic, $P\bar{1}$
 $a = 9.8318(3)$ Å
 $b = 10.3822(3)$ Å
 $c = 13.4393(4)$ Å
 $\alpha = 96.654(3)^\circ$
 $\beta = 103.085(3)^\circ$
 $\gamma = 107.714(3)^\circ$
 $V = 1247.36(6)$ Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 1.85$ mm⁻¹
 $T = 293$ K
 $0.30 \times 0.20 \times 0.20$ mm

Data collection

Oxford Diffraction Xcalibur CCD, Sapphire3 diffractometer
 Absorption correction: multi-scan (*CrysAlis PRO*; Oxford Diffraction, 2010)
 $T_{\min} = 0.764$, $T_{\max} = 1.000$
 41580 measured reflections
 4385 independent reflections
 3645 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.038$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.093$
 $S = 1.03$
 4385 reflections
 300 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.66$ e Å⁻³
 $\Delta\rho_{\min} = -0.52$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N8}-\text{H8}\cdots\text{O2}^i$	0.86	2.22	2.955 (4)	143

 Symmetry code: (i) $-x, -y + 1, -z + 1$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *PLATON* (Spek, 2009).

MK acknowledges the help of Bahubali College of Engineering, Shravanabelagola for his research work and RK acknowledges the Department of Science & Technology for the single-crystal X-ray diffractometer sanctioned as a National Facility under project No. SR/S2/CMP-47/2003.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6947).

References

- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Kant, R., Gupta, V. K., Kapoor, K., Kumar, M., Mallesha, L. & Sridhar, M. A. (2012). *Acta Cryst.* **E68**, o2590–o2591.
 Oxford Diffraction (2010). *CrysAlis PRO*. Oxford Diffraction Ltd, Yarnton, England.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supporting information

Acta Cryst. (2012). E68, o2831 [https://doi.org/10.1107/S1600536812037257]

***N*-(2-[[5-Bromo-2-(piperidin-1-yl)pyrimidin-4-yl]sulfanyl]-4-methoxyphenyl)-4-methylbenzenesulfonamide**

Mohan Kumar, L. Mallesha, M. A. Sridhar, Kamini Kapoor, Vivek K. Gupta and Rajni Kant

S1. Comment

As part of our ongoing studies of sulfonamides (Kant *et al.*, 2012), we now report the structure of the title compound, (I), (Fig. 1).

The piperidine ring is exhibiting a chair conformation. The two benzene rings (C1—C6/C9—C14) are tilted relative to each other by 69.7 (1)° and the dihedral angle between the sulfur bridged pyrimidine and benzene rings is 70.4 (1)°. The molecular conformation is stabilized by a weak intramolecular stacking interaction between the pyrimidine and the 4-methyl benzene rings [centroid–centroid distance = 3.633 (2) Å, interplanar spacing = 3.494 Å, and centroid shift = 1.00 Å]. In the crystal, molecules are linked into inversion dimers by pairs of N8—H8···O2 hydrogen bonds (Fig.2).

S2. Experimental

The reaction of *N*-[2-(5-bromo-2-piperidin-1-yl-pyrimidin-4-ylsulfanyl)-4-methoxy-phenyl]-4-methyl-benzene-sulfonamide (5.01 g, 0.01 mol) and piperidine (0.86 g, 0.01 mol) were carried out in the presence of triethylamine and the reaction mixture was allowed to stir at room temperature for 6–7 h in dry dichloromethane. The progress of the reaction was monitored by TLC. Upon completion, the solvent was removed under reduced pressure and residue was extracted with ethyl acetate. The compound was purified by successive recrystallization from methanol (yield 83%, m. p. 483–485 K) to yield light brown blocks of (I).

S3. Refinement

All H atoms were positioned geometrically and were treated as riding on their parent C atoms, with C—H distances of 0.93–0.97 Å and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{methyl C})$.




Figure 1

The molecular structure of (I) with displacement ellipsoids drawn at the 40% probability level.




Figure 2

A unit-cell packing view of the title compound down the a axis, showing intermolecular interactions. For clarity, hydrogen atoms which are not involved in hydrogen bonding have been omitted.

N-(2-[[5-Bromo-2-(piperidin-1-yl)pyrimidin-4-yl]sulfanyl]-4-methoxyphenyl)-4-methylbenzenesulfonamide

Crystal data

$C_{23}H_{25}BrN_4O_3S_2$

$M_r = 549.50$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 9.8318$ (3) Å

$b = 10.3822$ (3) Å

$c = 13.4393$ (4) Å

$\alpha = 96.654$ (3)°

$\beta = 103.085$ (3)°

$\gamma = 107.714$ (3)°

$V = 1247.36$ (6) Å³

$Z = 2$

$F(000) = 564$

$D_x = 1.463$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 16775 reflections

$\theta = 3.4\text{--}29.1^\circ$

$\mu = 1.85$ mm⁻¹

$T = 293$ K

Block, light-brown

$0.30 \times 0.20 \times 0.20$ mm

Data collection

Oxford Diffraction Xcalibur CCD, Sapphire3 diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 16.1049 pixels mm⁻¹

ω scans

Absorption correction: multi-scan

(*CrysAlis PRO*; Oxford Diffraction, 2010)

$T_{\min} = 0.764$, $T_{\max} = 1.000$

41580 measured reflections

4385 independent reflections
 3645 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.038$
 $\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 3.4^\circ$

$h = -11 \rightarrow 11$
 $k = -12 \rightarrow 12$
 $l = -15 \rightarrow 15$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.093$
 $S = 1.03$
 4385 reflections
 300 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0396P)^2 + 0.9744P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.66 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.52 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. *CrysAlis PRO*, Oxford Diffraction Ltd., Version 1.171.34.40 (release 27-08-2010 CrysAlis171. NET) (compiled Aug 27 2010, 11:50:40) Empirical absorption correction using spherical harmonics, implemented in SCALE3 ABSPACK scaling algorithm.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.00669 (4)	0.82027 (4)	0.88239 (3)	0.06852 (14)
S1	0.22673 (8)	0.63699 (8)	0.49289 (5)	0.04950 (19)
S2	0.20227 (7)	0.65245 (7)	0.79943 (5)	0.04366 (17)
O1	0.3610 (3)	0.6577 (2)	0.46369 (16)	0.0639 (6)
O2	0.0875 (3)	0.5665 (2)	0.41649 (15)	0.0656 (6)
C1	0.2248 (3)	0.7998 (3)	0.5459 (2)	0.0449 (6)
C2	0.0995 (4)	0.8109 (3)	0.5695 (3)	0.0582 (8)
H2	0.0164	0.7326	0.5593	0.070*
C3	0.0980 (4)	0.9391 (4)	0.6086 (3)	0.0696 (9)
H3	0.0134	0.9463	0.6252	0.084*
C4	0.2194 (4)	1.0571 (3)	0.6237 (3)	0.0642 (9)
C5	0.3433 (4)	1.0432 (3)	0.6002 (3)	0.0672 (9)
H5	0.4261	1.1216	0.6100	0.081*
C6	0.3486 (3)	0.9157 (3)	0.5622 (2)	0.0574 (8)
H6	0.4344	0.9082	0.5479	0.069*
C7	0.2160 (6)	1.1970 (4)	0.6657 (4)	0.1016 (15)
H7A	0.3102	1.2506	0.7142	0.152*
H7B	0.1391	1.1858	0.7005	0.152*

H7C	0.1965	1.2435	0.6092	0.152*
N8	0.2273 (3)	0.5458 (2)	0.58395 (17)	0.0485 (6)
H8	0.1476	0.4793	0.5805	0.058*
C9	0.3552 (3)	0.5726 (3)	0.6691 (2)	0.0428 (6)
C10	0.4798 (4)	0.5487 (3)	0.6516 (2)	0.0599 (8)
H10	0.4800	0.5182	0.5838	0.072*
C11	0.6028 (4)	0.5690 (4)	0.7322 (3)	0.0629 (8)
H11	0.6868	0.5558	0.7185	0.076*
C12	0.6023 (3)	0.6090 (3)	0.8336 (2)	0.0527 (7)
C13	0.4792 (3)	0.6322 (3)	0.8528 (2)	0.0451 (6)
H13	0.4783	0.6592	0.9210	0.054*
C14	0.3561 (3)	0.6157 (2)	0.7709 (2)	0.0391 (6)
O15	0.7290 (2)	0.6271 (3)	0.9099 (2)	0.0810 (8)
C16	0.7150 (4)	0.5974 (4)	1.0061 (3)	0.0660 (9)
H16A	0.6837	0.6649	1.0407	0.099*
H16B	0.8090	0.5995	1.0480	0.099*
H16C	0.6425	0.5074	0.9960	0.099*
C17	0.2812 (3)	0.8343 (3)	0.83612 (18)	0.0391 (6)
N18	0.4152 (2)	0.8963 (2)	0.82825 (16)	0.0399 (5)
C19	0.4729 (3)	1.0343 (3)	0.8579 (2)	0.0444 (6)
N20	0.4065 (3)	1.1143 (2)	0.8998 (2)	0.0560 (6)
C21	0.2713 (4)	1.0476 (3)	0.9059 (2)	0.0584 (8)
H21	0.2212	1.0990	0.9337	0.070*
C22	0.2008 (3)	0.9085 (3)	0.8739 (2)	0.0480 (7)
N23	0.6066 (3)	1.0965 (2)	0.8425 (2)	0.0548 (6)
C24	0.6954 (4)	1.0140 (3)	0.8157 (3)	0.0648 (9)
H24A	0.6300	0.9209	0.7819	0.078*
H24B	0.7626	1.0088	0.8789	0.078*
C25	0.7831 (4)	1.0749 (4)	0.7447 (3)	0.0768 (10)
H25A	0.7158	1.0658	0.6771	0.092*
H25B	0.8488	1.0242	0.7350	0.092*
C26	0.8740 (4)	1.2256 (4)	0.7879 (3)	0.0828 (11)
H26A	0.9498	1.2343	0.8512	0.099*
H26B	0.9233	1.2638	0.7377	0.099*
C27	0.7764 (4)	1.3046 (4)	0.8112 (3)	0.0788 (11)
H27A	0.8377	1.3992	0.8440	0.095*
H27B	0.7097	1.3061	0.7463	0.095*
C28	0.6865 (4)	1.2435 (3)	0.8813 (3)	0.0685 (9)
H28A	0.7519	1.2578	0.9506	0.082*
H28B	0.6160	1.2903	0.8865	0.082*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0502 (2)	0.0873 (3)	0.0799 (3)	0.03164 (18)	0.02998 (17)	0.01633 (19)
S1	0.0587 (4)	0.0511 (4)	0.0322 (3)	0.0106 (3)	0.0140 (3)	0.0029 (3)
S2	0.0408 (4)	0.0443 (4)	0.0436 (4)	0.0102 (3)	0.0162 (3)	0.0030 (3)
O1	0.0750 (15)	0.0739 (15)	0.0493 (12)	0.0249 (12)	0.0317 (11)	0.0101 (11)

O2	0.0735 (15)	0.0658 (14)	0.0344 (10)	0.0018 (11)	0.0058 (10)	-0.0003 (10)
C1	0.0497 (16)	0.0481 (16)	0.0329 (13)	0.0118 (13)	0.0089 (12)	0.0108 (12)
C2	0.0544 (18)	0.0514 (18)	0.066 (2)	0.0141 (14)	0.0163 (15)	0.0143 (15)
C3	0.070 (2)	0.072 (2)	0.079 (2)	0.0362 (19)	0.0254 (19)	0.0203 (19)
C4	0.081 (2)	0.0513 (19)	0.058 (2)	0.0264 (18)	0.0082 (17)	0.0154 (15)
C5	0.071 (2)	0.0516 (19)	0.061 (2)	0.0040 (16)	0.0074 (17)	0.0133 (16)
C6	0.0541 (18)	0.0562 (19)	0.0553 (18)	0.0088 (15)	0.0157 (14)	0.0115 (15)
C7	0.137 (4)	0.063 (2)	0.107 (3)	0.050 (3)	0.019 (3)	0.012 (2)
N8	0.0601 (15)	0.0382 (12)	0.0391 (12)	0.0074 (11)	0.0136 (11)	0.0029 (10)
C9	0.0565 (16)	0.0323 (13)	0.0421 (15)	0.0160 (12)	0.0181 (13)	0.0060 (11)
C10	0.082 (2)	0.065 (2)	0.0509 (18)	0.0386 (18)	0.0351 (17)	0.0115 (15)
C11	0.065 (2)	0.076 (2)	0.071 (2)	0.0401 (18)	0.0359 (18)	0.0246 (18)
C12	0.0499 (17)	0.0580 (18)	0.0585 (18)	0.0205 (14)	0.0216 (14)	0.0241 (15)
C13	0.0482 (15)	0.0457 (15)	0.0420 (15)	0.0128 (12)	0.0170 (12)	0.0111 (12)
C14	0.0446 (14)	0.0304 (13)	0.0436 (14)	0.0107 (11)	0.0181 (12)	0.0065 (11)
O15	0.0464 (13)	0.125 (2)	0.0810 (17)	0.0292 (13)	0.0216 (12)	0.0489 (16)
C16	0.062 (2)	0.065 (2)	0.062 (2)	0.0191 (17)	0.0039 (16)	0.0112 (17)
C17	0.0440 (15)	0.0459 (15)	0.0286 (12)	0.0183 (12)	0.0095 (11)	0.0053 (11)
N18	0.0460 (12)	0.0388 (12)	0.0372 (11)	0.0153 (10)	0.0164 (10)	0.0046 (9)
C19	0.0560 (17)	0.0430 (15)	0.0368 (14)	0.0187 (13)	0.0165 (12)	0.0052 (12)
N20	0.0729 (17)	0.0441 (14)	0.0582 (15)	0.0233 (13)	0.0301 (13)	0.0045 (11)
C21	0.074 (2)	0.0546 (19)	0.0634 (19)	0.0356 (17)	0.0342 (17)	0.0089 (15)
C22	0.0498 (16)	0.0595 (18)	0.0442 (15)	0.0268 (14)	0.0198 (13)	0.0107 (13)
N23	0.0581 (15)	0.0410 (13)	0.0651 (16)	0.0119 (11)	0.0273 (13)	0.0036 (11)
C24	0.0604 (19)	0.0529 (19)	0.085 (2)	0.0163 (15)	0.0339 (18)	0.0089 (17)
C25	0.069 (2)	0.091 (3)	0.084 (3)	0.033 (2)	0.037 (2)	0.021 (2)
C26	0.065 (2)	0.087 (3)	0.101 (3)	0.016 (2)	0.036 (2)	0.039 (2)
C27	0.080 (2)	0.062 (2)	0.094 (3)	0.0160 (19)	0.027 (2)	0.029 (2)
C28	0.076 (2)	0.0458 (18)	0.073 (2)	0.0069 (16)	0.0225 (18)	0.0026 (16)

Geometric parameters (Å, °)

Br1—C22	1.885 (3)	C13—C14	1.390 (4)
S1—O1	1.423 (2)	C13—H13	0.9300
S1—O2	1.429 (2)	O15—C16	1.390 (4)
S1—N8	1.632 (2)	C16—H16A	0.9600
S1—C1	1.764 (3)	C16—H16B	0.9600
S2—C17	1.769 (3)	C16—H16C	0.9600
S2—C14	1.778 (3)	C17—N18	1.313 (3)
C1—C2	1.374 (4)	C17—C22	1.393 (4)
C1—C6	1.380 (4)	N18—C19	1.343 (3)
C2—C3	1.379 (5)	C19—N20	1.350 (3)
C2—H2	0.9300	C19—N23	1.351 (4)
C3—C4	1.382 (5)	N20—C21	1.325 (4)
C3—H3	0.9300	C21—C22	1.365 (4)
C4—C5	1.370 (5)	C21—H21	0.9300
C4—C7	1.509 (5)	N23—C28	1.456 (4)
C5—C6	1.383 (5)	N23—C24	1.467 (4)

C5—H5	0.9300	C24—C25	1.494 (5)
C6—H6	0.9300	C24—H24A	0.9700
C7—H7A	0.9600	C24—H24B	0.9700
C7—H7B	0.9600	C25—C26	1.513 (5)
C7—H7C	0.9600	C25—H25A	0.9700
N8—C9	1.425 (4)	C25—H25B	0.9700
N8—H8	0.8600	C26—C27	1.499 (5)
C9—C10	1.385 (4)	C26—H26A	0.9700
C9—C14	1.386 (4)	C26—H26B	0.9700
C10—C11	1.371 (5)	C27—C28	1.498 (5)
C10—H10	0.9300	C27—H27A	0.9700
C11—C12	1.380 (4)	C27—H27B	0.9700
C11—H11	0.9300	C28—H28A	0.9700
C12—O15	1.369 (4)	C28—H28B	0.9700
C12—C13	1.376 (4)		
O1—S1—O2	119.46 (13)	O15—C16—H16A	109.5
O1—S1—N8	108.46 (14)	O15—C16—H16B	109.5
O2—S1—N8	104.72 (13)	H16A—C16—H16B	109.5
O1—S1—C1	107.97 (14)	O15—C16—H16C	109.5
O2—S1—C1	107.66 (14)	H16A—C16—H16C	109.5
N8—S1—C1	108.12 (12)	H16B—C16—H16C	109.5
C17—S2—C14	99.34 (12)	N18—C17—C22	121.4 (3)
C2—C1—C6	120.3 (3)	N18—C17—S2	119.23 (19)
C2—C1—S1	119.8 (2)	C22—C17—S2	119.4 (2)
C6—C1—S1	119.9 (2)	C17—N18—C19	117.9 (2)
C1—C2—C3	119.4 (3)	N18—C19—N20	125.1 (3)
C1—C2—H2	120.3	N18—C19—N23	116.7 (2)
C3—C2—H2	120.3	N20—C19—N23	118.2 (3)
C2—C3—C4	121.5 (3)	C21—N20—C19	115.0 (3)
C2—C3—H3	119.3	N20—C21—C22	124.3 (3)
C4—C3—H3	119.3	N20—C21—H21	117.8
C5—C4—C3	118.0 (3)	C22—C21—H21	117.8
C5—C4—C7	121.0 (4)	C21—C22—C17	116.3 (3)
C3—C4—C7	121.0 (4)	C21—C22—Br1	122.2 (2)
C4—C5—C6	121.7 (3)	C17—C22—Br1	121.5 (2)
C4—C5—H5	119.1	C19—N23—C28	121.4 (2)
C6—C5—H5	119.1	C19—N23—C24	120.1 (2)
C1—C6—C5	119.1 (3)	C28—N23—C24	115.9 (3)
C1—C6—H6	120.4	N23—C24—C25	111.5 (3)
C5—C6—H6	120.4	N23—C24—H24A	109.3
C4—C7—H7A	109.5	C25—C24—H24A	109.3
C4—C7—H7B	109.5	N23—C24—H24B	109.3
H7A—C7—H7B	109.5	C25—C24—H24B	109.3
C4—C7—H7C	109.5	H24A—C24—H24B	108.0
H7A—C7—H7C	109.5	C24—C25—C26	111.6 (3)
H7B—C7—H7C	109.5	C24—C25—H25A	109.3
C9—N8—S1	122.28 (19)	C26—C25—H25A	109.3

C9—N8—H8	118.9	C24—C25—H25B	109.3
S1—N8—H8	118.9	C26—C25—H25B	109.3
C10—C9—C14	118.4 (3)	H25A—C25—H25B	108.0
C10—C9—N8	120.0 (2)	C27—C26—C25	110.4 (3)
C14—C9—N8	121.5 (2)	C27—C26—H26A	109.6
C11—C10—C9	121.4 (3)	C25—C26—H26A	109.6
C11—C10—H10	119.3	C27—C26—H26B	109.6
C9—C10—H10	119.3	C25—C26—H26B	109.6
C10—C11—C12	120.0 (3)	H26A—C26—H26B	108.1
C10—C11—H11	120.0	C28—C27—C26	112.5 (3)
C12—C11—H11	120.0	C28—C27—H27A	109.1
O15—C12—C13	123.8 (3)	C26—C27—H27A	109.1
O15—C12—C11	116.7 (3)	C28—C27—H27B	109.1
C13—C12—C11	119.5 (3)	C26—C27—H27B	109.1
C12—C13—C14	120.4 (3)	H27A—C27—H27B	107.8
C12—C13—H13	119.8	N23—C28—C27	111.5 (3)
C14—C13—H13	119.8	N23—C28—H28A	109.3
C9—C14—C13	120.1 (2)	C27—C28—H28A	109.3
C9—C14—S2	121.1 (2)	N23—C28—H28B	109.3
C13—C14—S2	118.7 (2)	C27—C28—H28B	109.3
C12—O15—C16	118.1 (2)	H28A—C28—H28B	108.0
O1—S1—C1—C2	172.6 (2)	C12—C13—C14—S2	-178.0 (2)
O2—S1—C1—C2	42.4 (3)	C17—S2—C14—C9	-108.1 (2)
N8—S1—C1—C2	-70.2 (3)	C17—S2—C14—C13	71.3 (2)
O1—S1—C1—C6	-6.6 (3)	C13—C12—O15—C16	34.4 (5)
O2—S1—C1—C6	-136.8 (2)	C11—C12—O15—C16	-147.6 (3)
N8—S1—C1—C6	110.5 (2)	C14—S2—C17—N18	4.4 (2)
C6—C1—C2—C3	0.7 (5)	C14—S2—C17—C22	-174.8 (2)
S1—C1—C2—C3	-178.5 (2)	C22—C17—N18—C19	0.0 (4)
C1—C2—C3—C4	0.6 (5)	S2—C17—N18—C19	-179.23 (19)
C2—C3—C4—C5	-1.0 (5)	C17—N18—C19—N20	3.2 (4)
C2—C3—C4—C7	179.5 (3)	C17—N18—C19—N23	-175.7 (2)
C3—C4—C5—C6	0.1 (5)	N18—C19—N20—C21	-3.4 (4)
C7—C4—C5—C6	179.6 (3)	N23—C19—N20—C21	175.5 (3)
C2—C1—C6—C5	-1.6 (4)	C19—N20—C21—C22	0.5 (5)
S1—C1—C6—C5	177.6 (2)	N20—C21—C22—C17	2.2 (5)
C4—C5—C6—C1	1.2 (5)	N20—C21—C22—Br1	-178.8 (2)
O1—S1—N8—C9	43.7 (2)	N18—C17—C22—C21	-2.5 (4)
O2—S1—N8—C9	172.3 (2)	S2—C17—C22—C21	176.7 (2)
C1—S1—N8—C9	-73.1 (2)	N18—C17—C22—Br1	178.55 (19)
S1—N8—C9—C10	-67.8 (3)	S2—C17—C22—Br1	-2.3 (3)
S1—N8—C9—C14	115.5 (3)	N18—C19—N23—C28	-173.6 (3)
C14—C9—C10—C11	-1.1 (4)	N20—C19—N23—C28	7.4 (4)
N8—C9—C10—C11	-177.8 (3)	N18—C19—N23—C24	-12.3 (4)
C9—C10—C11—C12	2.5 (5)	N20—C19—N23—C24	168.7 (3)
C10—C11—C12—O15	179.9 (3)	C19—N23—C24—C25	146.5 (3)
C10—C11—C12—C13	-1.9 (5)	C28—N23—C24—C25	-51.2 (4)

O15—C12—C13—C14	178.0 (3)	N23—C24—C25—C26	52.4 (4)
C11—C12—C13—C14	0.0 (4)	C24—C25—C26—C27	-54.7 (4)
C10—C9—C14—C13	-0.9 (4)	C25—C26—C27—C28	54.2 (5)
N8—C9—C14—C13	175.8 (2)	C19—N23—C28—C27	-147.7 (3)
C10—C9—C14—S2	178.5 (2)	C24—N23—C28—C27	50.3 (4)
N8—C9—C14—S2	-4.8 (3)	C26—C27—C28—N23	-51.4 (5)
C12—C13—C14—C9	1.4 (4)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N8—H8 \cdots O2 ⁱ	0.86	2.22	2.955 (4)	143

Symmetry code: (i) $-x, -y+1, -z+1$.