

Redetermination of junitoite, CaZn₂Si₂O₇·H₂O

 Hexiong Yang,^{a*} Neil G. Jenkins^b and Robert T. Downs^a
^aDepartment of Geosciences, University of Arizona, 1040 E. 4th Street, Tucson, Arizona 85721-0077, USA, and ^bSabino High School, 5000 North Bowes Road, Tucson, Arizona 85749, USA

Correspondence e-mail: hyang@u.arizona.edu

Received 28 August 2012; accepted 31 August 2012

 Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{Si}-\text{O}) = 0.001$ Å; R factor = 0.017; wR factor = 0.044; data-to-parameter ratio = 19.3.

The crystal structure of the mineral junitoite, ideally CaZn₂Si₂O₇·H₂O (calcium dizinc disilicate monohydrate), was first determined by Hamilton & Finney [*Mineral. Mag.* (1985), **49**, 91–95] based on the space group *Ama2*, yielding a reliability factor R of 0.10, with isotropic displacement parameters for all non-H atoms. The present study reports a structure redetermination of junitoite using single-crystal X-ray diffraction data from a natural sample, demonstrating that the space group of this mineral is actually *Aea2*, which can be attained simply by shifting the origin. Topologically, the structure models in the space groups *Aea2* and *Ama2* are analogous, consisting of chains of corner-sharing ZnO₄ tetrahedra parallel to the b axis, cross-linked by Si₂O₇ tetrahedral dimers (the site symmetry of the bridging O atom is ..2) along a and c , forming a three-dimensional framework. The Ca²⁺ cations (site symmetry ..2) are situated in cavities of the framework and are bonded to five O atoms and one H₂O molecule (site symmetry ..2) in a distorted octahedral coordination environment. However, some bond lengths, especially for the SiO₄ tetrahedron, are noticeably different between the two structure models. Hydrogen bonding in junitoite is found between the water molecule and a framework O atom.

Related literature

For junitoite, see: Williams (1976); Hamilton & Finney (1985). For junitoite-related minerals and compounds, see: Lin *et al.* (1999); Fleet & Liu (2001); Kolitsch *et al.* (2009); Yang *et al.* (2012). Parameters for bond-valence calculations were taken from Brese & O'Keeffe (1991).

Experimental

Crystal data

CaZn ₂ Si ₂ O ₇ ·H ₂ O	$V = 676.5$ (3) Å ³
$M_r = 357.02$	$Z = 4$
Orthorhombic, <i>Aea2</i>	Mo $K\alpha$ radiation
$a = 12.530$ (4) Å	$\mu = 8.21$ mm ⁻¹
$b = 6.3056$ (18) Å	$T = 293$ K
$c = 8.562$ (3) Å	$0.06 \times 0.06 \times 0.05$ mm

Data collection

Bruker APEXII CCD area-detector diffractometer	2719 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 2005)	1257 independent reflections
$T_{\min} = 0.639$, $T_{\max} = 0.684$	1189 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.021$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.017$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.044$	$\Delta\rho_{\text{max}} = 0.52$ e Å ⁻³
$S = 1.07$	$\Delta\rho_{\text{min}} = -0.65$ e Å ⁻³
1257 reflections	Absolute structure: Flack (1983),
65 parameters	580 Friedel pairs
1 restraint	Flack parameter: 0.023 (12)

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
OW5–H···O1	0.70 (3)	2.18 (2)	2.875 (2)	170 (3)

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *Xtal-Draw* (Downs & Hall-Wallace, 2003); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We gratefully acknowledge support of this study by the Arizona Science Foundation.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WM2677).

References

- Brese, N. E. & O'Keeffe, M. (1991). *Acta Cryst.* **B47**, 192–197.
 Bruker (2004). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Downs, R. T. & Hall-Wallace, M. (2003). *Am. Mineral.* **88**, 247–250.
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
 Fleet, M. E. & Liu, X. (2001). *J. Solid State Chem.* **161**, 166–172.
 Hamilton, R. D. & Finney, J. J. (1985). *Mineral. Mag.* **49**, 91–95.
 Kolitsch, U., Wierzbicka-Wieczorek, M. & Tillmanns, E. (2009). *Can. Mineral.* **47**, 421–431.
 Lin, J. H., Lu, G. X., Du, J., Su, M. Z., Loong, C.-K. & Richardson, J. W. Jr (1999). *J. Phys. Chem. Solids*, **60**, 975–983.
 Sheldrick, G. M. (2005). *SADABS*. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
 Williams, S. A. (1976). *Am. Mineral.* **61**, 1255–1258.
 Yang, H., Downs, R. T., Evans, S. H. & Pinch, W. W. (2012). *Am. Mineral.* In the press.

supporting information

Acta Cryst. (2012). E68, i73 [https://doi.org/10.1107/S1600536812037622]

Redetermination of junitoite, $\text{CaZn}_2\text{Si}_2\text{O}_7 \cdot \text{H}_2\text{O}$

Hexiong Yang, Neil G. Jenkins and Robert T. Downs

S1. Comment

Junitoite, $\text{CaZn}_2\text{Si}_2\text{O}_7 \cdot \text{H}_2\text{O}$, from the Christmas Mine, Gila County, Arizona was first described by Williams (1976) with orthorhombic symmetry in space group *Bbm2* (non-standard setting of space group No. 40) and unit-cell parameters $a = 6.309$, $b = 12.503$, $c = 8.549$ Å. By adopting the standard unit-cell setting of this space group in *Ama2* ($a = 12.510$, $b = 6.318$, $c = 8.561$ Å) for this mineral, Hamilton & Finney (1985) noted that while the Weissenberg photographic data pointed to *Ama2*, the X-ray diffractometer data were also compatible with the space group *Aea2*. Although the two space groups yielded similar reliability factors $R1 \sim 0.10$ with isotropic displacement parameters for all atoms (H atoms were not located), Hamilton & Finney (1985) chose *Ama2* for their final structure report because it "produces less distortion of the coordination polyhedra and provides a structure in which the site symmetry of the cations is more similar to other zinc silicates". Their attempts at refinement with anisotropic displacement parameters resulted in non-positive definite displacement parameters for a number of atoms. In our efforts to understand the hydrogen bonding environments in minerals and their relationships to Raman spectra, we concluded that the structural model for junitoite needed improvement. This study reports a structure redetermination of junitoite from the type locality by means of single-crystal X-ray diffraction data, demonstrating that the space group of this mineral is actually *Aea2*, rather than *Ama2*.

The crystal structure of junitoite consists of chains of corner-sharing ZnO_4 tetrahedra parallel to the b axis, cross-linked by Si_2O_7 tetrahedral dimers along a and c to form a three-dimensional framework. The Ca^{2+} cations, situated in cavities of the framework, are bonded to five O atoms and one H_2O molecule in a distorted octahedral $[\text{CaO}_5(\text{H}_2\text{O})]$ coordination environment (Figs. 1, 2). As described below, it may be useful to consider that there is a $\text{Ca}-\text{H}_2\text{O}$ bonded pair in the cavity. The structure of junitoite in space group *Aea2* resembles that in space group *Ama2* (Hamilton & Finney, 1985). In fact, as noted by Hamilton & Finney (1985), the structure model in *Aea2* can be attained simply by shifting the origin of the structure model in *Ama2* from (x, y, z) to $(x - 1/4, y - 1/4, z)$. Upon this shift, the only major structural change is that the two unique Zn atoms at the $4a$ sites in the *Ama2* structure model are transformed into a single atom at the $8b$ site in the *Aea2* structure model. The numbers and coordination polyhedra of the distinct Ca, Si, and O sites remain unaffected. However, some bond lengths are noticeably different between the two structure models. For example, the Si—O, Zn—O, and Ca—O bond lengths range from 1.55 (5) to 1.69 (5) Å, 1.93 (4) to 1.99 (4) Å, and 2.29 (7) to 2.44 (5) Å in the *Ama2* structure model, respectively, but from 1.6130 (14) to 1.6719 (12) Å, 1.9454 (13) to 1.9691 (13) Å, and 2.286 (2) to 2.439 (2) Å in the *Aea2* structure model. The Si—O—Si angle within the Si_2O_7 disilicate group is $124.8 (1)^\circ$ in our study, which is slightly greater than that (122.4°) determined by Hamilton & Finney (1985).

The hydrogen bond in junitoite is found between Ow5 and O1, with Ow5 as the donor and O1 as the acceptor. This agrees with the calculated bond-valence sums of 0.42 valence units for Ow5 and 1.77 valence units for O1 by using the parameters given by Brese & O'Keeffe (1991). For numerical details of the hydrogen-bonding geometry, see: Table 1.

Remarkably, junitoite is topologically related to a group of compounds with the general formula $BaM^{2+}_2Si_2O_7$, where M = Be (barylite and clinobarylite), Fe (andremeyerite), Cu (scottyite), and Mg, Mn, Co, and Zn in synthetic phases. These Ba-silicates are all comprised of corner-sharing MO_4 tetrahedral chains that are interlinked by Si_2O_7 tetrahedral dimers and Ba^{2+} cations, despite their diverse structural symmetries (Yang *et al.*, 2012). Intriguingly, there is no documentation for any $SrM_2Si_2O_7$ compounds. It then begs the question whether the $BaM_2Si_2O_7$ compounds are capable of accommodating a significant amount of cations smaller than Ba^{2+} . Similar to the pair ($Ca^{2+} + H_2O$) in junitoite, the Ba^{2+} cations in the $BaM_2Si_2O_7$ structures are also situated in the cavities of the framework formed by the Si_2O_7 dimers and the MO_4 tetrahedral chains. Conceivably, any substantial replacement of large Ba^{2+} by smaller divalent cations (such as Sr^{2+}) would require, in addition to the other structural adjustments (such as the tilting or distortion of MO_4 and/or SiO_4 tetrahedra), a further narrowing of the Si—O—Si angle in the Si_2O_7 group in order to satisfy the bonding environment for smaller cations. This, however, would not be energetically favorable, because the Si—O—Si angles in the $BaM_2Si_2O_7$ compounds, ranging from 124 to 135°, are already among the smallest of disilicate materials, *e.g.* for high-temperature $BaZn_2Si_2O_7$ (Lin *et al.*, 1999), high-pressure rare earth (*RE*) disilicates $RE_2Si_2O_7$ (Fleet & Liu, 2001) or $BaKY(Si_2O_7)$ (Kolitsch *et al.*, 2009). Accordingly, any sizable substitution of smaller Sr^{2+} for Ba^{2+} would worsen the bonding energetics for this site and thus destabilize the structure. For junitoite, Ca^{2+} by itself, which is even smaller than Sr^{2+} , is apparently too small to occupy the cavities in the framework. Therefore, the presence of the $H_2O—Ca^{2+}$ bonded pair is essential to stabilize its structure. By the same token, one could argue that the pair ($Sr^{2+} + H_2O$) together may be too large for the cavities in the structures analogous to those for the $BaM_2Si_2O_7$ materials, since there is no report for any $SrM^{2+}_2Si_2O_7.H_2O$ compound up to date. Based on this reasoning, we postulate that more compounds with composition $CaM^{2+}_2Si_2O_7.H_2O$ may be found in nature or synthesized in laboratories. Furthermore, it would be interesting if the Sr— H_2O pair might be found in digermanates, where this structural unit is even larger.

S2. Experimental

The junitoite crystal used in this study is from the type locality, the Christmas Mine, Gila County, Arizona and is in the collection of the RRUFF project (<http://rruff.info/R120100>). Its chemical composition measured by Williams (1976) is $Ca_{0.98}Zn_{1.96}Si_{1.84}O_{6.6}.1.13H_2O$.

S3. Refinement

The H atom was located near Ow5 from difference Fourier syntheses and its position refined freely with a fixed isotropic displacement ($U_{iso} = 0.03$). For simplicity, the ideal chemistry, $CaZn_2Si_2O_7.H_2O$, was assumed during the final refinement. The highest residual peak in the difference Fourier maps was located at (0.6270, 0.1150, 0.9598), 0.69 Å from O1, and the deepest hole at (0.3072, 0.2596, 0.0874), 0.81 Å from Zn.




Figure 1

Crystal structure of junitoite. The gray, green, and small blue spheres represent Ca, Ow5, and H atoms, respectively. The yellow and red tetrahedra represent ZnO_4 and SiO_4 groups, respectively.




Figure 2

Atoms in junitoite with corresponding ellipsoids at the 99.9% probability level. The gray, yellow, purple, red, and green ellipsoids represent Ca, Zn, Si, O, and Ow5, respectively. The small blue spheres represent H atoms.

calcium dizinc disilicate monohydrate

Crystal data

CaZn₂Si₂O₇·H₂O $M_r = 357.02$ Orthorhombic, *Aea2*

Hall symbol: A 2 -2ac

 $a = 12.530 (4) \text{ \AA}$ $b = 6.3056 (18) \text{ \AA}$ $c = 8.562 (3) \text{ \AA}$ $V = 676.5 (3) \text{ \AA}^3$ $Z = 4$ $F(000) = 696$ $D_x = 3.506 \text{ Mg m}^{-3}$ Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 1387 reflections

 $\theta = 4.3\text{--}33.2^\circ$ $\mu = 8.21 \text{ mm}^{-1}$ $T = 293 \text{ K}$

Cuboid, colorless

 $0.06 \times 0.06 \times 0.05 \text{ mm}$

Data collection

Bruker APEXII CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 φ and ω scanAbsorption correction: multi-scan
(*SADABS*; Sheldrick, 2005) $T_{\min} = 0.639$, $T_{\max} = 0.684$

2719 measured reflections

1257 independent reflections

1189 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.021$ $\theta_{\max} = 33.3^\circ$, $\theta_{\min} = 3.3^\circ$ $h = -8 \rightarrow 19$ $k = -9 \rightarrow 8$ $l = -12 \rightarrow 13$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.017$ $wR(F^2) = 0.044$ $S = 1.07$

1257 reflections

65 parameters

1 restraint

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
map

Hydrogen site location: difference Fourier map

H atoms treated by a mixture of independent
and constrained refinement $w = 1/[\sigma^2(F_o^2) + (0.0094P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} = 0.001$ $\Delta\rho_{\max} = 0.52 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -0.65 \text{ e \AA}^{-3}$ Extinction correction: *SHELXL97* (Sheldrick,
2008), $F_c^* = kFc^*[1 + 0.001x\lambda^3/\sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0120 (4)

Absolute structure: Flack (1983), 580 Friedel
pairs

Absolute structure parameter: 0.023 (12)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Ca	0.0000	0.5000	0.51882 (6)	0.00899 (12)
Zn	0.252164 (9)	0.244531 (17)	0.13494 (6)	0.00808 (7)

Si	0.11827 (4)	0.00274 (4)	0.39406 (7)	0.00615 (9)
O1	0.12311 (9)	0.2992 (2)	0.01003 (15)	0.0107 (3)
O2	0.12739 (10)	0.21991 (19)	0.49272 (17)	0.0099 (3)
O3	0.20596 (13)	-0.00482 (13)	0.25347 (15)	0.0101 (3)
O4	0.0000	0.0000	0.3037 (2)	0.0082 (3)
OW5	0.0000	0.5000	0.2518 (3)	0.0209 (7)
H	0.035 (3)	0.448 (4)	0.200 (3)	0.030*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ca	0.0106 (2)	0.0078 (2)	0.0085 (2)	0.00019 (13)	0.000	0.000
Zn	0.00755 (10)	0.00739 (10)	0.00931 (11)	-0.00067 (7)	0.00070 (15)	-0.00072 (18)
Si	0.00563 (18)	0.00687 (18)	0.00596 (19)	-0.00022 (11)	0.00019 (16)	0.00035 (16)
O1	0.0084 (5)	0.0112 (5)	0.0126 (8)	-0.0010 (4)	-0.0020 (4)	0.0046 (5)
O2	0.0086 (5)	0.0109 (5)	0.0102 (8)	0.0019 (4)	-0.0026 (4)	-0.0035 (5)
O3	0.0124 (7)	0.0080 (6)	0.0099 (6)	-0.0001 (3)	0.0045 (4)	0.0003 (4)
O4	0.0063 (8)	0.0112 (8)	0.0071 (9)	0.0000 (4)	0.000	0.000
OW5	0.0272 (15)	0.0278 (18)	0.0076 (11)	0.0100 (7)	0.000	0.000

Geometric parameters (Å, °)

Ca—OW5	2.286 (2)	Zn—O3 ^v	1.9501 (11)
Ca—O2 ⁱ	2.3910 (12)	Zn—O3	1.9589 (11)
Ca—O2	2.3910 (12)	Zn—O1	1.9691 (13)
Ca—O1 ⁱⁱ	2.4381 (13)	Si—O2	1.6130 (14)
Ca—O1 ⁱⁱⁱ	2.4381 (13)	Si—O1 ^{vi}	1.6239 (14)
Ca—O4 ⁱⁱ	2.439 (2)	Si—O3	1.6305 (16)
Zn—O2 ^{iv}	1.9454 (13)	Si—O4	1.6719 (12)
OW5—Ca—O2 ⁱ	84.64 (4)	O1 ⁱⁱⁱ —Ca—O4 ⁱⁱ	91.77 (4)
OW5—Ca—O2	84.64 (4)	O2 ^{iv} —Zn—O3 ^v	100.47 (6)
O2 ⁱ —Ca—O2	169.28 (8)	O2 ^{iv} —Zn—O3	119.31 (6)
OW5—Ca—O1 ⁱⁱ	88.23 (4)	O3 ^v —Zn—O3	117.42 (7)
O2 ⁱ —Ca—O1 ⁱⁱ	81.25 (6)	O2 ^{iv} —Zn—O1	108.13 (8)
O2—Ca—O1 ⁱⁱ	98.41 (6)	O3 ^v —Zn—O1	111.21 (6)
OW5—Ca—O1 ⁱⁱⁱ	88.23 (4)	O3—Zn—O1	100.32 (6)
O2 ⁱ —Ca—O1 ⁱⁱⁱ	98.41 (6)	O2—Si—O1 ^{vi}	110.37 (11)
O2—Ca—O1 ⁱⁱⁱ	81.25 (6)	O2—Si—O3	111.33 (7)
O1 ⁱⁱ —Ca—O1 ⁱⁱⁱ	176.46 (7)	O1 ^{vi} —Si—O3	113.77 (7)
OW5—Ca—O4 ⁱⁱ	180.0	O2—Si—O4	108.30 (6)
O2 ⁱ —Ca—O4 ⁱⁱ	95.36 (4)	O1 ^{vi} —Si—O4	107.94 (6)
O2—Ca—O4 ⁱⁱ	95.36 (4)	O3—Si—O4	104.79 (10)
O1 ⁱⁱ —Ca—O4 ⁱⁱ	91.77 (4)		

Symmetry codes: (i) $-x, -y+1, z$; (ii) $x, y+1/2, z+1/2$; (iii) $-x, -y+1/2, z+1/2$; (iv) $-x+1/2, y, z-1/2$; (v) $-x+1/2, y+1/2, z$; (vi) $x, y-1/2, z+1/2$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
OW5—H···O1	0.70 (3)	2.18 (2)	2.875 (2)	170 (3)