

## 1H-Indole-3-carbaldehyde

C. S. Dileep,<sup>a</sup> M. M. M. Abdoh,<sup>b</sup> M. P. Chakravarthy,<sup>c</sup>  
K. N. Mohana<sup>c</sup> and M. A. Sridhar<sup>a\*</sup>

<sup>a</sup>Department of Studies in Physics, Manasagangotri, University of Mysore, Mysore 570 006, India, <sup>b</sup>Department of Physics, Faculty of Science, An Najah National University, Nabtus, West Bank, Palestinian Territories, and <sup>c</sup>Department of Studies in Chemistry, Manasagangotri, University of Mysore, Mysore 570 006, India  
Correspondence e-mail: mas@physics.uni-mysore.ac.in


Received 14 September 2012; accepted 25 September 2012

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$ ;  $R$  factor = 0.029;  $wR$  factor = 0.069; data-to-parameter ratio = 7.1.

In the title compound,  $\text{C}_9\text{H}_7\text{NO}$ , the benzene ring forms a dihedral angle of  $3.98(12)^\circ$  with the pyrrole ring. In the crystal,  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds links the molecules into chains which run parallel to  $[02\bar{1}]$ .

### Related literature

For a related structure, see: Rizal *et al.* (2008).



### Experimental

#### Crystal data

$\text{C}_9\text{H}_7\text{NO}$

$M_r = 145.16$

Orthorhombic,  $Pca2_1$   
 $a = 14.0758(9)\text{ \AA}$   
 $b = 5.8059(4)\text{ \AA}$   
 $c = 8.6909(5)\text{ \AA}$   
 $V = 710.24(8)\text{ \AA}^3$

$Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 0.09\text{ mm}^{-1}$   
 $T = 293\text{ K}$   
 $0.30 \times 0.20 \times 0.20\text{ mm}$

#### Data collection

Bruker Kappa APEXII CCD  
diffractometer  
3791 measured reflections

775 independent reflections  
699 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.024$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.029$   
 $wR(F^2) = 0.069$   
 $S = 1.08$   
775 reflections  
109 parameters  
1 restraint

H atoms treated by a mixture of  
independent and constrained  
refinement  
 $\Delta\rho_{\text{max}} = 0.11\text{ e \AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.09\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1A $\cdots$ O1 <sup>i</sup>	0.94 (3)	1.92 (3)	2.831 (2)	165 (3)

Symmetry code: (i)  $-x + \frac{1}{2}, y - 1, z + \frac{1}{2}$ .

Data collection: *APEX2* (Bruker, 2004); cell refinement: *APEX2* and *SAINT* (Bruker, 2004); data reduction: *SAINT* and *XPREP* (Bruker, 2004); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *PLATON*.

CSD would like to acknowledge the UGC–BRS and the University of Mysore for financial assistance.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GO2070).

### References

- Bruker (2004). *APEX2, SAINT* and *XPREP*. Bruker AXS Inc., Madison, Wisconsin, USA.  
Rizal, M. R., Ali, H. M. & Ng, S. W. (2008). *Acta Cryst. E64*, o555.  
Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.  
Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

# supporting information

*Acta Cryst.* (2012). E68, o3135 [doi:10.1107/S1600536812040573]

## 1*H*-Indole-3-carbaldehyde

C. S. Dileep, M. M. M. Abdoh, M. P. Chakravarthy, K. N. Mohana and M. A. Sridhar

### S1. Comment


The molecule is shown with its labelling in Figure 1. The molecules are connected into one-dimensional chains by the N1—H1A···O1(3/2-x,-1+y,1/2+z) hydrogen bond which links the molecules into one dimensional chains which run parallel to [2-10], Table 1 and Figure2.

### S2. Experimental

Indole was converted to 1*H*-indole-3-carbaldehyde in the presence of DMF, POCl<sub>3</sub>, NaOH. 1*H*-indole-3-carbaldehyde was taken and recrystallized in methanol solvent. The purity of the compound is confirmed by the TLC. A little quantity of compound was taken again for recrystallization to get a pure crystal in methanol solvent medium.


### S3. Refinement

H atoms were treated as riding atoms with C—H(aromatic), 0.93 Å with  $U_{\text{iso}} = 1.2U_{\text{eq}}(\text{C})$ . The H atoms attached to C1 and N1 were located on a difference map and refined isotropically. Friedel pairs were merged.



**Figure 1**

ORTEP of the title compound with the ellipsoids for non-H atoms are drawn at the 50% probability.

**Figure 2**

Molecular packing view of down the *b* axis. Dashed lines indicate hydrogen bonds.

### **1*H*-Indole-3-carbaldehyde**

#### *Crystal data*

$C_9H_7NO$   
 $M_r = 145.16$   
Orthorhombic,  $Pca2_1$   
Hall symbol: P 2c -2ac  
 $a = 14.0758 (9) \text{ \AA}$   
 $b = 5.8059 (4) \text{ \AA}$   
 $c = 8.6909 (5) \text{ \AA}$   
 $V = 710.24 (8) \text{ \AA}^3$

$Z = 4$   
 $F(000) = 304$   
 $D_x = 1.357 \text{ Mg m}^{-3}$   
Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$   
 $\mu = 0.09 \text{ mm}^{-1}$   
 $T = 293 \text{ K}$   
Block, colourless  
 $0.30 \times 0.20 \times 0.20 \text{ mm}$

#### *Data collection*

Bruker Kappa APEXII CCD  
diffractometer  
Radiation source: fine-focus sealed tube

Graphite monochromator  
 $\omega$  and  $\varphi$  scan  
3791 measured reflections

775 independent reflections  
 699 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.024$   
 $\theta_{\text{max}} = 26.5^\circ, \theta_{\text{min}} = 2.9^\circ$

$h = -17 \rightarrow 16$   
 $k = -6 \rightarrow 7$   
 $l = -10 \rightarrow 8$

#### Refinement

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.029$   
 $wR(F^2) = 0.069$   
 $S = 1.08$   
 775 reflections  
 109 parameters  
 1 restraint  
 Primary atom site location: structure-invariant direct methods  
 Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites  
 H atoms treated by a mixture of independent and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.0297P)^2 + 0.0895P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\text{max}} = 0.002$   
 $\Delta\rho_{\text{max}} = 0.11 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.09 \text{ e } \text{\AA}^{-3}$   
 Extinction correction: *SHELXL*,  $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$   
 Extinction coefficient: 0.031 (5)

#### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

#### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.76788 (16)	0.4092 (3)	-0.0642 (3)	0.0443 (5)
C2	0.72368 (14)	0.2453 (4)	0.0341 (3)	0.0401 (5)
C3	0.76781 (17)	0.0501 (4)	0.0871 (3)	0.0489 (6)
H3	0.8297	0.0074	0.0628	0.059*
C4	0.62446 (14)	0.0451 (3)	0.1907 (3)	0.0418 (5)
C5	0.54556 (17)	-0.0080 (4)	0.2797 (3)	0.0500 (6)
H5	0.5427	-0.1430	0.3371	0.060*
C6	0.47250 (16)	0.1462 (4)	0.2795 (3)	0.0535 (6)
H6	0.4189	0.1166	0.3388	0.064*
C7	0.47646 (15)	0.3467 (4)	0.1924 (3)	0.0526 (6)
H7	0.4255	0.4487	0.1951	0.063*
C8	0.55405 (16)	0.3976 (4)	0.1024 (3)	0.0449 (5)
H8	0.5555	0.5310	0.0433	0.054*
C9	0.63026 (14)	0.2450 (3)	0.1019 (2)	0.0370 (5)
N1	0.70988 (14)	-0.0700 (3)	0.1783 (2)	0.0510 (5)
O1	0.72964 (12)	0.5795 (2)	-0.1176 (2)	0.0565 (5)
H1	0.8360 (16)	0.376 (4)	-0.086 (3)	0.054 (6)*
H1A	0.7253 (17)	-0.204 (4)	0.233 (4)	0.073 (8)*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0516 (13)	0.0432 (10)	0.0382 (12)	-0.0002 (9)	0.0027 (11)	-0.0014 (10)
C2	0.0514 (11)	0.0368 (9)	0.0323 (10)	0.0032 (9)	0.0004 (9)	0.0009 (8)
C3	0.0557 (14)	0.0476 (11)	0.0435 (14)	0.0109 (11)	0.0050 (11)	0.0022 (10)
C4	0.0557 (12)	0.0341 (9)	0.0354 (11)	-0.0017 (8)	-0.0038 (10)	0.0013 (10)
C5	0.0666 (14)	0.0435 (12)	0.0399 (13)	-0.0135 (11)	-0.0021 (12)	0.0050 (10)
C6	0.0499 (13)	0.0624 (14)	0.0484 (14)	-0.0136 (11)	0.0036 (11)	-0.0011 (12)
C7	0.0466 (12)	0.0576 (13)	0.0536 (15)	0.0036 (10)	-0.0017 (12)	-0.0041 (14)
C8	0.0512 (12)	0.0406 (10)	0.0428 (13)	0.0013 (9)	-0.0046 (11)	0.0037 (9)
C9	0.0473 (11)	0.0344 (10)	0.0293 (10)	-0.0027 (8)	-0.0052 (9)	-0.0017 (8)
N1	0.0680 (12)	0.0395 (9)	0.0455 (11)	0.0102 (8)	0.0016 (10)	0.0113 (10)
O1	0.0658 (11)	0.0442 (8)	0.0596 (11)	-0.0002 (7)	0.0058 (8)	0.0163 (8)

Geometric parameters ( $\text{\AA}$ ,  $\text{^\circ}$ )

C1—O1	1.218 (2)	C5—C6	1.363 (3)
C1—C2	1.422 (3)	C5—H5	0.9300
C1—H1	1.00 (2)	C6—C7	1.390 (3)
C2—C3	1.372 (3)	C6—H6	0.9300
C2—C9	1.441 (3)	C7—C8	1.375 (3)
C3—N1	1.334 (3)	C7—H7	0.9300
C3—H3	0.9300	C8—C9	1.391 (3)
C4—N1	1.380 (3)	C8—H8	0.9300
C4—C5	1.388 (3)	N1—H1A	0.94 (2)
C4—C9	1.396 (3)		
O1—C1—C2	125.4 (2)	C5—C6—C7	121.4 (2)
O1—C1—H1	120.6 (14)	C5—C6—H6	119.3
C2—C1—H1	114.0 (14)	C7—C6—H6	119.3
C3—C2—C1	123.8 (2)	C8—C7—C6	121.4 (2)
C3—C2—C9	105.93 (19)	C8—C7—H7	119.3
C1—C2—C9	130.22 (19)	C6—C7—H7	119.3
N1—C3—C2	110.8 (2)	C7—C8—C9	118.5 (2)
N1—C3—H3	124.6	C7—C8—H8	120.7
C2—C3—H3	124.6	C9—C8—H8	120.7
N1—C4—C5	129.30 (19)	C8—C9—C4	118.84 (19)
N1—C4—C9	107.95 (18)	C8—C9—C2	134.75 (19)
C5—C4—C9	122.65 (19)	C4—C9—C2	106.31 (18)
C6—C5—C4	117.2 (2)	C3—N1—C4	109.04 (17)
C6—C5—H5	121.4	C3—N1—H1A	126.4 (16)
C4—C5—H5	121.4	C4—N1—H1A	124.4 (16)
O1—C1—C2—C3	177.2 (2)	N1—C4—C9—C8	-176.6 (2)
O1—C1—C2—C9	-4.9 (4)	C5—C4—C9—C8	0.1 (3)
C1—C2—C3—N1	179.3 (2)	N1—C4—C9—C2	0.2 (2)
C9—C2—C3—N1	1.0 (3)	C5—C4—C9—C2	176.9 (2)

N1—C4—C5—C6	175.0 (2)	C3—C2—C9—C8	175.3 (2)
C9—C4—C5—C6	-0.9 (3)	C1—C2—C9—C8	-2.8 (4)
C4—C5—C6—C7	0.7 (4)	C3—C2—C9—C4	-0.7 (2)
C5—C6—C7—C8	0.3 (4)	C1—C2—C9—C4	-178.9 (2)
C6—C7—C8—C9	-1.1 (3)	C2—C3—N1—C4	-0.8 (3)
C7—C8—C9—C4	0.9 (3)	C5—C4—N1—C3	-176.0 (2)
C7—C8—C9—C2	-174.8 (2)	C9—C4—N1—C3	0.4 (2)

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1A···O1 <sup>i</sup>	0.94 (3)	1.92 (3)	2.831 (2)	165 (3)

Symmetry code: (i)  $-x+3/2, y-1, z+1/2$ .