

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# 3,5-Dimethylpyrazolium 3,5-dinitro-salicylate

Shuaishuai Wei, Shouwen Jin,\* Zhaofeng Hu, Yong Zhou and Yingping Zhou

Tianmu College of Zhejiang A &amp; F University, Lin'an 311300, People's Republic of China

Correspondence e-mail: shouwenjin@yahoo.cn


Received 10 September 2012; accepted 6 October 2012

 Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.005$  Å;  $R$  factor = 0.047;  $wR$  factor = 0.105; data-to-parameter ratio = 11.1.

In the title molecular salt,  $\text{C}_5\text{H}_9\text{N}_2^+\cdot\text{C}_7\text{H}_3\text{N}_2\text{O}_7^-$ , the roughly planar anion (r.m.s. deviation = 0.120 Å) has been deprotonated at the phenol group. An intramolecular  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bond in the anion generates an  $S(6)$  ring. In the crystal, the components are linked by cation-to-anion  $\text{N}-\text{H}\cdots\text{O}$  and  $\text{N}-\text{H}\cdots(\text{O},\text{O})$  hydrogen bonds, generating [010] double chains. Weak  $\text{C}-\text{H}\cdots\text{O}$  interactions consolidate the packing.

## Related literature

For a related structure and background to hydrogen-bonding interactions, see: Jin *et al.* (2010). For another related structure, see: Smith *et al.* (2011).



## Experimental

### Crystal data

 $\text{C}_5\text{H}_9\text{N}_2^+\cdot\text{C}_7\text{H}_3\text{N}_2\text{O}_7^-$ 
 $M_r = 324.26$ 

 Monoclinic,  $P2_1$ 
 $a = 8.1183$  (7) Å

 $b = 6.0636$  (5) Å

 $c = 14.1453$  (11) Å

 $\beta = 91.904$  (1)°

 $V = 695.93$  (10) Å<sup>3</sup>
 $Z = 2$ 

 Mo  $K\alpha$  radiation

 $\mu = 0.13$  mm<sup>-1</sup>
 $T = 293$  K

 $0.40 \times 0.27 \times 0.11$  mm

### Data collection

Bruker SMART CCD

diffractometer

Absorption correction: multi-scan

(SADABS; Bruker, 2002)

 $T_{\min} = 0.959$ ,  $T_{\max} = 0.986$ 

3523 measured reflections

2301 independent reflections

 1659 reflections with  $I > 2\sigma(I)$ 
 $R_{\text{int}} = 0.040$ 

### Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.047$ 
 $wR(F^2) = 0.105$ 
 $S = 1.02$ 

2301 reflections

208 parameters

1 restraint

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.16$  e Å<sup>-3</sup>
 $\Delta\rho_{\text{min}} = -0.18$  e Å<sup>-3</sup>
**Table 1**

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1}\cdots\text{O7}^{\text{i}}$	0.86	2.09	2.859 (4)	148
$\text{N1}-\text{H1}\cdots\text{O1}^{\text{i}}$	0.86	2.16	2.809 (4)	132
$\text{N2}-\text{H2}\cdots\text{O3}$	0.86	1.88	2.684 (3)	156
$\text{O2}-\text{H2A}\cdots\text{O1}$	0.82	1.72	2.481 (3)	154
$\text{C1}-\text{H1A}\cdots\text{O7}^{\text{i}}$	0.96	2.32	3.166 (5)	147
$\text{C5}-\text{H5B}\cdots\text{O4}^{\text{ii}}$	0.96	2.49	3.414 (5)	160
$\text{C10}-\text{H10}\cdots\text{O6}^{\text{iii}}$	0.93	2.48	3.379 (4)	164

 Symmetry codes: (i)  $-x+1, y+\frac{1}{2}, -z+1$ ; (ii)  $-x, y-\frac{3}{2}, -z+1$ ; (iii)  $-x+1, y+\frac{1}{2}, -z+2$ .

Data collection: *SMART* (Bruker, 2002); cell refinement: *SAINTE* (Bruker, 2002); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

We gratefully acknowledge the financial support of the Education Office Foundation of Zhejiang Province (project No. Y201017321) and the innovation project of Zhejiang A & F University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6958).

## References

- Bruker (2002). *SADABS*, *SMART* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Jin, S. W., Zhang, W. B., Liu, L., Gao, H. F., Wang, D. Q., Chen, R. P. & Xu, X. L. (2010). *J. Mol. Struct.* **975**, 128–136.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Smith, G., Wermuth, U. D., Healy, P. C. & White, J. M. (2011). *J. Chem. Crystallogr.* **41**, 1649–1662.

## supporting information

*Acta Cryst.* (2012). E68, o3117 [doi:10.1107/S1600536812041906]

## 3,5-Dimethylpyrazolium 3,5-dinitrosalicylate

Shuaishuai Wei, Shouwen Jin, Zhaofeng Hu, Yong Zhou and Yingping Zhou

### S1. Comment

Organic acid-base adducts based on hydrogen bonding are a research field receiving great attention in recent years. As an extension of our study concentrating on hydrogen bonded assembly of organic acid and organic base (Jin *et al.*, 2010), herein we report the crystal structure of 3,5-dimethylpyrazolium 3,5-dinitrosalicylate.

The crystal of the title compound of the formula  $C_{12}H_{12}N_4O_7$  was obtained by recrystallization of the mixture of 3,5-dimethylpyrazole and 3,5-dinitrosalicylic acid from the MeOH solution.

In this case (Fig. 1) it is the phenol H that has been deprotonated. The C—O distance 1.284 (4) Å concerning the phenolate is similar to the proton transfer compound bearing the 3,5-dinitrosalicylate in which only the phenol group has been deprotonated (Smith *et al.*, 2011). The C—O distances O(2)—C(6), 1.300 (4) Å, O(3)—C(6), 1.223 (4) Å; in the COOH show characteristic C—O, and C=O distances which are also confirming the reliability of adding H atoms experimentally by different electron density onto O atoms.

One anion is bonded to one cation *via* N—H $\cdots$ O, and CH<sub>3</sub>—O associations to form a heteroadduct.

For the presence of these interactions, there are close joint motifs with descriptors of  $R_21(6)$ , and  $R_12(6)$ . The usual intramolecular hydrogen bond is found between the phenolate and the carboxyl group to exhibit a  $S_11(6)$  graph. The heteroadducts were linked together by the CH<sub>3</sub>—O association between the methyl group of the pyrazole and the 5-nitro group with C—O distance of 3.415 Å to form one-dimensional chain running along the direction that made an angle of *ca* 60° with the *a* axis (Fig. 2). The chains were further stacked along the direction that is perpendicular with its extending direction *via* the interchain N— $\pi$  interaction between the 5-nitro group and the phenyl ring of the anion with N—Cg distance of 3.236 Å to form two-dimensional sheet extending parallel to the *ab* plane. The sheets were further stacked along the *c* axis direction by the CH—O, N—H $\cdots$ O, and O—H $\cdots$ O associations to form three-dimensional ABAB layer network structure. Herein the chains at adjacent layers intersect at an angle of *ca* 120° with each other.


### S2. Experimental

A solution of 3,5-dimethyl pyrazole (19.2 mg, 0.2 mmol) in 5 ml of MeOH was added to a MeOH solution (6 ml) containing 3,5-dinitrosalicylic acid (22.8 mg, 0.1 mmol) under continuous stirring. The solution was stirred for about 1 h at room temperature, then the solution was filtered into a test tube. The solution was left standing at room temperature for several days, yellow blocks were isolated after slow evaporation of the solution in air at ambient temperature. The crystals were collected and dried in air to give the title compound.

### S3. Refinement


The absolute structure could not be determined in the present refinement. Hydrogen atoms attached to the C atoms were placed in calculated positions with  $d(C—H) = 0.93–0.96$  Å. Positions of the hydrogen atoms at the NH, and COOH groups were located from the Fourier difference syntheses and refined independently. All  $U_{iso}$  values were restrained on

$U_{eq}$  values of the parent atoms.



**Figure 1**

The structure of the title compound, showing displacement ellipsoids drawn at the 30% probability level.



**Figure 2**

The one-dimensional chain formed through the CH<sub>3</sub>—O interaction.

### 3,5-Dimethylpyrazolium 3,5-dinitrosalicylate

#### Crystal data

$C_5H_9N_2^+ \cdot C_7H_3N_2O_7^-$

$M_r = 324.26$

Monoclinic,  $P2_1$

$a = 8.1183$  (7) Å

$b = 6.0636$  (5) Å

$c = 14.1453$  (11) Å

$\beta = 91.904$  (1)°

$V = 695.93$  (10) Å<sup>3</sup>

$Z = 2$

$F(000) = 336$

$D_x = 1.547$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 1025 reflections

$\theta = 2.5$ – $22.6$ °

$\mu = 0.13$  mm<sup>-1</sup>

$T = 293$  K

Block, colorless

$0.40 \times 0.27 \times 0.11$  mm

#### Data collection

Bruker SMART CCD  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\omega$  scans

Absorption correction: multi-scan  
(*SADABS*; Bruker, 2002)

$T_{min} = 0.959$ ,  $T_{max} = 0.986$

3523 measured reflections

2301 independent reflections

1659 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.040$   
 $\theta_{\text{max}} = 25.0^\circ$ ,  $\theta_{\text{min}} = 2.5^\circ$   
 $h = -9 \rightarrow 9$

$k = -7 \rightarrow 7$   
 $l = -16 \rightarrow 12$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.047$   
 $wR(F^2) = 0.105$   
 $S = 1.02$   
 2301 reflections  
 208 parameters  
 1 restraint  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0405P)^2]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\text{max}} < 0.001$   
 $\Delta\rho_{\text{max}} = 0.16 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.18 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.2255 (4)	0.7148 (5)	0.2340 (2)	0.0463 (7)
H1	0.2726	0.8381	0.2218	0.056*
N2	0.2198 (3)	0.6193 (5)	0.31994 (19)	0.0469 (8)
H2	0.2635	0.6724	0.3713	0.056*
C6	0.3512 (4)	0.7229 (6)	0.5523 (2)	0.0405 (8)
N3	0.2184 (3)	1.3159 (5)	0.7707 (2)	0.0487 (7)
N4	0.5550 (4)	0.7118 (5)	0.8914 (2)	0.0484 (8)
O1	0.5414 (3)	0.5277 (4)	0.70183 (15)	0.0454 (6)
O2	0.4254 (3)	0.5367 (4)	0.53738 (15)	0.0581 (7)
H2A	0.4770	0.4990	0.5855	0.087*
O3	0.2657 (3)	0.8102 (4)	0.48999 (14)	0.0493 (6)
O4	0.1417 (3)	1.4008 (4)	0.70451 (18)	0.0632 (7)
O5	0.2271 (3)	1.3968 (5)	0.85007 (18)	0.0689 (8)
O6	0.5255 (4)	0.7837 (5)	0.96945 (17)	0.0833 (10)
O7	0.6547 (4)	0.5660 (5)	0.87978 (17)	0.0698 (8)
C1	0.1300 (5)	0.6449 (8)	0.0686 (2)	0.0717 (13)
H1A	0.1787	0.7868	0.0584	0.108*
H1B	0.0155	0.6492	0.0495	0.108*
H1C	0.1854	0.5360	0.0320	0.108*
C2	0.1462 (4)	0.5868 (6)	0.1713 (2)	0.0464 (9)
C3	0.0892 (4)	0.4066 (7)	0.2187 (3)	0.0544 (10)
H3	0.0298	0.2892	0.1925	0.065*

C4	0.1370 (4)	0.4321 (6)	0.3134 (2)	0.0464 (9)
C5	0.1105 (4)	0.2906 (7)	0.3972 (3)	0.0625 (11)
H5A	0.2148	0.2570	0.4278	0.094*
H5B	0.0572	0.1562	0.3774	0.094*
H5C	0.0422	0.3672	0.4406	0.094*
C12	0.4688 (4)	0.7102 (5)	0.7195 (2)	0.0353 (8)
C7	0.3732 (3)	0.8231 (6)	0.64774 (19)	0.0339 (7)
C8	0.2969 (4)	1.0188 (6)	0.6644 (2)	0.0376 (8)
H8	0.2396	1.0905	0.6154	0.045*
C9	0.3041 (4)	1.1108 (6)	0.7533 (2)	0.0368 (8)
C10	0.3900 (4)	1.0088 (6)	0.8269 (2)	0.0396 (8)
H10	0.3935	1.0711	0.8870	0.048*
C11	0.4701 (4)	0.8142 (6)	0.8101 (2)	0.0366 (7)

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
N1	0.0503 (17)	0.0446 (18)	0.0438 (17)	−0.0035 (14)	−0.0020 (13)	−0.0021 (14)
N2	0.0493 (18)	0.051 (2)	0.0397 (16)	−0.0042 (16)	−0.0060 (13)	−0.0055 (16)
C6	0.0382 (18)	0.048 (2)	0.0356 (19)	−0.0017 (17)	0.0045 (16)	−0.0016 (17)
N3	0.0479 (17)	0.0412 (19)	0.057 (2)	0.0007 (16)	0.0029 (15)	−0.0032 (18)
N4	0.064 (2)	0.044 (2)	0.0372 (18)	−0.0012 (16)	−0.0046 (15)	0.0033 (15)
O1	0.0544 (14)	0.0412 (14)	0.0404 (13)	0.0081 (12)	−0.0011 (10)	−0.0056 (11)
O2	0.0701 (17)	0.0628 (18)	0.0408 (14)	0.0203 (15)	−0.0077 (11)	−0.0155 (13)
O3	0.0554 (13)	0.0590 (16)	0.0329 (12)	0.0050 (13)	−0.0072 (11)	0.0004 (12)
O4	0.0679 (17)	0.0510 (17)	0.0699 (17)	0.0183 (14)	−0.0112 (14)	0.0030 (15)
O5	0.088 (2)	0.0582 (18)	0.0602 (17)	0.0083 (16)	0.0045 (14)	−0.0181 (15)
O6	0.144 (3)	0.073 (2)	0.0317 (15)	0.033 (2)	−0.0088 (15)	−0.0034 (14)
O7	0.088 (2)	0.070 (2)	0.0509 (15)	0.0294 (18)	−0.0100 (14)	0.0072 (15)
C1	0.079 (3)	0.093 (3)	0.043 (2)	−0.017 (3)	−0.005 (2)	−0.009 (2)
C2	0.044 (2)	0.052 (3)	0.0424 (19)	0.0036 (19)	−0.0059 (16)	−0.0103 (19)
C3	0.050 (2)	0.056 (3)	0.057 (2)	−0.006 (2)	−0.0065 (18)	−0.023 (2)
C4	0.0363 (19)	0.042 (2)	0.061 (2)	0.0018 (18)	0.0040 (16)	−0.0034 (19)
C5	0.061 (2)	0.057 (3)	0.069 (3)	−0.001 (2)	−0.0008 (19)	0.010 (2)
C12	0.0346 (18)	0.038 (2)	0.0338 (18)	−0.0056 (16)	0.0023 (14)	0.0013 (16)
C7	0.0348 (16)	0.0367 (19)	0.0302 (16)	−0.0059 (16)	0.0012 (13)	0.0009 (15)
C8	0.0393 (19)	0.039 (2)	0.0346 (18)	−0.0019 (17)	−0.0034 (14)	0.0036 (16)
C9	0.0394 (19)	0.0310 (19)	0.0399 (18)	−0.0045 (16)	0.0016 (14)	−0.0010 (16)
C10	0.049 (2)	0.040 (2)	0.0299 (17)	−0.0064 (18)	0.0020 (15)	−0.0014 (16)
C11	0.0403 (18)	0.0381 (19)	0.0312 (17)	−0.0023 (18)	−0.0024 (13)	0.0046 (16)

*Geometric parameters (Å, °)*

N1—C2	1.329 (4)	C1—H1B	0.9600
N1—N2	1.348 (4)	C1—H1C	0.9600
N1—H1	0.8600	C2—C3	1.371 (5)
N2—C4	1.321 (4)	C3—C4	1.391 (5)
N2—H2	0.8600	C3—H3	0.9300

C6—O3	1.224 (4)	C4—C5	1.484 (5)
C6—O2	1.301 (4)	C5—H5A	0.9600
C6—C7	1.486 (4)	C5—H5B	0.9600
N3—O4	1.221 (3)	C5—H5C	0.9600
N3—O5	1.226 (3)	C12—C11	1.427 (4)
N3—C9	1.449 (4)	C12—C7	1.431 (4)
N4—O7	1.213 (4)	C7—C8	1.363 (4)
N4—O6	1.218 (3)	C8—C9	1.375 (4)
N4—C11	1.460 (4)	C8—H8	0.9300
O1—C12	1.283 (4)	C9—C10	1.379 (4)
O2—H2A	0.8200	C10—C11	1.372 (4)
C1—C2	1.496 (5)	C10—H10	0.9300
C1—H1A	0.9600		
C2—N1—N2	108.7 (3)	N2—C4—C3	106.7 (3)
C2—N1—H1	125.6	N2—C4—C5	121.9 (3)
N2—N1—H1	125.6	C3—C4—C5	131.4 (3)
C4—N2—N1	109.8 (3)	C4—C5—H5A	109.5
C4—N2—H2	125.1	C4—C5—H5B	109.5
N1—N2—H2	125.1	H5A—C5—H5B	109.5
O3—C6—O2	120.9 (3)	C4—C5—H5C	109.5
O3—C6—C7	121.7 (3)	H5A—C5—H5C	109.5
O2—C6—C7	117.4 (3)	H5B—C5—H5C	109.5
O4—N3—O5	123.1 (3)	O1—C12—C11	124.5 (3)
O4—N3—C9	117.9 (3)	O1—C12—C7	121.0 (3)
O5—N3—C9	119.0 (3)	C11—C12—C7	114.4 (3)
O7—N4—O6	122.4 (3)	C8—C7—C12	122.2 (3)
O7—N4—C11	120.2 (3)	C8—C7—C6	118.1 (3)
O6—N4—C11	117.4 (3)	C12—C7—C6	119.7 (3)
C6—O2—H2A	109.5	C7—C8—C9	120.4 (3)
C2—C1—H1A	109.5	C7—C8—H8	119.8
C2—C1—H1B	109.5	C9—C8—H8	119.8
H1A—C1—H1B	109.5	C8—C9—C10	120.8 (3)
C2—C1—H1C	109.5	C8—C9—N3	119.8 (3)
H1A—C1—H1C	109.5	C10—C9—N3	119.4 (3)
H1B—C1—H1C	109.5	C11—C10—C9	119.1 (3)
N1—C2—C3	107.6 (3)	C11—C10—H10	120.5
N1—C2—C1	122.4 (4)	C9—C10—H10	120.5
C3—C2—C1	130.0 (3)	C10—C11—C12	123.1 (3)
C2—C3—C4	107.2 (3)	C10—C11—N4	116.3 (3)
C2—C3—H3	126.4	C12—C11—N4	120.6 (3)
C4—C3—H3	126.4		
C2—N1—N2—C4	-0.4 (4)	C7—C8—C9—C10	0.8 (5)
N2—N1—C2—C3	0.0 (4)	C7—C8—C9—N3	-178.3 (3)
N2—N1—C2—C1	-179.8 (3)	O4—N3—C9—C8	0.0 (4)
N1—C2—C3—C4	0.3 (4)	O5—N3—C9—C8	-179.4 (3)
C1—C2—C3—C4	-179.9 (4)	O4—N3—C9—C10	-179.1 (3)

N1—N2—C4—C3	0.6 (3)	O5—N3—C9—C10	1.5 (4)
N1—N2—C4—C5	-179.9 (3)	C8—C9—C10—C11	0.6 (5)
C2—C3—C4—N2	-0.6 (4)	N3—C9—C10—C11	179.7 (3)
C2—C3—C4—C5	180.0 (4)	C9—C10—C11—C12	-0.2 (4)
O1—C12—C7—C8	-179.0 (3)	C9—C10—C11—N4	-178.4 (3)
C11—C12—C7—C8	2.9 (4)	O1—C12—C11—C10	-179.5 (3)
O1—C12—C7—C6	2.8 (4)	C7—C12—C11—C10	-1.4 (4)
C11—C12—C7—C6	-175.3 (3)	O1—C12—C11—N4	-1.3 (5)
O3—C6—C7—C8	-0.9 (4)	C7—C12—C11—N4	176.7 (3)
O2—C6—C7—C8	179.7 (3)	O7—N4—C11—C10	-165.3 (3)
O3—C6—C7—C12	177.4 (3)	O6—N4—C11—C10	12.4 (4)
O2—C6—C7—C12	-2.1 (4)	O7—N4—C11—C12	16.5 (5)
C12—C7—C8—C9	-2.7 (5)	O6—N4—C11—C12	-165.9 (3)
C6—C7—C8—C9	175.5 (3)		

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1 $\cdots$ O7 <sup>i</sup>	0.86	2.09	2.859 (4)	148
N1—H1 $\cdots$ O1 <sup>i</sup>	0.86	2.16	2.809 (4)	132
N2—H2 $\cdots$ O3	0.86	1.88	2.684 (3)	156
O2—H2A $\cdots$ O1	0.82	1.72	2.481 (3)	154
C1—H1A $\cdots$ O7 <sup>i</sup>	0.96	2.32	3.166 (5)	147
C5—H5B $\cdots$ O4 <sup>ii</sup>	0.96	2.49	3.414 (5)	160
C10—H10 $\cdots$ O6 <sup>iii</sup>	0.93	2.48	3.379 (4)	164

Symmetry codes: (i)  $-x+1, y+1/2, -z+1$ ; (ii)  $-x, y-3/2, -z+1$ ; (iii)  $-x+1, y+1/2, -z+2$ .