

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(Acetato- κ O)(acetato- κ O, O')bis(1,3-diazinane-2-thione- κ S)cadmium(II)

Rashid Mahmood,^a Saima Ghulam Hussain,^b Mohammed Fettouhi,^c Anvarhusein A. Isab^c and Saeed Ahmad^{b*}

^aDivision of Science and Technology, University of Education, Township, Lahore, Pakistan, ^bDepartment of Chemistry, University of Engineering and Technology, Lahore 54890, Pakistan, and ^cDepartment of Chemistry, King Fahd University of Petroleum and Minerals, Dhahran 31261, Saudi Arabia

Correspondence e-mail: saeed_a786@hotmail.com


Received 29 August 2012; accepted 6 October 2012

Key indicators: single-crystal X-ray study; $T = 294$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.033; wR factor = 0.082; data-to-parameter ratio = 22.7.

In the title complex, $[\text{Cd}(\text{CH}_3\text{COO})_2(\text{C}_4\text{H}_8\text{N}_2\text{S})_2]$, the Cd^{II} cation is coordinated by three acetate O atoms and two S atoms of Diaz [Diaz = 1,3-diazinane-2-thione = 3,4,5,6-tetrahydropyrimidine-2(1*H*)-thione]. The Cd^{II} coordination is augmented by one considerably longer Cd—O bond of 2.782 (3) Å to a carboxylate O atom. The resulting coordination polyhedron around the Cd^{II} cations can be described as a highly distorted octahedron. The Diaz ligand and the acetate anions are linked by N—H...O hydrogen-bonding interactions.

Related literature

For crystal structures of Cd^{II} complexes of thiones, see: Ahmad *et al.* (2011, 2012); Altaf *et al.* (2011); Beheshti *et al.* (2007); Lobana *et al.* (2008); Nawaz *et al.* (2010); Moloto *et al.* (2003, 2007); Wang *et al.* (2002); Wazeer *et al.* (2007). For van der Waals radii, see: Bondi (1964).



Experimental

Crystal data

$[\text{Cd}(\text{C}_2\text{H}_3\text{O}_2)_2(\text{C}_4\text{H}_8\text{N}_2\text{S})_2]$
 $M_r = 462.86$
 Triclinic, $P\bar{1}$
 $a = 8.6930$ (17) Å
 $b = 10.175$ (2) Å
 $c = 12.203$ (2) Å
 $\alpha = 97.452$ (4)°
 $\beta = 100.683$ (4)°

$\gamma = 111.610$ (3)°
 $V = 962.6$ (3) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 1.37$ mm⁻¹
 $T = 294$ K
 $0.19 \times 0.18 \times 0.10$ mm

Data collection

Bruker SMART APEX area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\text{min}} = 0.781$, $T_{\text{max}} = 0.875$

13173 measured reflections
 4775 independent reflections
 3705 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.024$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.033$
 $wR(F^2) = 0.082$
 $S = 1.02$
 4775 reflections

210 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.47$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.43$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1...O3	0.86	1.94	2.779 (3)	167
N3—H3...O4	0.86	2.10	2.897 (4)	154
N2—H2...O2 ⁱ	0.86	2.01	2.829 (3)	160
N4—H4...O1 ⁱⁱ	0.86	2.03	2.836 (3)	156

Symmetry codes: (i) $x + 1, y, z$; (ii) $-x + 1, -y, -z + 1$.

Data collection: SMART (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997); software used to prepare material for publication: publCIF (Westrip, 2010).

The authors gratefully acknowledge King Fahd University of Petroleum and Minerals, Dhahran, Saudi Arabia, for providing X-ray facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NC2292).

References

- Ahmad, S., Altaf, M., Stoeckli-Evans, H., Isab, A. A., Malik, M. R., Ali, S. & Shuja, S. (2011). *J. Chem. Crystallogr.* **41**, 1099–1104.
 Ahmad, S., Amir, Q., Naz, G., Fettouhi, M., Isab, A. A., Rüffer, T. & Lang, H. (2012). *J. Chem. Crystallogr.* **42**, 615–620.
 Altaf, M., Stoeckli-Evans, H., Murtaza, G., Isab, A. A., Ahmad, S. & Shaheen, M. A. (2011). *J. Struct. Chem.* **52**, 625–630.
 Beheshti, A., Clegg, W., Dale, S. H. & Hyvadi, R. (2007). *Inorg. Chim. Acta*, **360**, 2967–2972.
 Bondi, A. (1964). *J. Phys. Chem.* **68**, 441–451.
 Bruker (2007). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Lobana, T. S., Sharma, R., Sharma, R., Sultana, R. & Butcher, R. J. (2008). *Z. Anorg. Allg. Chem.* **634**, 718–723.

- Moloto, M. J., Malik, M. A., O'Brien, P., Motevalli, M. & Kolawole, G. A. (2003). *Polyhedron*, **22**, 595–603.
- Moloto, N., Revaprasadu, N., Moloto, M. J., O'Brien, P. & Helliwell, M. (2007). *Polyhedron*, **26**, 3947–3955.
- Nawaz, S., Sadaf, S., Fettouhi, M., Fazal, A. & Ahmad, S. (2010). *Acta Cryst. E* **66**, m950.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Wang, X. Q., Yu, W. T., Xu, D., Lu, M. K. & Yuan, D. R. (2002). *Acta Cryst. C* **58**, m336–m337.
- Wazeer, M. I. M., Isab, A. A. & Fettouhi, M. (2007). *Polyhedron*, **26**, 1725–1730.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2012). E68, m1352–m1353 [doi:10.1107/S1600536812041852]

(Acetato- κ O)(acetato- κ O,O')bis(1,3-diazinane-2-thione- κ S)cadmium(II)

Rashid Mahmood, Saima Ghulam Hussain, Mohammed Fettouhi, Anvarhusein A. Isab and Saeed Ahmad

S1. Comment


The structural reports on cadmium(II) complexes of thiones describe that in most of the cases, the cadmium(II) complexes exist as neutral monomeric (Ahmad *et al.* 2011, 2012; Beheshti *et al.*, 2007; Lobana *et al.* 2008; Nawaz *et al.*, 2010; Wazeer *et al.*, 2007) or polymeric (Moloto *et al.*, 2007; Wang *et al.*, 2002) with cadmium atom possessing a tetrahedral or distorted octahedral coordination environment respectively. In continuation of our studies on the structural chemistry of cadmium(II) complexes with thione ligands, we report here the crystal structure of the title compound. The crystal structure of the title complex (Figure 1) consists of discrete monomeric species in which the Cd(II) ion is bound to two sulfur Diaz atoms, two oxygen atoms belonging to a chelating acetate and a third oxygen atom of a second acetate anion. The Cd—S bond distances are 2.5787 (8) and 2.529 (1) Å respectively while the Cd—O bond distances are in the range of 2.266 (2)- 2.421 (2) Å. In addition, a secondary bonding interaction takes place with the second oxygen of one acetate anion with a distance of 2.782 (3) Å. The van der Waals radii for cadmium and oxygen are 1.58 and 1.52 Å respectively (Bondi, 1964). Considering the secondary bonding, the geometry could be considered as highly distorted octahedral. The S—Cd—S bond angle is 97.02 (3) °, the O—Cd—O bond angle for the bidentate acetate ligand is 53.86 (7) ° and the S—Cd—O bond angles vary from 92.37 (5) ° to 144.34 (6) °. The Cd—S and Cd—O bond lengths are in agreement with those reported for related compounds. Two intramolecular (N—H···O) hydrogen bonds are found between the Diaz ligands and acetate anions. The discrete complex molecules are also linked by intermolecular N—H···O hydrogen bonding (Table 1).

S2. Experimental

The title complex was prepared by adding 0.24 g (2.0 mmol) of 1,3-diazinane-2-thione in 15 ml of methanol to an aqueous solution (5 ml) of 0.26 g (1.0 mmol) cadmium sulfate in water followed by addition of 2 equivalents of sodium acetate in 10 ml water and stirring the mixture for 30 minutes. The colorless solution was filtered and the filtrate was kept at room temperature for crystallization. As a result, white crystalline product was obtained, that was washed with methanol and dried.

S3. Refinement

All H atoms were placed in calculated positions and were refined isotropic with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$ (1.5 for methyl H atoms) using a riding model with C—H distances of 0.96 Å and 0.97 Å and N—H distances of 0.86 Å.

**Figure 1**

Molecular structure of the title compound showing the atomic numbering scheme. Displacement ellipsoids are drawn at the 30% probability level and dashed lines indicate intramolecular hydrogen bonding and the secondary Cd \cdots O interaction.

(Acetato- κ O)(acetato- κ O,O')bis(1,3-diazinane-2-thione- κ S)cadmium(II)

Crystal data

[Cd(C₂H₃O₂)₂(C₄H₈N₂S)₂]

M_r = 462.86

Triclinic, $P\bar{1}$

Hall symbol: -P 1

a = 8.6930 (17) Å

b = 10.175 (2) Å

c = 12.203 (2) Å

α = 97.452 (4)°

β = 100.683 (4)°

γ = 111.610 (3)°

V = 962.6 (3) Å³

Z = 2

$F(000)$ = 468

D_x = 1.597 Mg m⁻³

Mo $K\alpha$ radiation, λ = 0.71073 Å

Cell parameters from 13173 reflections

θ = 1.7–28.3°

μ = 1.37 mm⁻¹

T = 294 K

Block, colourless

0.19 × 0.18 × 0.10 mm

Data collection

Bruker SMART APEX area-detector
diffractometer

Radiation source: normal-focus sealed tube

Graphite monochromator

ω scans

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 1996)

T_{\min} = 0.781, T_{\max} = 0.875

13173 measured reflections

4775 independent reflections

3705 reflections with $I > 2\sigma(I)$

R_{int} = 0.024

θ_{max} = 28.3°, θ_{min} = 1.7°

h = -11→11

k = -13→13

l = -16→16

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.033$
 $wR(F^2) = 0.082$
 $S = 1.02$
 4775 reflections
 210 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0408P)^2 + 0.0579P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.47 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.43 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cd1	0.71521 (3)	0.17566 (2)	0.769239 (17)	0.05732 (9)
S1	0.98165 (9)	0.11886 (8)	0.77361 (6)	0.05721 (17)
S2	0.73948 (12)	0.30728 (12)	0.60691 (8)	0.0827 (3)
O1	0.5230 (3)	-0.0768 (2)	0.69890 (17)	0.0679 (5)
O2	0.4199 (3)	0.0743 (2)	0.76067 (19)	0.0713 (6)
O3	0.7698 (3)	0.2558 (2)	0.96021 (19)	0.0753 (6)
O4	0.6916 (4)	0.4146 (3)	0.8923 (2)	0.1107 (10)
N1	1.0593 (3)	0.1951 (2)	1.00174 (18)	0.0515 (5)
H1	0.9615	0.2007	0.9917	0.062*
N2	1.2447 (3)	0.1265 (3)	0.92374 (19)	0.0569 (6)
H2	1.2731	0.0989	0.8638	0.068*
N3	0.4916 (3)	0.3743 (3)	0.6630 (2)	0.0655 (6)
H3	0.5640	0.4152	0.7283	0.079*
N4	0.4398 (3)	0.2662 (3)	0.4764 (2)	0.0677 (7)
H4	0.4757	0.2314	0.4233	0.081*
C1	1.1027 (3)	0.1503 (3)	0.9105 (2)	0.0464 (5)
C2	1.1658 (4)	0.2360 (3)	1.1187 (2)	0.0595 (7)
H2A	1.1236	0.1581	1.1579	0.071*
H2B	1.1597	0.3216	1.1596	0.071*
C3	1.3496 (4)	0.2664 (3)	1.1175 (3)	0.0623 (7)
H3A	1.4004	0.3573	1.0945	0.075*
H3B	1.4150	0.2745	1.1935	0.075*
C4	1.3539 (4)	0.1449 (3)	1.0350 (3)	0.0612 (7)
H4A	1.4702	0.1670	1.0292	0.073*
H4B	1.3143	0.0559	1.0624	0.073*

C5	0.5411 (4)	0.3152 (3)	0.5810 (2)	0.0593 (7)
C6	0.2707 (4)	0.2682 (4)	0.4463 (3)	0.0800 (10)
H6A	0.1956	0.1855	0.3854	0.096*
H6B	0.2784	0.3556	0.4196	0.096*
C7	0.2012 (5)	0.2632 (5)	0.5474 (3)	0.0978 (13)
H7A	0.1735	0.1679	0.5645	0.117*
H7B	0.0962	0.2783	0.5305	0.117*
C8	0.3232 (4)	0.3746 (4)	0.6498 (3)	0.0768 (9)
H8A	0.3282	0.4695	0.6413	0.092*
H8B	0.2835	0.3544	0.7174	0.092*
C9	0.4005 (3)	-0.0511 (3)	0.7195 (2)	0.0561 (6)
C10	0.2272 (4)	-0.1727 (4)	0.6978 (4)	0.0892 (11)
H10A	0.2125	-0.2038	0.7676	0.134*
H10B	0.2183	-0.2523	0.6416	0.134*
H10C	0.1401	-0.1395	0.6704	0.134*
C11	0.7493 (3)	0.3726 (3)	0.9725 (2)	0.0574 (7)
C12	0.7991 (5)	0.4611 (3)	1.0914 (3)	0.0779 (10)
H12A	0.9214	0.5038	1.1188	0.117*
H12B	0.7504	0.4001	1.1406	0.117*
H12C	0.7573	0.5363	1.0911	0.117*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cd1	0.05297 (13)	0.06353 (14)	0.05545 (14)	0.02626 (10)	0.00898 (9)	0.01124 (9)
S1	0.0544 (4)	0.0699 (4)	0.0480 (4)	0.0297 (3)	0.0097 (3)	0.0057 (3)
S2	0.0714 (5)	0.1273 (8)	0.0748 (6)	0.0542 (6)	0.0299 (4)	0.0473 (5)
O1	0.0557 (11)	0.0753 (13)	0.0716 (13)	0.0327 (10)	0.0119 (10)	-0.0007 (10)
O2	0.0577 (12)	0.0688 (13)	0.0832 (15)	0.0238 (10)	0.0231 (11)	-0.0012 (11)
O3	0.0630 (13)	0.0762 (14)	0.0839 (15)	0.0388 (11)	0.0061 (11)	-0.0071 (11)
O4	0.164 (3)	0.0721 (16)	0.0651 (16)	0.0344 (17)	-0.0127 (16)	0.0102 (12)
N1	0.0450 (11)	0.0592 (13)	0.0509 (13)	0.0223 (10)	0.0126 (9)	0.0096 (10)
N2	0.0503 (13)	0.0674 (14)	0.0554 (13)	0.0291 (11)	0.0123 (10)	0.0068 (11)
N3	0.0695 (16)	0.0747 (16)	0.0498 (14)	0.0336 (13)	0.0068 (11)	0.0033 (12)
N4	0.0780 (17)	0.0855 (18)	0.0496 (14)	0.0462 (15)	0.0164 (12)	0.0077 (12)
C1	0.0436 (13)	0.0442 (13)	0.0503 (14)	0.0163 (11)	0.0129 (10)	0.0093 (11)
C2	0.0680 (18)	0.0619 (17)	0.0464 (15)	0.0261 (14)	0.0098 (13)	0.0115 (12)
C3	0.0562 (17)	0.0582 (16)	0.0622 (18)	0.0188 (13)	0.0008 (13)	0.0114 (13)
C4	0.0504 (15)	0.0641 (17)	0.0674 (18)	0.0255 (14)	0.0046 (13)	0.0162 (14)
C5	0.0674 (18)	0.0646 (17)	0.0522 (16)	0.0305 (15)	0.0159 (14)	0.0206 (13)
C6	0.078 (2)	0.100 (3)	0.0587 (19)	0.045 (2)	0.0041 (16)	-0.0017 (17)
C7	0.071 (2)	0.147 (4)	0.068 (2)	0.046 (2)	0.0096 (18)	0.000 (2)
C8	0.078 (2)	0.093 (2)	0.066 (2)	0.0462 (19)	0.0191 (16)	0.0012 (17)
C9	0.0519 (15)	0.0685 (18)	0.0475 (15)	0.0267 (14)	0.0096 (12)	0.0082 (13)
C10	0.066 (2)	0.077 (2)	0.110 (3)	0.0162 (18)	0.0223 (19)	0.004 (2)
C11	0.0501 (15)	0.0525 (16)	0.0573 (17)	0.0102 (12)	0.0139 (12)	0.0025 (13)
C12	0.093 (2)	0.0612 (19)	0.0603 (19)	0.0145 (17)	0.0210 (17)	-0.0023 (15)

Geometric parameters (Å, °)

Cd1—O3	2.266 (2)	C2—C3	1.515 (4)
Cd1—O2	2.364 (2)	C2—H2A	0.9700
Cd1—O1	2.421 (2)	C2—H2B	0.9700
Cd1—S2	2.5291 (10)	C3—C4	1.507 (4)
Cd1—S1	2.5787 (8)	C3—H3A	0.9700
Cd1—O4	2.782 (3)	C3—H3B	0.9700
S1—C1	1.721 (3)	C4—H4A	0.9700
S2—C5	1.728 (3)	C4—H4B	0.9700
O1—C9	1.245 (3)	C6—C7	1.470 (5)
O2—C9	1.247 (3)	C6—H6A	0.9700
O3—C11	1.259 (4)	C6—H6B	0.9700
O4—C11	1.215 (4)	C7—C8	1.492 (5)
N1—C1	1.319 (3)	C7—H7A	0.9700
N1—C2	1.462 (3)	C7—H7B	0.9700
N1—H1	0.8600	C8—H8A	0.9700
N2—C1	1.329 (3)	C8—H8B	0.9700
N2—C4	1.451 (3)	C9—C10	1.505 (4)
N2—H2	0.8600	C10—H10A	0.9600
N3—C5	1.320 (4)	C10—H10B	0.9600
N3—C8	1.444 (4)	C10—H10C	0.9600
N3—H3	0.8600	C11—C12	1.499 (4)
N4—C5	1.322 (4)	C12—H12A	0.9600
N4—C6	1.456 (4)	C12—H12B	0.9600
N4—H4	0.8600	C12—H12C	0.9600
O3—Cd1—O2	89.05 (8)	H3A—C3—H3B	108.3
O3—Cd1—O1	114.48 (8)	N2—C4—C3	109.3 (2)
O2—Cd1—O1	53.86 (7)	N2—C4—H4A	109.8
O3—Cd1—S2	132.00 (7)	C3—C4—H4A	109.8
O2—Cd1—S2	105.07 (6)	N2—C4—H4B	109.8
O1—Cd1—S2	110.65 (6)	C3—C4—H4B	109.8
O3—Cd1—S1	96.66 (5)	H4A—C4—H4B	108.3
O2—Cd1—S1	144.34 (6)	N3—C5—N4	119.3 (3)
O1—Cd1—S1	92.37 (5)	N3—C5—S2	121.1 (2)
S2—Cd1—S1	97.02 (3)	N4—C5—S2	119.5 (2)
O3—Cd1—O4	49.63 (8)	N4—C6—C7	109.1 (3)
O2—Cd1—O4	81.01 (8)	N4—C6—H6A	109.9
O1—Cd1—O4	134.07 (8)	C7—C6—H6A	109.9
S2—Cd1—O4	86.90 (6)	N4—C6—H6B	109.9
S1—Cd1—O4	128.48 (7)	C7—C6—H6B	109.9
C1—S1—Cd1	112.35 (9)	H6A—C6—H6B	108.3
C5—S2—Cd1	98.79 (10)	C6—C7—C8	112.5 (3)
C9—O1—Cd1	91.30 (17)	C6—C7—H7A	109.1
C9—O2—Cd1	93.95 (17)	C8—C7—H7A	109.1
C11—O3—Cd1	105.62 (19)	C6—C7—H7B	109.1
C11—O4—Cd1	81.81 (19)	C8—C7—H7B	109.1

C1—N1—C2	124.6 (2)	H7A—C7—H7B	107.8
C1—N1—H1	117.7	N3—C8—C7	110.3 (3)
C2—N1—H1	117.7	N3—C8—H8A	109.6
C1—N2—C4	122.8 (2)	C7—C8—H8A	109.6
C1—N2—H2	118.6	N3—C8—H8B	109.6
C4—N2—H2	118.6	C7—C8—H8B	109.6
C5—N3—C8	124.0 (2)	H8A—C8—H8B	108.1
C5—N3—H3	118.0	O1—C9—O2	120.9 (3)
C8—N3—H3	118.0	O1—C9—C10	120.2 (3)
C5—N4—C6	123.5 (2)	O2—C9—C10	118.9 (3)
C5—N4—H4	118.3	C9—C10—H10A	109.5
C6—N4—H4	118.3	C9—C10—H10B	109.5
N1—C1—N2	119.2 (2)	H10A—C10—H10B	109.5
N1—C1—S1	122.82 (19)	C9—C10—H10C	109.5
N2—C1—S1	117.95 (19)	H10A—C10—H10C	109.5
N1—C2—C3	110.1 (2)	H10B—C10—H10C	109.5
N1—C2—H2A	109.6	O4—C11—O3	122.5 (3)
C3—C2—H2A	109.6	O4—C11—C12	119.9 (3)
N1—C2—H2B	109.6	O3—C11—C12	117.6 (3)
C3—C2—H2B	109.6	C11—C12—H12A	109.5
H2A—C2—H2B	108.2	C11—C12—H12B	109.5
C4—C3—C2	109.2 (2)	H12A—C12—H12B	109.5
C4—C3—H3A	109.8	C11—C12—H12C	109.5
C2—C3—H3A	109.8	H12A—C12—H12C	109.5
C4—C3—H3B	109.8	H12B—C12—H12C	109.5
C2—C3—H3B	109.8		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1 \cdots O3	0.86	1.94	2.779 (3)	167
N3—H3 \cdots O4	0.86	2.10	2.897 (4)	154
N2—H2 \cdots O2 ⁱ	0.86	2.01	2.829 (3)	160
N4—H4 \cdots O1 ⁱⁱ	0.86	2.03	2.836 (3)	156

Symmetry codes: (i) $x+1, y, z$; (ii) $-x+1, -y, -z+1$.