

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1-Allyl-6-nitro-1*H*-indazoleNabil El Brahmī,^{a*} Mohammed Benchidmi,^a El Mokhtar Essassi,^a Sonia Ladeira^b and Lahcen El Ammari^c^aLaboratoire de Chimie Organique Hétérocyclique URAC21, Faculté des Sciences, Université Mohammed V-Agdal, Avenue Ibn Battouta, BP 1014, Rabat, Morocco,^bLaboratoire de Chimie de Coordination, route de Narbonne, 31077 Toulouse, France, and ^cLaboratoire de Chimie du Solide Appliquée, Faculté des Sciences, Université Mohammed V-Agdal, Avenue Ibn Battouta, BP 1014, Rabat, Morocco
Correspondence e-mail: n_elbrahmi@yahoo.fr


Received 20 October 2012; accepted 9 November 2012

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.034; wR factor = 0.100; data-to-parameter ratio = 15.5.

The fused five- and six-membered rings in the title molecule, $\text{C}_{10}\text{H}_9\text{N}_3\text{O}_2$, are essentially coplanar, the largest deviation from the mean plane being 0.012 (1) Å for the C atom linked to the nitro group. The fused-ring system makes a dihedral angle of 11.34 (6)° with the nitro group, leading to a syn-periplanar conformation. The plane through the atoms forming the allyl group is nearly perpendicular to the indazole fused-ring system, as indicated by the dihedral angle of 73.3 (5)°. In the crystal, each molecule is linked to its symmetry equivalent about the center of inversion by pairs of non-classical C—H...O hydrogen bonds, forming an extended tape motif parallel to the $(\bar{1}\bar{2}\bar{1})$ plane.

Related literature

For the pharmacological and biochemical properties of substituted indazoles, see: Saczewski *et al.* (2008); Jones *et al.* (2009); Bouissane *et al.* (2006). For compounds with similar structures, see: El Brahmī *et al.* (2009, 2011).



Experimental

Crystal data

 $\text{C}_{10}\text{H}_9\text{N}_3\text{O}_2$
 $M_r = 203.20$
 Triclinic, $P\bar{1}$
 $a = 4.3630$ (16) Å
 $b = 8.3245$ (7) Å
 $c = 13.541$ (5) Å

 $\alpha = 95.647$ (2)°
 $\beta = 98.46$ (2)°
 $\gamma = 97.770$ (2)°
 $V = 478.5$ (3) Å³
 $Z = 2$

 Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹
 $T = 296$ K
 $0.38 \times 0.29 \times 0.27$ mm

Data collection

 Bruker Kappa APEXII Quazar
 area-detector diffractometer
 Absorption correction: multi-scan
 (SADABS; Bruker, 2009)
 $T_{\min} = 0.957$, $T_{\max} = 0.997$

 8258 measured reflections
 2109 independent reflections
 1675 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.021$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.100$
 $S = 1.06$
 2109 reflections

 136 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.18$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.20$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C6}-\text{H6}\cdots\text{O1}^{\text{i}}$	0.93	2.51	3.3973 (17)	160
$\text{C8}-\text{H8A}\cdots\text{O1}^{\text{i}}$	0.97	2.53	3.4475 (19)	157
$\text{C2}-\text{H2}\cdots\text{O2}^{\text{ii}}$	0.93	2.66	3.3911 (17)	136

Symmetry codes: (i) $-x, -y, -z$; (ii) $-x + 2, -y + 1, -z$.

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 2012) and Mercury (Macrae *et al.* 2008); software used to prepare material for publication: PLATON (Spek, 2009) and publCIF (Westrip, 2010).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PK2452).

References

- Bouissane, L., El Kazzouli, S., Léonce, S., Pfeiffer, B., Rakib, E. M., Khouili, M. & Guillaumet, G. (2006). *Bioorg. Med. Chem.* **14**, 1078–1088.
- Bruker (2009). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- El Brahmī, N., Benchidmi, M., Essassi, E. M., Ladeira, S. & Ng, S. W. (2011). *Acta Cryst.* **E67**, o3260.
- El Brahmī, N., Mohamed, B., Essassi, E. M., Zouihri, H. & Ng, S. W. (2009). *Acta Cryst.* **E65**, o2320.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Jones, L. H., Allan, G., Barba, O., Burt, C., Corbau, R., Dupont, T., Knöchel, T., Irving, S., Middleton, D. S., Mowbray, C. E., Perros, M., Ringrose, H., Swain, N. A., Webster, R., Westby, M. & Phillips, C. (2009). *J. Med. Chem.* **52**, 1219–1223.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Saczewski, F., Kornicka, A., Rybczyn'ska, A., Hudson, A. L., Miao, S. S., Gdaniec, M., Boblewski, K. & Lehmann, A. (2008). *J. Med. Chem.* **51**, 3599–3608.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2012). E68, o3368 [doi:10.1107/S1600536812046478]

1-Allyl-6-nitro-1*H*-indazole

Nabil El Brahmi, Mohammed BENCHIDMI, El Mokhtar ESSASSI, Sonia LADEIRA and Lahcen EL AMMARI

S1. Comment

Indazole derivatives constitute an exciting heterocyclic family because of their important biological activities. Thus substituted indazoles are generally found to be of pharmaceutical interest in a variety of therapeutic areas (Saczewski *et al.*, 2008, Jones *et al.*, 2009) and with significant cytotoxicities against human (colon and prostate) and murine (leukemia) cell lines (Bouissane *et al.*, 2006).

The plot of the structure of the title compound shows the indazole ring system is linked to a C₃H₅ chain and to a nitro group (Fig.1). The fused-ring system is essentially planar, with a maximum deviation of 0.012 (1) Å for C1. The allyl group is nearly perpendicular to indazole plane as indicated by the torsion angle of C9 C8 N2 N3 = 88.35 (18)°. Moreover, the dihedral angle of 11.34 (6)° between the fused ring system and the nitro group lead to a synperiplanar conformation. The structure of the 1-Allyl-6-nitro-indazole is similar to that reported for the following molecules: 1-allyl-3-chloro-6-nitro-1*H*-indazole and 3-bromo-6-nitro-1-(prop-2-ynyl)-1*H*-indazole (El Brahmi *et al.* 2009, 2011).


In the crystal, each molecule and its symmetry equivalent through the inversion center are linked by C6–H6···O1, C8–H8B···O1 and C2–H2···O2, non-classical hydrogen bonds, which results in an extended tape motif parallel to the plane (-1 2 -1) as shown in Fig.2.

S2. Experimental


6-nitroindazole (5 mmol) and allyl bromide (10 mmol) were reacted in THF (40 ml) in the presence of potassium carbonate (10 mmol) and tetra-*n*-butylammonium bromide (0.5 mmol). The mixture was stirred for 24 h, filtered, and the THF removed under vacuum. The product was separated by chromatography on silica gel with a hexane:ethyl acetate (9:1) solvent system. The compound was obtained as yellow crystals in 68% yield.

S3. Refinement

H atoms were located in a difference map and treated as riding with N—H = 0.86 Å, C—H = 0.93 Å (aromatic), and C—H = 0.97 Å (methylene). with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}$ (aromatic, methylene).

**Figure 1**

Molecular structure of the title compound, with displacement ellipsoids drawn at the 50% probability level. H atoms are represented as small circles.

**Figure 2**

The title molecule and its symmetry equivalent through the inversion center linked by hydrogen bonds and building extended tape motifs parallel to the plane (-1 2 -1).

1-Allyl-6-nitro-1H-indazole

Crystal data

C₁₀H₉N₃O₂ $M_r = 203.20$ Triclinic, $P\bar{1}$

Hall symbol: -P 1

 $a = 4.3630 (16) \text{ \AA}$ $b = 8.3245 (7) \text{ \AA}$ $c = 13.541 (5) \text{ \AA}$ $\alpha = 95.647 (2)^\circ$ $\beta = 98.46 (2)^\circ$ $\gamma = 97.770 (2)^\circ$ $V = 478.5 (3) \text{ \AA}^3$ $Z = 2$ $F(000) = 212$ $D_x = 1.410 \text{ Mg m}^{-3}$ Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 2109 reflections

 $\theta = 3.7\text{--}27.1^\circ$ $\mu = 0.10 \text{ mm}^{-1}$ $T = 296 \text{ K}$

Block, yellow

 $0.38 \times 0.29 \times 0.27 \text{ mm}$

Data collection

Bruker Kappa APEXII Quazar area-detector
diffractometer

Radiation source: microfocus sealed tube

Multilayer optics monochromator

 φ and ω scans

Absorption correction: multi-scan

(SADABS; Bruker, 2009)

 $T_{\min} = 0.957$, $T_{\max} = 0.997$

8258 measured reflections

2109 independent reflections

1675 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.021$ $\theta_{\max} = 27.1^\circ$, $\theta_{\min} = 3.7^\circ$ $h = -5 \rightarrow 5$ $k = -10 \rightarrow 10$ $l = -17 \rightarrow 17$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.034$ $wR(F^2) = 0.100$ $S = 1.06$

2109 reflections

136 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
map

Hydrogen site location: difference Fourier map

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0593P)^2 + 0.0394P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.18 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -0.20 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against all reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on all data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.5250 (3)	0.28071 (13)	0.07294 (8)	0.0285 (3)
C2	0.7333 (3)	0.42576 (14)	0.10756 (9)	0.0336 (3)
H2	0.8493	0.4767	0.0639	0.040*

C3	0.7649 (3)	0.49196 (14)	0.20582 (9)	0.0347 (3)
H3	0.9001	0.5887	0.2297	0.042*
C4	0.5890 (3)	0.41073 (13)	0.26971 (9)	0.0296 (3)
C5	0.3813 (3)	0.26597 (13)	0.23161 (8)	0.0269 (3)
C6	0.3433 (3)	0.19722 (13)	0.13128 (8)	0.0279 (3)
H6	0.2053	0.1020	0.1061	0.034*
C7	0.5573 (3)	0.43449 (15)	0.37259 (9)	0.0357 (3)
H7	0.6669	0.5214	0.4182	0.043*
C8	0.0238 (3)	0.06785 (15)	0.31003 (9)	0.0337 (3)
H8A	-0.1083	0.0416	0.2447	0.040*
H8B	-0.1104	0.0882	0.3595	0.040*
C9	0.1822 (3)	-0.07536 (15)	0.33350 (11)	0.0430 (3)
H9	0.3288	-0.1040	0.2942	0.052*
C10	0.1274 (4)	-0.16305 (18)	0.40592 (12)	0.0599 (4)
H10A	0.2369	-0.2555	0.4167	0.072*
H10B	-0.0296	-0.1344	0.4488	0.072*
N1	0.5029 (2)	0.21201 (12)	-0.03272 (7)	0.0335 (3)
N2	0.2434 (2)	0.21521 (11)	0.30948 (7)	0.0306 (2)
N3	0.3528 (3)	0.31791 (12)	0.39570 (7)	0.0360 (3)
O1	0.2915 (2)	0.09935 (12)	-0.06728 (7)	0.0468 (3)
O2	0.6971 (3)	0.26879 (12)	-0.08121 (7)	0.0515 (3)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0332 (6)	0.0299 (5)	0.0236 (6)	0.0101 (5)	0.0050 (5)	0.0012 (4)
C2	0.0385 (7)	0.0296 (6)	0.0340 (6)	0.0042 (5)	0.0099 (5)	0.0051 (5)
C3	0.0399 (7)	0.0263 (5)	0.0361 (7)	0.0005 (5)	0.0068 (5)	0.0003 (5)
C4	0.0342 (6)	0.0265 (5)	0.0271 (6)	0.0064 (5)	0.0028 (5)	-0.0015 (4)
C5	0.0279 (5)	0.0280 (5)	0.0255 (6)	0.0073 (4)	0.0036 (4)	0.0031 (4)
C6	0.0290 (6)	0.0276 (5)	0.0257 (6)	0.0046 (4)	0.0018 (4)	-0.0008 (4)
C7	0.0429 (7)	0.0339 (6)	0.0266 (6)	0.0020 (5)	0.0029 (5)	-0.0044 (5)
C8	0.0300 (6)	0.0421 (6)	0.0261 (6)	-0.0028 (5)	0.0041 (5)	0.0023 (5)
C9	0.0370 (7)	0.0356 (7)	0.0521 (8)	-0.0037 (5)	0.0074 (6)	-0.0032 (6)
C10	0.0817 (12)	0.0388 (7)	0.0518 (9)	0.0053 (8)	-0.0079 (8)	0.0024 (7)
N1	0.0409 (6)	0.0351 (5)	0.0263 (5)	0.0117 (4)	0.0070 (4)	0.0021 (4)
N2	0.0343 (5)	0.0336 (5)	0.0223 (5)	0.0018 (4)	0.0048 (4)	0.0005 (4)
N3	0.0426 (6)	0.0387 (6)	0.0240 (5)	0.0034 (5)	0.0036 (4)	-0.0031 (4)
O1	0.0512 (6)	0.0529 (6)	0.0297 (5)	-0.0007 (5)	0.0020 (4)	-0.0088 (4)
O2	0.0659 (7)	0.0549 (6)	0.0364 (5)	0.0031 (5)	0.0255 (5)	0.0017 (4)

Geometric parameters (Å, °)

C1—C6	1.3706 (16)	C7—H7	0.9300
C1—C2	1.4036 (17)	C8—N2	1.4522 (15)
C1—N1	1.4720 (15)	C8—C9	1.4942 (18)
C2—C3	1.3691 (17)	C8—H8A	0.9700
C2—H2	0.9300	C8—H8B	0.9700

C3—C4	1.4016 (16)	C9—C10	1.309 (2)
C3—H3	0.9300	C9—H9	0.9300
C4—C5	1.4087 (15)	C10—H10A	0.9711
C4—C7	1.4174 (17)	C10—H10B	0.9983
C5—N2	1.3631 (14)	N1—O2	1.2203 (13)
C5—C6	1.3985 (16)	N1—O1	1.2256 (14)
C6—H6	0.9300	N2—N3	1.3605 (14)
C7—N3	1.3204 (16)		
C6—C1—C2	124.41 (11)	C4—C7—H7	124.2
C6—C1—N1	117.64 (10)	N2—C8—C9	112.95 (10)
C2—C1—N1	117.95 (10)	N2—C8—H8A	109.0
C3—C2—C1	119.75 (11)	C9—C8—H8A	109.0
C3—C2—H2	120.1	N2—C8—H8B	109.0
C1—C2—H2	120.1	C9—C8—H8B	109.0
C2—C3—C4	118.53 (11)	H8A—C8—H8B	107.8
C2—C3—H3	120.7	C10—C9—C8	124.06 (14)
C4—C3—H3	120.7	C10—C9—H9	118.0
C3—C4—C5	119.81 (11)	C8—C9—H9	118.0
C3—C4—C7	136.30 (11)	C9—C10—H10A	119.9
C5—C4—C7	103.89 (10)	C9—C10—H10B	119.2
N2—C5—C6	130.53 (10)	H10A—C10—H10B	120.9
N2—C5—C4	106.94 (10)	O2—N1—O1	123.28 (10)
C6—C5—C4	122.53 (10)	O2—N1—C1	118.53 (10)
C1—C6—C5	114.96 (10)	O1—N1—C1	118.18 (10)
C1—C6—H6	122.5	N3—N2—C5	111.12 (10)
C5—C6—H6	122.5	N3—N2—C8	120.52 (9)
N3—C7—C4	111.61 (11)	C5—N2—C8	128.23 (9)
N3—C7—H7	124.2	C7—N3—N2	106.44 (10)
C6—C1—C2—C3	0.27 (18)	N2—C8—C9—C10	125.37 (14)
N1—C1—C2—C3	-178.80 (10)	C6—C1—N1—O2	-168.29 (10)
C1—C2—C3—C4	0.77 (17)	C2—C1—N1—O2	10.84 (16)
C2—C3—C4—C5	-1.17 (16)	C6—C1—N1—O1	11.05 (16)
C2—C3—C4—C7	178.55 (13)	C2—C1—N1—O1	-169.83 (10)
C3—C4—C5—N2	-179.75 (10)	C6—C5—N2—N3	178.95 (11)
C7—C4—C5—N2	0.45 (12)	C4—C5—N2—N3	-0.67 (12)
C3—C4—C5—C6	0.59 (16)	C6—C5—N2—C8	3.07 (19)
C7—C4—C5—C6	-179.21 (10)	C4—C5—N2—C8	-176.55 (11)
C2—C1—C6—C5	-0.84 (16)	C9—C8—N2—N3	-88.19 (13)
N1—C1—C6—C5	178.23 (9)	C9—C8—N2—C5	87.34 (14)
N2—C5—C6—C1	-179.17 (10)	C4—C7—N3—N2	-0.31 (14)
C4—C5—C6—C1	0.40 (15)	C5—N2—N3—C7	0.62 (13)
C3—C4—C7—N3	-179.84 (13)	C8—N2—N3—C7	176.85 (10)
C5—C4—C7—N3	-0.09 (13)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
C6—H6···O1 ⁱ	0.93	2.51	3.3973 (17)	160
C8—H8A···O1 ⁱ	0.97	2.53	3.4475 (19)	157
C2—H2···O2 ⁱⁱ	0.93	2.66	3.3911 (17)	136

Symmetry codes: (i) $-x, -y, -z$; (ii) $-x+2, -y+1, -z$.