

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(E)-2,4-Dimethyl-N'-(2-methylbenzylidene)benzohydrazide

Muhammad Taha,^a Nor Hadiani Ismail,^b
Faridahanim Mohd Jaafar,^b Khalid M. Khan^c and Sammer
Yousuf^{c*}

^aAtta-ur-Rahman Institute for Natural Product Discovery, Universiti Teknologi MARA (UiTM), Puncak Alam Campus, 42300 Bandar Puncak Alam, Selangor, Malaysia,

^bFaculty of Applied Science, Universiti Teknologi MARA (UiTM), 40450 Shah Alam, Malaysia, and ^cH.E.J. Research Institute of Chemistry, International Center for Chemical and Biological Sciences, University of Karachi, Karachi 75270, Pakistan
Correspondence e-mail: dr.sammer.yousuf@gmail.com


Received 11 February 2013; accepted 14 February 2013

Key indicators: single-crystal X-ray study; $T = 273$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.030; wR factor = 0.080; data-to-parameter ratio = 13.6.

In the title benzoylhydrazide derivative, $\text{C}_{17}\text{H}_{18}\text{N}_2\text{O}$, the dihedral angle between the benzene rings is $88.45(8)^\circ$ and the azomethine double bond adopts an *E* conformation. In the crystal, molecules are linked by $\text{N}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds, forming a chain along the *b* axis.

Related literature

For the applications and biological activity of Schiff bases, see: Musharraf *et al.* (2012); Khan *et al.* (2012). For the crystal structures of related compounds, see: Taha *et al.* (2012a,b); Naz *et al.* (2012).



Experimental

Crystal data

$\text{C}_{17}\text{H}_{18}\text{N}_2\text{O}$
 $M_r = 266.33$
Orthorhombic, $Pca2_1$
 $a = 26.1151(10)$ Å
 $b = 4.9484(2)$ Å
 $c = 11.3933(4)$ Å

$V = 1472.33(10)$ Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.08$ mm⁻¹
 $T = 273$ K
 $0.56 \times 0.55 \times 0.23$ mm

Data collection

Bruker SMART APEX CCD area-detector diffractometer
Absorption correction: multi-scan (SADABS; Bruker, 2000)
 $T_{\min} = 0.959$, $T_{\max} = 0.983$
8023 measured reflections
2577 independent reflections
2483 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.021$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.080$
 $S = 1.04$
2577 reflections
189 parameters
1 restraint
H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.12$ e Å⁻³
 $\Delta\rho_{\min} = -0.14$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1A}\cdots\text{O1}^i$	0.833 (15)	2.000 (15)	2.8150 (14)	166.1 (14)
$\text{C8}-\text{H8A}\cdots\text{O1}^i$	0.93	2.52	3.2696 (19)	138

Symmetry code: (i) $x, y - 1, z$.

Data collection: SMART (Bruker, 2000); cell refinement: SAINT (Bruker, 2000); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL, PARST (Nardelli, 1995) and PLATON (Spek, 2009).

The authors would like to acknowledge the Research Management Institute of UiTM for financial support under the Dana Kecemerlangan Grant Scheme [grant No. 600-RMI/DANA 5/3 RIF (143/2012)].

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS5246).

References

- Bruker (2000). *SADABS*, *SMART* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Khan, K. M., Taha, M., Naz, F., Siddiqui, S., Rahim, F., Perveen, S. & Choudhary, M. I. (2012). *Med. Chem.* **8**, 705–710.
- Musharraf, S. G., Bibi, A., Shahid, N., Najam-ul-Haq, M., Khan, M., Taha, M., Mughal, U. R. & Khan, K. M. (2012). *Am. J. Anal. Chem.* **3**, 779–789.
- Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
- Naz, H., Taha, M., Rahman, A. A., Ismail, N. H. & Yousuf, S. (2012). *Acta Cryst.* **E68**, o2671.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Taha, M., Naz, H., Rahman, A. A., Ismail, N. H. & Sammer, Y. (2012a). *Acta Cryst.* **E68**, o2778.
- Taha, M., Naz, H., Rahman, A. A., Ismail, N. H. & Sammer, Y. (2012b). *Acta Cryst.* **E68**, o2780.

supporting information

Acta Cryst. (2013). E69, o400 [doi:10.1107/S1600536813004388]

(E)-2,4-Dimethyl-N'-(2-methylbenzylidene)benzohydrazide

Muhammad Taha, Nor Hadiani Ismail, Faridahanim Mohd Jaafar, Khalid M. Khan and Sammer Yousuf

S1. Comment


Benzohydrazides represents an important class of organic compounds with a wide range of biological applications (Musharraf *et al.*, 2012; Khan *et al.*, 2012). The title compound was obtained in continuation of our work to synthesize and study the biological activities of benzohydrazide derivatives. Previously, we have published crystal structures of many benzohydrazides derivatives with different substitution pattern on two phenyl rings (Taha *et al.*, 2012*a,b*; Naz *et al.*, 2012). In the title compound two methyl substituted phenyl rings (C1–C6 and C9–C14) are each planner with dihedral angle of 88.45 (8)° between them. The bond lengths and angles were found to be similar as in structurally related benzohydrazide derivatives (Taha *et al.*, 2012*a,b*; Naz *et al.*, 2012). The crystal structure stabilize by intermolecular N1—H1A···O1ⁱ and C8—H8A···O1ⁱ interactions to form a chain along the *b* axis (symmetry code as in Table 1).

S2. Experimental


The title compound was synthesized by using (0.328 g) 2 mmol of 2,4-dimethylbenzohydrazide and (0.240 g) 2 mmol 2-methylbenzaldehyde as starting material under same conditions and solvents as described in our previous publications (Taha *et al.*, 2012*a,b*; Naz *et al.*, 2012). The compound was recrystallized by dissolving in methanol to obtain colorless needles (0.474 g, 89% yield). All chemicals were purchased by sigma Aldrich Germany.

S3. Refinement

H atoms on methyl and phenyl groups were positioned geometrically with C—H = 0.96 and 0.93 Å, respectively, and constrained to ride on their parent atoms with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C}_{\text{methyl}})$ or $1.2U_{\text{eq}}(\text{C}_{\text{phenyl}})$. A rotating group model was applied to the methyl groups. The H atoms on the nitrogen was located in a difference Fourier map and refined isotropically [N—H = 0.832 (15) Å]. The Hooft *y* parameter was -0.3 (5).

**Figure 1**

The molecular structure of the title compound, with displacement ellipsoids drawn at 30% probability level.

**Figure 2**

A crystal packing diagram of the title compound. Only hydrogen atoms involved in the hydrogen bonds (dashed lines) are shown.

(E)-2,4-Dimethyl-N'-(2-methylbenzylidene)benzohydrazide*Crystal data*C₁₇H₁₈N₂O $M_r = 266.33$ Orthorhombic, *Pca*2₁

Hall symbol: P 2c -2ac

 $a = 26.1151 (10) \text{ \AA}$ $b = 4.9484 (2) \text{ \AA}$ $c = 11.3933 (4) \text{ \AA}$ $V = 1472.33 (10) \text{ \AA}^3$ $Z = 4$ $F(000) = 568$ $D_x = 1.202 \text{ Mg m}^{-3}$ Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 4590 reflections

 $\theta = 2.4\text{--}27.4^\circ$ $\mu = 0.08 \text{ mm}^{-1}$ $T = 273 \text{ K}$

Block, colorless

 $0.56 \times 0.55 \times 0.23 \text{ mm}$ *Data collection*Bruker SMART APEX CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 ω scanAbsorption correction: multi-scan
(*SADABS*; Bruker, 2000) $T_{\min} = 0.959, T_{\max} = 0.983$

8023 measured reflections

2577 independent reflections

2483 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.021$ $\theta_{\max} = 25.5^\circ, \theta_{\min} = 2.4^\circ$ $h = -25 \rightarrow 31$ $k = -5 \rightarrow 5$ $l = -13 \rightarrow 13$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.030$ $wR(F^2) = 0.080$ $S = 1.04$

2577 reflections

189 parameters

1 restraint

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sitesH atoms treated by a mixture of independent
and constrained refinement $w = 1/[\sigma^2(F_o^2) + (0.0459P)^2 + 0.1205P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.12 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -0.14 \text{ e \AA}^{-3}$ Extinction correction: *SHELXTL* (Sheldrick,
2008), $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.016 (2)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.40241 (5)	0.5493 (2)	0.43046 (18)	0.0938 (6)
N1	0.38352 (4)	0.1086 (2)	0.44326 (11)	0.0438 (3)
H1A	0.3925 (5)	-0.050 (3)	0.4301 (13)	0.039 (4)*

N2	0.33498 (4)	0.1558 (2)	0.48752 (10)	0.0448 (3)
C1	0.49419 (6)	0.0399 (3)	0.44104 (14)	0.0500 (3)
H1B	0.4797	-0.0346	0.5082	0.060*
C2	0.54235 (6)	-0.0412 (3)	0.40641 (15)	0.0600 (4)
H2B	0.5601	-0.1679	0.4510	0.072*
C3	0.56461 (6)	0.0633 (4)	0.30653 (15)	0.0605 (4)
C4	0.53740 (6)	0.2556 (4)	0.24413 (14)	0.0588 (4)
H4A	0.5524	0.3297	0.1775	0.071*
C5	0.48869 (6)	0.3437 (3)	0.27612 (13)	0.0488 (4)
C6	0.46699 (5)	0.2308 (3)	0.37737 (12)	0.0426 (3)
C7	0.41514 (5)	0.3145 (3)	0.41905 (14)	0.0480 (3)
C8	0.30625 (5)	-0.0517 (3)	0.49163 (13)	0.0459 (3)
H8A	0.3181	-0.2157	0.4624	0.055*
C9	0.25464 (5)	-0.0345 (3)	0.54203 (13)	0.0458 (3)
C10	0.24424 (6)	0.1476 (3)	0.63203 (14)	0.0553 (4)
H10A	0.2703	0.2579	0.6602	0.066*
C11	0.19599 (7)	0.1674 (4)	0.68005 (15)	0.0656 (5)
H11A	0.1893	0.2922	0.7392	0.079*
C12	0.15778 (7)	0.0008 (4)	0.63967 (17)	0.0665 (5)
H12A	0.1252	0.0108	0.6722	0.080*
C13	0.16772 (6)	-0.1808 (3)	0.55115 (18)	0.0628 (4)
H13A	0.1414	-0.2922	0.5248	0.075*
C14	0.21564 (5)	-0.2029 (3)	0.50006 (14)	0.0500 (3)
C15	0.22448 (7)	-0.3941 (3)	0.39994 (17)	0.0652 (4)
H15A	0.1940	-0.4979	0.3862	0.098*
H15B	0.2522	-0.5136	0.4189	0.098*
H15C	0.2329	-0.2933	0.3306	0.098*
C16	0.61613 (7)	-0.0348 (6)	0.2648 (2)	0.0969 (8)
H16A	0.6383	-0.0606	0.3311	0.145*
H16B	0.6308	0.0969	0.2129	0.145*
H16C	0.6120	-0.2030	0.2239	0.145*
C17	0.46107 (8)	0.5471 (4)	0.20175 (18)	0.0736 (5)
H17A	0.4250	0.5091	0.2028	0.110*
H17B	0.4735	0.5370	0.1226	0.110*
H17C	0.4670	0.7251	0.2323	0.110*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0731 (8)	0.0328 (5)	0.1753 (17)	0.0011 (5)	0.0407 (10)	-0.0067 (8)
N1	0.0394 (6)	0.0335 (5)	0.0587 (7)	0.0032 (4)	0.0055 (5)	-0.0042 (5)
N2	0.0388 (6)	0.0451 (6)	0.0505 (6)	0.0038 (5)	0.0034 (5)	-0.0020 (5)
C1	0.0471 (8)	0.0556 (8)	0.0472 (8)	-0.0015 (6)	0.0033 (7)	0.0003 (7)
C2	0.0467 (8)	0.0745 (10)	0.0586 (10)	0.0108 (7)	-0.0058 (8)	-0.0036 (8)
C3	0.0421 (8)	0.0810 (11)	0.0583 (9)	-0.0034 (7)	0.0019 (8)	-0.0175 (8)
C4	0.0589 (10)	0.0728 (10)	0.0448 (8)	-0.0218 (8)	0.0111 (7)	-0.0100 (7)
C5	0.0555 (9)	0.0431 (7)	0.0477 (8)	-0.0127 (6)	-0.0011 (7)	-0.0043 (6)
C6	0.0429 (7)	0.0367 (6)	0.0482 (8)	-0.0057 (5)	0.0015 (6)	-0.0066 (5)

C7	0.0460 (8)	0.0349 (6)	0.0630 (9)	-0.0006 (5)	0.0038 (7)	-0.0038 (6)
C8	0.0415 (7)	0.0427 (6)	0.0534 (8)	0.0024 (6)	0.0011 (6)	-0.0018 (6)
C9	0.0414 (7)	0.0469 (7)	0.0492 (7)	0.0032 (6)	0.0032 (6)	0.0059 (6)
C10	0.0508 (9)	0.0632 (9)	0.0520 (8)	-0.0002 (7)	0.0027 (7)	-0.0012 (7)
C11	0.0663 (11)	0.0764 (11)	0.0541 (9)	0.0055 (9)	0.0177 (8)	-0.0022 (8)
C12	0.0499 (9)	0.0789 (11)	0.0707 (11)	0.0030 (8)	0.0207 (9)	0.0056 (9)
C13	0.0458 (8)	0.0650 (9)	0.0774 (11)	-0.0079 (7)	0.0060 (9)	0.0036 (9)
C14	0.0454 (8)	0.0459 (7)	0.0586 (8)	-0.0005 (6)	0.0026 (7)	0.0066 (6)
C15	0.0578 (9)	0.0578 (9)	0.0800 (12)	-0.0053 (7)	0.0007 (9)	-0.0111 (9)
C16	0.0501 (11)	0.150 (2)	0.0910 (16)	0.0072 (12)	0.0129 (10)	-0.0378 (15)
C17	0.0953 (15)	0.0592 (10)	0.0664 (11)	-0.0078 (9)	-0.0073 (11)	0.0132 (8)

Geometric parameters (Å, °)

O1—C7	1.2156 (17)	C9—C14	1.400 (2)
N1—C7	1.3401 (17)	C10—C11	1.377 (2)
N1—N2	1.3842 (16)	C10—H10A	0.9300
N1—H1A	0.832 (15)	C11—C12	1.374 (3)
N2—C8	1.2723 (18)	C11—H11A	0.9300
C1—C2	1.378 (2)	C12—C13	1.376 (3)
C1—C6	1.387 (2)	C12—H12A	0.9300
C1—H1B	0.9300	C13—C14	1.384 (2)
C2—C3	1.378 (3)	C13—H13A	0.9300
C2—H2B	0.9300	C14—C15	1.500 (2)
C3—C4	1.384 (3)	C15—H15A	0.9600
C3—C16	1.507 (2)	C15—H15B	0.9600
C4—C5	1.393 (2)	C15—H15C	0.9600
C4—H4A	0.9300	C16—H16A	0.9600
C5—C6	1.401 (2)	C16—H16B	0.9600
C5—C17	1.500 (2)	C16—H16C	0.9600
C6—C7	1.4933 (19)	C17—H17A	0.9600
C8—C9	1.4675 (19)	C17—H17B	0.9600
C8—H8A	0.9300	C17—H17C	0.9600
C9—C10	1.392 (2)		
C7—N1—N2	120.73 (11)	C11—C10—H10A	119.4
C7—N1—H1A	120.4 (10)	C9—C10—H10A	119.4
N2—N1—H1A	118.9 (10)	C12—C11—C10	119.26 (16)
C8—N2—N1	114.66 (11)	C12—C11—H11A	120.4
C2—C1—C6	121.07 (15)	C10—C11—H11A	120.4
C2—C1—H1B	119.5	C11—C12—C13	120.04 (15)
C6—C1—H1B	119.5	C11—C12—H12A	120.0
C1—C2—C3	120.80 (16)	C13—C12—H12A	120.0
C1—C2—H2B	119.6	C12—C13—C14	122.03 (16)
C3—C2—H2B	119.6	C12—C13—H13A	119.0
C2—C3—C4	117.73 (14)	C14—C13—H13A	119.0
C2—C3—C16	121.07 (18)	C13—C14—C9	117.84 (15)
C4—C3—C16	121.17 (18)	C13—C14—C15	120.58 (15)

C3—C4—C5	123.35 (14)	C9—C14—C15	121.54 (14)
C3—C4—H4A	118.3	C14—C15—H15A	109.5
C5—C4—H4A	118.3	C14—C15—H15B	109.5
C4—C5—C6	117.36 (14)	H15A—C15—H15B	109.5
C4—C5—C17	120.09 (15)	C14—C15—H15C	109.5
C6—C5—C17	122.54 (15)	H15A—C15—H15C	109.5
C1—C6—C5	119.67 (13)	H15B—C15—H15C	109.5
C1—C6—C7	119.13 (13)	C3—C16—H16A	109.5
C5—C6—C7	121.19 (13)	C3—C16—H16B	109.5
O1—C7—N1	122.42 (14)	H16A—C16—H16B	109.5
O1—C7—C6	123.16 (13)	C3—C16—H16C	109.5
N1—C7—C6	114.42 (11)	H16A—C16—H16C	109.5
N2—C8—C9	120.62 (12)	H16B—C16—H16C	109.5
N2—C8—H8A	119.7	C5—C17—H17A	109.5
C9—C8—H8A	119.7	C5—C17—H17B	109.5
C10—C9—C14	119.67 (14)	H17A—C17—H17B	109.5
C10—C9—C8	120.32 (13)	C5—C17—H17C	109.5
C14—C9—C8	120.01 (13)	H17A—C17—H17C	109.5
C11—C10—C9	121.14 (15)	H17B—C17—H17C	109.5
C7—N1—N2—C8	170.77 (14)	C5—C6—C7—O1	-49.4 (2)
C6—C1—C2—C3	-0.8 (2)	C1—C6—C7—N1	-49.99 (18)
C1—C2—C3—C4	1.3 (2)	C5—C6—C7—N1	130.66 (14)
C1—C2—C3—C16	-176.88 (17)	N1—N2—C8—C9	176.93 (12)
C2—C3—C4—C5	-1.2 (2)	N2—C8—C9—C10	-30.9 (2)
C16—C3—C4—C5	177.01 (16)	N2—C8—C9—C14	149.45 (15)
C3—C4—C5—C6	0.5 (2)	C14—C9—C10—C11	-0.6 (2)
C3—C4—C5—C17	-178.19 (16)	C8—C9—C10—C11	179.71 (15)
C2—C1—C6—C5	0.0 (2)	C9—C10—C11—C12	1.2 (3)
C2—C1—C6—C7	-179.36 (14)	C10—C11—C12—C13	-0.9 (3)
C4—C5—C6—C1	0.16 (19)	C11—C12—C13—C14	0.0 (3)
C17—C5—C6—C1	178.77 (14)	C12—C13—C14—C9	0.6 (2)
C4—C5—C6—C7	179.50 (12)	C12—C13—C14—C15	-177.14 (17)
C17—C5—C6—C7	-1.9 (2)	C10—C9—C14—C13	-0.2 (2)
N2—N1—C7—O1	-2.7 (3)	C8—C9—C14—C13	179.43 (14)
N2—N1—C7—C6	177.21 (12)	C10—C9—C14—C15	177.45 (15)
C1—C6—C7—O1	129.9 (2)	C8—C9—C14—C15	-2.9 (2)

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1A \cdots O1 ⁱ	0.833 (15)	2.000 (15)	2.8150 (14)	166.1 (14)
C8—H8A \cdots O1 ⁱ	0.93	2.52	3.2696 (19)	138

Symmetry code: (i) *x*, *y*-1, *z*.