

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N-Cyclohexyl-3-methylbenzamidine

Rui-Qin Liu, Sheng-Di Bai* and Tao Wang

Institute of Applied Chemistry, Shanxi University, Taiyuan 030006, People's Republic of China

Correspondence e-mail: sdbai@sxu.edu.cn


Received 18 February 2013; accepted 5 March 2013

Key indicators: single-crystal X-ray study; $T = 200$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.040; wR factor = 0.098; data-to-parameter ratio = 14.6.

The title amidine compound, $\text{C}_{14}\text{H}_{20}\text{N}_2$, prepared by a one-pot reaction, is asymmetric as only one N atom has an alkyl substituent. The terminal cyclohexyl group connected to the amino N atom is located on the other side of the N—C—N skeleton to the 4-methylbenzene ring and has a chair conformation. The dihedral angle between the phenyl ring and the NCN plane is 47.87 (12)°. In the crystal, molecules are linked *via* N—H...N hydrogen bonds, forming chains propagating along the a -axis direction.

Related literature

For reviews of related metal amidinates and their applications in olefin polymerization, see: Edelmann (1994); Barker & Kilner (1994); Collins (2011); Bai *et al.* (2010); Yang *et al.* (2013). For a review of neutral amidines, see: Coles (2006). For a related synthetic method for amidines, see: Wang *et al.* (2008). For related silyl-linked bis(amidinate) ligands, see: Bai *et al.* (2006).



Experimental

Crystal data

 $\text{C}_{14}\text{H}_{20}\text{N}_2$ $M_r = 216.32$ Orthorhombic, $P2_12_12_1$ $a = 9.064$ (2) Å $b = 11.417$ (3) Å $c = 12.311$ (3) Å $V = 1274.0$ (5) Å³ $Z = 4$ Mo $K\alpha$ radiation $\mu = 0.07$ mm⁻¹ $T = 200$ K $0.30 \times 0.25 \times 0.20$ mm

Data collection

Bruker SMART CCD diffractometer
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.980$, $T_{\max} = 0.987$ 7147 measured reflections
2244 independent reflections
1758 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.042$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.098$
 $S = 1.02$
2244 reflections
154 parametersH atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.15$ e Å⁻³
 $\Delta\rho_{\min} = -0.12$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1A}\cdots\text{N2}^i$	0.90 (2)	2.08 (2)	2.975 (2)	168.0 (18)

Symmetry code: (i) $x + \frac{1}{2}, -y + \frac{1}{2}, -z$.

Data collection: SMART (Bruker, 2000); cell refinement: SAINT (Bruker, 2000); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXL97.

This work was supported by grants from the Natural Science Foundation of China (20702029) and the Natural Science Foundation of Shanxi Province (2008011024).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RK2395).

References

- Bai, S. D., Guo, J. P. & Liu, D. S. (2006). *Dalton Trans.* pp. 2244–2250.
Bai, S. D., Tong, H. B., Guo, J. P., Zhou, M. S., Liu, D. S. & Yuan, S. F. (2010). *Polyhedron*, **29**, 262–269.
Barker, J. & Kilner, M. (1994). *Coord. Chem. Rev.* **133**, 219–300.
Bruker (2000). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
Coles, M. P. (2006). *Dalton Trans.* pp. 985–1001.
Collins, S. (2011). *Coord. Chem. Rev.* **255**, 118–138.
Edelmann, F. T. (1994). *Coord. Chem. Rev.* **137**, 403–481.
Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Wang, J. F., Xu, F., Cai, T. & Shen, Q. (2008). *Org. Lett.* **10**, 445–448.
Yang, Q. K., Zhou, M. S., Tong, H. B., Guo, D. L. & Liu, D. S. (2013). *Inorg. Chem. Commun.* **29**, 1–3.

supporting information

Acta Cryst. (2013). E69, o520 [doi:10.1107/S1600536813006272]

N-Cyclohexyl-3-methylbenzamidinium

Rui-Qin Liu, Sheng-Di Bai and Tao Wang

S1. Comment

The exploration of ancillary ligand systems supporting catalytically active metal centers is a long-standing demand in the coordination chemistry. Amidinates represent an important class in the array comparable to the cyclopentadienyl system (Edelmann, 1994; Barker & Kilner, 1994; Collins, 2011). They are four-electron, monoanionic and *N*-donor bidentate chelates, demonstrating a great diversity by variation of substituents on the conjugated N—C—N backbone. Their steric and electronic properties are easily tunable to meet their requirements of different metal centers. In the course of extending amidinate chemistry, we explored a synthetic pathway to the silyl-linked bis(amidinate) ligands, $[\text{SiMe}_2\{\text{NC}(\text{Ph})\text{N}(\text{R})\}_2]^{2-}$ (Bai *et al.*, 2006). They were applied to synthesize the Group 4 complexes, which were good catalysts for ethylene polymerization (Bai *et al.*, 2010; Yang *et al.*, 2013). Amidines are convenient precursors for both monoanionic amidinate ligands and dianionic *ansa*-bis(amidinate) ligands (Coles, 2006). Some amidines could be prepared by Yb complex catalyzed addition reactions of aromatic amines and nitriles (Wang *et al.*, 2008). Here, the synthesis and crystal structure of a new amidine will be described.

The title compound **I** was prepared by a one-pot reaction of cyclohexylamine, *LiBu*, *m*-tolunitrile and H₂O. The intermediate process involved an addition reaction of lithium amide and nitrile to yield lithium monoamidinate. The suitable for X-ray investigation single-crystal of the title compound was obtained by recrystallization in CH₂Cl₂. Its molecular structure is shown in Fig. 1. Two N atoms connect the central C atom in different lengths of 1.357 (2) Å and 1.284 (2) Å, composing the characteristic N—C—N skeleton for amidine species. The terminal cyclohexyl with chair-like configuration connects the amino N atom. The phenyl group is attached to the central C atom. The angle between the phenyl plane and the [NCN] plane is 47.87 (12)°. Cyclohexyl and phenyl are in opposite directions. Fig. 2 displays the packing view of compound **I**. Molecules of **I** can form the one-dimensional chain extending along *a*-axis through the intermolecular hydrogen bond. The imino N atom is the acceptor for hydrogen atom. In the chain, every adjacent two molecules have *C*₂ rotation symmetrical relationship with each other and the couple serves as the repeatable unit.

S2. Experimental

A solution of *LiBu* (2.2 M, 2.7 ml, 6.0 mmol) in hexane was slowly added into a stirred solution of cyclohexylamine (0.69 ml, 6.0 mmol) in *Et*₂O (*ca* 30 ml) by syringe at 273 K. The reaction mixture was warmed to room temperature and kept stirring for 3 h. Then *m*-tolunitrile (0.71 ml, 6.0 mmol) was added by syringe at 273 K. The reaction mixture was warmed to room temperature and kept stirring for 4 h. H₂O (0.11 ml, 6.0 mmol) was added by syringe at 273 K. After stirred at room temperature for 4 h, the mixture was filtered and the filtrate was dried in vacuum to remove all volatiles. The residue was recrystallized with CH₂Cl₂ and gave colourless crystals of the title compound (yield 0.96 g, 74%). ¹H NMR (300 MHz, CDCl₃): δ = 7.38–7.25 (m, 4H; phenyl), 3.81 (br, 2H; NH), 2.43 (s, 3H; *m*-MePh), 2.15–1.21 (m, 11H; Cy). ¹³C NMR (75 MHz, CDCl₃): δ = 139.3–123.6 (*Ph*), 50.4 (*m*-MePh), 33.8, 26.5, 25.6, 22.0 (*Cy*). Anal. Calc. for C₁₄H₂₀N₂ (216.32): C, 77.73; H, 9.32; N, 12.95%. Found: C, 77.46; H, 9.22; N, 13.04%.

S3. Refinement

The methyl H atoms were constrained to an ideal geometry, with C—H distances of 0.98 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$, but each group was allowed to rotate freely about its C—C bond. The methylene H atoms were constrained with C—H distances of 0.99 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The methine H atom was constrained with C—H distance of 1.00 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The phenyl H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms, with C—H distances in the range 0.95 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

The Flack parameter was omitted in CIF because no any atoms heavy Si.




Figure 1

The molecular structure, showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level. H atoms are presented as a small spheres of arbitrary radius.




Figure 2

The view of one-dimensional chain in crystal structure of **I**. Symmetry codes: (i) $x+1/2, -y+1/2, -z$.

N-Cyclohexyl-3-methylbenzamidinium*Crystal data*C₁₄H₂₀N₂ $M_r = 216.32$ Orthorhombic, $P2_12_12_1$

Hall symbol: P 2ac 2ab

 $a = 9.064 (2) \text{ \AA}$ $b = 11.417 (3) \text{ \AA}$ $c = 12.311 (3) \text{ \AA}$ $V = 1274.0 (5) \text{ \AA}^3$ $Z = 4$ $F(000) = 472$ $D_x = 1.128 \text{ Mg m}^{-3}$ Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 1347 reflections

 $\theta = 2.8\text{--}20.0^\circ$ $\mu = 0.07 \text{ mm}^{-1}$ $T = 200 \text{ K}$

Block, colourless

 $0.30 \times 0.25 \times 0.20 \text{ mm}$ *Data collection*

Bruker SMART CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 φ and ω scan

Absorption correction: multi-scan

(SADABS; Sheldrick, 1996)

 $T_{\min} = 0.980$, $T_{\max} = 0.987$

7147 measured reflections

2244 independent reflections

1758 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.042$ $\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 2.4^\circ$ $h = -10 \rightarrow 7$ $k = -13 \rightarrow 12$ $l = -14 \rightarrow 14$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.040$ $wR(F^2) = 0.098$ $S = 1.02$

2244 reflections

154 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier

map

Hydrogen site location: inferred from

neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

 $w = 1/[\sigma^2(F_o^2) + (0.0452P)^2 + 0.0788P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.15 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -0.12 \text{ e \AA}^{-3}$ Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.020 (3)

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.12298 (18)	0.22773 (14)	0.08238 (13)	0.0434 (4)
N2	-0.08387 (18)	0.17325 (15)	-0.02421 (15)	0.0475 (5)
C1	0.1003 (2)	0.14588 (16)	0.17183 (15)	0.0391 (5)

H1B	-0.0079	0.1422	0.1872	0.047*
C2	0.1528 (2)	0.02235 (17)	0.14538 (16)	0.0502 (6)
H2C	0.0991	-0.0073	0.0810	0.060*
H2D	0.2593	0.0240	0.1277	0.060*
C3	0.1267 (3)	-0.05956 (17)	0.24111 (18)	0.0611 (7)
H3A	0.0195	-0.0656	0.2553	0.073*
H3B	0.1638	-0.1387	0.2228	0.073*
C4	0.2041 (3)	-0.0156 (2)	0.34235 (18)	0.0612 (7)
H4A	0.3122	-0.0170	0.3308	0.073*
H4B	0.1809	-0.0679	0.4042	0.073*
C5	0.1552 (3)	0.10824 (19)	0.36872 (17)	0.0612 (6)
H5A	0.2124	0.1376	0.4316	0.073*
H5B	0.0496	0.1075	0.3894	0.073*
C6	0.1769 (2)	0.19043 (17)	0.27290 (16)	0.0485 (5)
H6A	0.1375	0.2687	0.2916	0.058*
H6B	0.2837	0.1989	0.2581	0.058*
C7	0.0332 (2)	0.23343 (16)	-0.00570 (16)	0.0382 (5)
C8	0.0761 (2)	0.32275 (16)	-0.08846 (16)	0.0387 (5)
C9	0.1108 (2)	0.43699 (16)	-0.05829 (17)	0.0460 (5)
H9A	0.1104	0.4587	0.0162	0.055*
C10	0.1457 (3)	0.51857 (19)	-0.13683 (18)	0.0559 (6)
H10A	0.1683	0.5967	-0.1162	0.067*
C11	0.1480 (3)	0.48753 (19)	-0.24477 (19)	0.0578 (6)
H11A	0.1724	0.5446	-0.2980	0.069*
C12	0.1153 (2)	0.37453 (19)	-0.27700 (17)	0.0520 (6)
C13	0.0782 (2)	0.29362 (17)	-0.19718 (16)	0.0446 (5)
H13A	0.0536	0.2160	-0.2181	0.054*
C14	0.1184 (3)	0.3393 (2)	-0.3945 (2)	0.0864 (9)
H14A	0.0928	0.2562	-0.4010	0.130*
H14B	0.2174	0.3523	-0.4240	0.130*
H14C	0.0469	0.3864	-0.4352	0.130*
H2A	-0.098 (2)	0.1201 (17)	0.0303 (17)	0.054 (6)*
H1A	0.213 (2)	0.2603 (17)	0.0752 (16)	0.052 (6)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0400 (10)	0.0501 (10)	0.0401 (9)	-0.0084 (9)	-0.0028 (8)	0.0139 (8)
N2	0.0419 (10)	0.0512 (10)	0.0495 (11)	-0.0071 (9)	-0.0050 (8)	0.0109 (9)
C1	0.0377 (10)	0.0413 (11)	0.0384 (11)	-0.0025 (9)	0.0019 (9)	0.0065 (8)
C2	0.0573 (14)	0.0481 (12)	0.0451 (13)	0.0004 (11)	-0.0039 (11)	-0.0022 (10)
C3	0.0705 (17)	0.0405 (11)	0.0721 (16)	-0.0032 (12)	-0.0081 (14)	0.0079 (12)
C4	0.0729 (16)	0.0571 (14)	0.0536 (15)	0.0025 (14)	-0.0025 (13)	0.0189 (11)
C5	0.0785 (16)	0.0640 (15)	0.0410 (13)	0.0052 (13)	-0.0004 (12)	0.0068 (11)
C6	0.0565 (13)	0.0438 (11)	0.0450 (12)	0.0025 (11)	-0.0040 (10)	-0.0009 (10)
C7	0.0364 (10)	0.0379 (11)	0.0402 (12)	0.0025 (9)	0.0009 (9)	0.0016 (9)
C8	0.0342 (10)	0.0392 (11)	0.0427 (12)	0.0062 (9)	-0.0006 (9)	0.0042 (9)
C9	0.0502 (13)	0.0430 (11)	0.0448 (12)	0.0024 (10)	-0.0028 (10)	0.0033 (10)

C10	0.0701 (16)	0.0387 (11)	0.0590 (15)	-0.0026 (11)	-0.0040 (12)	0.0082 (11)
C11	0.0638 (16)	0.0524 (13)	0.0570 (15)	-0.0022 (12)	0.0016 (12)	0.0212 (12)
C12	0.0552 (14)	0.0590 (13)	0.0418 (13)	0.0052 (12)	0.0000 (11)	0.0089 (11)
C13	0.0460 (12)	0.0448 (11)	0.0431 (12)	0.0012 (10)	-0.0011 (9)	0.0006 (10)
C14	0.123 (2)	0.0921 (19)	0.0440 (14)	-0.0059 (19)	0.0068 (16)	0.0102 (14)

Geometric parameters (Å, °)

N1—C7	1.357 (2)	C5—H5B	0.9900
N1—C1	1.459 (2)	C6—H6A	0.9900
N1—H1A	0.90 (2)	C6—H6B	0.9900
N2—C7	1.284 (2)	C7—C8	1.493 (3)
N2—H2A	0.91 (2)	C8—C13	1.379 (3)
C1—C6	1.513 (3)	C8—C9	1.392 (3)
C1—C2	1.524 (3)	C9—C10	1.379 (3)
C1—H1B	1.0000	C9—H9A	0.9500
C2—C3	1.523 (3)	C10—C11	1.375 (3)
C2—H2C	0.9900	C10—H10A	0.9500
C2—H2D	0.9900	C11—C12	1.382 (3)
C3—C4	1.516 (3)	C11—H11A	0.9500
C3—H3A	0.9900	C12—C13	1.390 (3)
C3—H3B	0.9900	C12—C14	1.502 (3)
C4—C5	1.517 (3)	C13—H13A	0.9500
C4—H4A	0.9900	C14—H14A	0.9800
C4—H4B	0.9900	C14—H14B	0.9800
C5—C6	1.520 (3)	C14—H14C	0.9800
C5—H5A	0.9900		
C7—N1—C1	123.32 (16)	C1—C6—C5	111.80 (17)
C7—N1—H1A	116.5 (13)	C1—C6—H6A	109.3
C1—N1—H1A	117.7 (13)	C5—C6—H6A	109.3
C7—N2—H2A	110.1 (13)	C1—C6—H6B	109.3
N1—C1—C6	109.93 (15)	C5—C6—H6B	109.3
N1—C1—C2	112.82 (15)	H6A—C6—H6B	107.9
C6—C1—C2	110.11 (16)	N2—C7—N1	127.70 (18)
N1—C1—H1B	107.9	N2—C7—C8	117.35 (18)
C6—C1—H1B	107.9	N1—C7—C8	114.94 (16)
C2—C1—H1B	107.9	C13—C8—C9	118.79 (18)
C3—C2—C1	110.77 (16)	C13—C8—C7	120.08 (17)
C3—C2—H2C	109.5	C9—C8—C7	121.12 (18)
C1—C2—H2C	109.5	C10—C9—C8	119.8 (2)
C3—C2—H2D	109.5	C10—C9—H9A	120.1
C1—C2—H2D	109.5	C8—C9—H9A	120.1
H2C—C2—H2D	108.1	C11—C10—C9	120.4 (2)
C4—C3—C2	111.18 (18)	C11—C10—H10A	119.8
C4—C3—H3A	109.4	C9—C10—H10A	119.8
C2—C3—H3A	109.4	C10—C11—C12	121.0 (2)
C4—C3—H3B	109.4	C10—C11—H11A	119.5

C2—C3—H3B	109.4	C12—C11—H11A	119.5
H3A—C3—H3B	108.0	C11—C12—C13	118.0 (2)
C3—C4—C5	110.44 (19)	C11—C12—C14	121.5 (2)
C3—C4—H4A	109.6	C13—C12—C14	120.5 (2)
C5—C4—H4A	109.6	C8—C13—C12	121.95 (19)
C3—C4—H4B	109.6	C8—C13—H13A	119.0
C5—C4—H4B	109.6	C12—C13—H13A	119.0
H4A—C4—H4B	108.1	C12—C14—H14A	109.5
C4—C5—C6	111.80 (18)	C12—C14—H14B	109.5
C4—C5—H5A	109.3	H14A—C14—H14B	109.5
C6—C5—H5A	109.3	C12—C14—H14C	109.5
C4—C5—H5B	109.3	H14A—C14—H14C	109.5
C6—C5—H5B	109.3	H14B—C14—H14C	109.5
H5A—C5—H5B	107.9		

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
N1—H1A...N2 ⁱ	0.90 (2)	2.08 (2)	2.975 (2)	168.0 (18)

Symmetry code: (i) $x+1/2, -y+1/2, -z$.