

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Ethyl 3-bromo-4-cyano-5-[(2-ethoxy-2-oxoethyl)sulfanyl]thiophene-2-carboxylate

Xiuping Li,* Xiaochuan Jia and Jing Li

Tianjin Entry–Exit Inspection and Quarantine Bureau, Youyi road No. 33, Tianjin, People's Republic of China

Correspondence e-mail: xpli0705@163.com


Received 14 March 2013; accepted 18 April 2013

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}–\text{C}) = 0.006$ Å; R factor = 0.052; wR factor = 0.157; data-to-parameter ratio = 19.5.

The title compound, $\text{C}_{12}\text{H}_{12}\text{BrNO}_4\text{S}_2$, was obtained by the Sandmeyer reaction from ethyl 3-amino-4-cyano-5-[(2-ethoxy-2-oxoethyl)sulfanyl]thiophene-2-carboxylate. The dihedral angle between the thiophene ring and linked CO_2 ester group is $2.0(5)^\circ$.

Related literature

For background literature on the use of 3-amino-4-cyano-5-ethoxycarbonylmethylsulfanyl-thiophene-2-carboxylic acid ethyl ester as an important intermediate compound for the synthesis of thienopyrimidine derivatives, which are thought to be potential biologically active compounds or pharmaceuticals, see: Liu *et al.* (2008). For a related compound, see: Padmavathi *et al.* (2011).



Experimental

Crystal data

$\text{C}_{12}\text{H}_{12}\text{BrNO}_4\text{S}_2$
 $M_r = 378.26$
 Monoclinic, $P2_1/c$
 $a = 8.5896(17)$ Å
 $b = 10.837(2)$ Å
 $c = 16.584(3)$ Å
 $\beta = 99.44(3)^\circ$

$V = 1522.8(5)$ Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 2.98$ mm⁻¹
 $T = 293$ K
 $0.20 \times 0.18 \times 0.12$ mm

Data collection

Rigaku Saturn diffractometer
 Absorption correction: multi-scan
 (*CrystalClear*; Rigaku, 2005)
 $T_{\min} = 0.587$, $T_{\max} = 0.716$

15211 measured reflections
 3596 independent reflections
 2224 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.067$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.052$
 $wR(F^2) = 0.157$
 $S = 1.00$
 3596 reflections

184 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.43$ e Å⁻³
 $\Delta\rho_{\min} = -0.71$ e Å⁻³

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors appreciate the help of Dr Haibin Song in Nankai University for the crystal X-ray measurement.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZJ2102).

References

- Liu, M. G., Hu, Y. G. & Ding, M. W. (2008). *Tetrahedron*, **64**, 9052–9059.
 Padmavathi, V., Reddy, G. D., Reddy, S. N. & Mahesh, K. (2011). *Eur. J. Med. Chem.* **46**, 1367–1373.
 Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2013). E69, o848 [doi:10.1107/S1600536813010611]

Ethyl 3-bromo-4-cyano-5-[(2-ethoxy-2-oxoethyl)sulfanyl]thiophene-2-carboxylate

Xiuping Li, Xiaochuan Jia and Jing Li

Comment

3-Amino-4-cyano-5-ethoxycarbonylmethylsulfanyl-thiophene-2-carboxylic acid ethyl ester is an important intermediate compound for synthesis of thienothienopyrimidines derivatives, which are thought to be potential biological active compounds or pharmaceuticals (Liu, *et al.*, 2008). We obtained the title compound by the Sandmeyer reaction from compound 3-Amino-4-cyano-5-ethoxycarbonylmethylsulfanyl-thiophene-2-carboxylic acid ethyl ester. The crystal for X-ray crystal structure analysis was obtained by recrystallizing the title compound in petroleum ether. In the crystal, the thiophene ring together with its four adjoint groups, *i.e.* CN, Br, S—CH₂ and COO⁻, was located at one perfect plane, which is consistent with the crystal of 3-Amino-4-cyano-5-ethoxycarbonylmethylsulfanyl-thiophene-2-carboxylic acid ethyl ester reported in the literature (Padmavathi *et al.*, 2011). The title compound crystal demonstrated a crystal system of monoclinic and a space group of P2(1)/c. There existed hydrogen bond with length of 2.554 Å between one of —SCH₂ H atom and the O atom of carbonyl adjoined with thiophene ring of neighbor molecule.

Experimental

To a solution of 3-Amino-4-cyano-5-ethoxycarbonylmethylsulfanyl -thiophene-2-carboxylic acid ethyl ester (1.57 g, 5 mmol) in 70% H₂SO₄ (13 mL) was added NaNO₂ (0.4 g, 5.7 mmol) in 5 minutes under ice water temperature. After addition, the solution was stirred for 30 min at room temperature. Then, the reaction mixture was transferred to HBr solution containing CuBr (1 g, 7 mmol). After standing overnight, water (100 mL) was added. The precipitate was collected by filtration and recrystallized from petroleum ether to afford the title compound as colourless crystals, yield 50%.

Computing details

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear* (Rigaku, 2005); data reduction: *CrystalClear* (Rigaku, 2005); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).




Figure 1

The structure of the title compound showing 30% probability ellipsoids.




Figure 2

Packing structure of the title compound.

Ethyl 3-bromo-4-cyano-5-[(2-ethoxy-2-oxoethyl)sulfanyl]thiophene-2-carboxylate

Crystal data

C₁₂H₁₂BrNO₄S₂
M_r = 378.26
 Monoclinic, *P*2₁/*c*
 Hall symbol: -*P* 2ybc
a = 8.5896 (17) Å
b = 10.837 (2) Å
c = 16.584 (3) Å
 β = 99.44 (3)°
V = 1522.8 (5) Å³
Z = 4

F(000) = 760
D_x = 1.650 Mg m⁻³
 Mo *K*α radiation, λ = 0.71073 Å
 Cell parameters from 3404 reflections
 θ = 3.1–27.9°
 μ = 2.98 mm⁻¹
T = 293 K
 Block, colorless
 0.20 × 0.18 × 0.12 mm

Data collection

Rigaku Saturn
 diffractometer
 Radiation source: rotating anode
 Confocal monochromator
 ω scans
 Absorption correction: multi-scan
 (*CrystalClear*; Rigaku, 2005)
T_{min} = 0.587, *T_{max}* = 0.716

15211 measured reflections
 3596 independent reflections
 2224 reflections with *I* > 2σ(*I*)
R_{int} = 0.067
 θ_{\max} = 27.9°, θ_{\min} = 3.1°
h = -11→11
k = -14→14
l = -21→21

Refinement

Refinement on *F*²
 Least-squares matrix: full
R[*F*² > 2σ(*F*²)] = 0.052
wR(*F*²) = 0.157
S = 1.00
 3596 reflections
 184 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map

Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0817P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.003$
 $\Delta\rho_{\max} = 0.43 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.71 \text{ e \AA}^{-3}$
 Extinction correction: *SHELXL97* (Sheldrick,
 2008), $F_c^* = kF_c[1 + 0.001xF_c^2\lambda^3/\sin(2\theta)]^{-1/4}$
 Extinction coefficient: 0.018 (2)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of *F*² against ALL reflections. The weighted *R*-factor *wR* and goodness of fit *S* are based on *F*², conventional *R*-factors *R* are based on *F*, with *F* set to zero for negative *F*². The threshold expression of *F*² > σ(*F*²) is used only for calculating *R*-factors(gt) etc. and is not relevant to the choice of reflections for refinement. *R*-factors based on *F*² are statistically about twice as large as those based on *F*, and *R*-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	<i>U_{iso}</i> */ <i>U_{eq}</i>
Br1	1.34684 (5)	-0.05401 (5)	0.13435 (3)	0.0723 (3)
S1	0.95059 (11)	0.19092 (9)	0.04312 (6)	0.0421 (3)
S2	0.75972 (12)	0.13383 (10)	0.17935 (7)	0.0505 (3)

O1	1.3519 (4)	0.1042 (3)	-0.0246 (2)	0.0703 (9)
O2	1.1477 (3)	0.2261 (3)	-0.07630 (18)	0.0577 (8)
O3	0.4240 (3)	0.3685 (3)	0.10494 (18)	0.0559 (8)
O4	0.4796 (4)	0.2478 (3)	0.2147 (2)	0.0744 (10)
N1	1.0569 (5)	-0.0869 (4)	0.2904 (3)	0.0784 (13)
C1	1.3411 (7)	0.3617 (5)	-0.1215 (3)	0.0825 (16)
H1A	1.2890	0.4277	-0.0980	0.124*
H1B	1.3794	0.3912	-0.1692	0.124*
H1C	1.4282	0.3325	-0.0823	0.124*
C2	1.2285 (6)	0.2598 (5)	-0.1446 (3)	0.0667 (13)
H2A	1.1510	0.2847	-0.1911	0.080*
H2B	1.2845	0.1885	-0.1607	0.080*
C3	1.2245 (5)	0.1481 (4)	-0.0215 (3)	0.0468 (10)
C4	1.1325 (4)	0.1209 (3)	0.0445 (2)	0.0420 (9)
C5	1.1667 (5)	0.0433 (3)	0.1088 (3)	0.0440 (9)
C6	1.0474 (4)	0.0400 (3)	0.1587 (2)	0.0427 (9)
C7	0.9215 (4)	0.1161 (3)	0.1300 (2)	0.0390 (8)
C8	1.0542 (5)	-0.0323 (4)	0.2309 (3)	0.0551 (11)
C9	0.6413 (4)	0.2381 (4)	0.1105 (2)	0.0473 (10)
H9A	0.7050	0.3072	0.0979	0.057*
H9B	0.6003	0.1960	0.0599	0.057*
C10	0.5069 (4)	0.2838 (4)	0.1508 (3)	0.0464 (10)
C11	0.2964 (6)	0.4242 (5)	0.1395 (3)	0.0632 (13)
H11A	0.3373	0.4630	0.1914	0.076*
H11B	0.2205	0.3617	0.1488	0.076*
C12	0.2186 (6)	0.5190 (5)	0.0800 (4)	0.0903 (18)
H12A	0.2957	0.5782	0.0691	0.135*
H12B	0.1371	0.5603	0.1029	0.135*
H12C	0.1733	0.4791	0.0299	0.135*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0499 (4)	0.0728 (4)	0.0911 (5)	0.0186 (2)	0.0023 (3)	0.0000 (3)
S1	0.0381 (5)	0.0419 (5)	0.0472 (6)	0.0045 (4)	0.0093 (4)	0.0038 (4)
S2	0.0452 (6)	0.0526 (6)	0.0570 (7)	0.0040 (5)	0.0186 (5)	0.0114 (5)
O1	0.0539 (19)	0.088 (2)	0.075 (2)	0.0161 (18)	0.0267 (16)	0.0067 (19)
O2	0.0532 (18)	0.0659 (19)	0.0574 (18)	0.0027 (15)	0.0195 (14)	0.0111 (16)
O3	0.0464 (16)	0.0658 (19)	0.0585 (19)	0.0130 (15)	0.0177 (14)	-0.0037 (15)
O4	0.074 (2)	0.088 (2)	0.070 (2)	0.0212 (18)	0.0363 (18)	0.0164 (19)
N1	0.073 (3)	0.077 (3)	0.087 (3)	0.012 (2)	0.020 (2)	0.035 (3)
C1	0.100 (4)	0.089 (4)	0.064 (3)	-0.022 (3)	0.030 (3)	0.002 (3)
C2	0.076 (3)	0.072 (3)	0.056 (3)	-0.004 (3)	0.022 (2)	0.008 (2)
C3	0.040 (2)	0.049 (2)	0.052 (2)	-0.0011 (19)	0.0102 (18)	-0.005 (2)
C4	0.034 (2)	0.039 (2)	0.054 (2)	0.0016 (16)	0.0101 (17)	-0.0058 (18)
C5	0.036 (2)	0.041 (2)	0.054 (2)	-0.0013 (16)	0.0032 (18)	-0.0033 (18)
C6	0.041 (2)	0.039 (2)	0.047 (2)	-0.0030 (17)	0.0034 (18)	0.0045 (17)
C7	0.036 (2)	0.0346 (19)	0.046 (2)	-0.0006 (16)	0.0064 (16)	-0.0012 (16)
C8	0.045 (2)	0.051 (2)	0.070 (3)	0.006 (2)	0.011 (2)	0.011 (2)
C9	0.036 (2)	0.059 (3)	0.048 (2)	0.0034 (19)	0.0114 (18)	0.003 (2)

C10	0.039 (2)	0.053 (2)	0.049 (2)	0.0003 (19)	0.0099 (18)	-0.003 (2)
C11	0.050 (3)	0.070 (3)	0.074 (3)	0.018 (2)	0.023 (2)	-0.008 (3)
C12	0.068 (3)	0.097 (4)	0.108 (5)	0.036 (3)	0.021 (3)	0.015 (4)

Geometric parameters (Å, °)

Br1—C5	1.864 (4)	C2—H2A	0.9700
S1—C7	1.707 (4)	C2—H2B	0.9700
S1—C4	1.734 (4)	C3—C4	1.480 (6)
S2—C7	1.735 (4)	C4—C5	1.352 (5)
S2—C9	1.798 (4)	C5—C6	1.418 (6)
O1—C3	1.202 (5)	C6—C7	1.381 (5)
O2—C3	1.335 (5)	C6—C8	1.425 (6)
O2—C2	1.468 (5)	C9—C10	1.509 (5)
O3—C10	1.323 (5)	C9—H9A	0.9700
O3—C11	1.449 (5)	C9—H9B	0.9700
O4—C10	1.188 (5)	C11—C12	1.504 (7)
N1—C8	1.147 (6)	C11—H11A	0.9700
C1—C2	1.477 (7)	C11—H11B	0.9700
C1—H1A	0.9600	C12—H12A	0.9600
C1—H1B	0.9600	C12—H12B	0.9600
C1—H1C	0.9600	C12—H12C	0.9600
C7—S1—C4	92.15 (19)	C5—C6—C8	124.7 (4)
C7—S2—C9	100.55 (18)	C6—C7—S1	111.1 (3)
C3—O2—C2	116.1 (3)	C6—C7—S2	123.1 (3)
C10—O3—C11	115.6 (3)	S1—C7—S2	125.8 (2)
C2—C1—H1A	109.5	N1—C8—C6	177.4 (5)
C2—C1—H1B	109.5	C10—C9—S2	108.6 (3)
H1A—C1—H1B	109.5	C10—C9—H9A	110.0
C2—C1—H1C	109.5	S2—C9—H9A	110.0
H1A—C1—H1C	109.5	C10—C9—H9B	110.0
H1B—C1—H1C	109.5	S2—C9—H9B	110.0
O2—C2—C1	111.0 (4)	H9A—C9—H9B	108.4
O2—C2—H2A	109.4	O4—C10—O3	125.0 (4)
C1—C2—H2A	109.4	O4—C10—C9	124.4 (4)
O2—C2—H2B	109.4	O3—C10—C9	110.6 (4)
C1—C2—H2B	109.4	O3—C11—C12	108.0 (4)
H2A—C2—H2B	108.0	O3—C11—H11A	110.1
O1—C3—O2	124.9 (4)	C12—C11—H11A	110.1
O1—C3—C4	123.6 (4)	O3—C11—H11B	110.1
O2—C3—C4	111.4 (3)	C12—C11—H11B	110.1
C5—C4—C3	129.3 (3)	H11A—C11—H11B	108.4
C5—C4—S1	111.1 (3)	C11—C12—H12A	109.5
C3—C4—S1	119.5 (3)	C11—C12—H12B	109.5
C4—C5—C6	113.1 (3)	H12A—C12—H12B	109.5
C4—C5—Br1	126.6 (3)	C11—C12—H12C	109.5
C6—C5—Br1	120.4 (3)	H12A—C12—H12C	109.5
C7—C6—C5	112.6 (3)	H12B—C12—H12C	109.5
C7—C6—C8	122.7 (4)		

C3—O2—C2—C1	-83.9 (5)	C5—C6—C7—S1	-0.2 (4)
C2—O2—C3—O1	-0.4 (6)	C8—C6—C7—S1	179.4 (3)
C2—O2—C3—C4	179.9 (3)	C5—C6—C7—S2	-178.6 (3)
O1—C3—C4—C5	-2.5 (7)	C8—C6—C7—S2	0.9 (6)
O2—C3—C4—C5	177.2 (4)	C4—S1—C7—C6	-0.1 (3)
O1—C3—C4—S1	179.4 (4)	C4—S1—C7—S2	178.3 (3)
O2—C3—C4—S1	-1.0 (5)	C9—S2—C7—C6	-178.3 (3)
C7—S1—C4—C5	0.4 (3)	C9—S2—C7—S1	3.5 (3)
C7—S1—C4—C3	178.9 (3)	C7—C6—C8—N1	-36 (12)
C3—C4—C5—C6	-178.9 (4)	C5—C6—C8—N1	144 (11)
S1—C4—C5—C6	-0.6 (4)	C7—S2—C9—C10	-169.7 (3)
C3—C4—C5—Br1	0.6 (6)	C11—O3—C10—O4	3.5 (6)
S1—C4—C5—Br1	178.9 (2)	C11—O3—C10—C9	-176.8 (4)
C4—C5—C6—C7	0.5 (5)	S2—C9—C10—O4	-5.4 (6)
Br1—C5—C6—C7	-179.0 (3)	S2—C9—C10—O3	174.9 (3)
C4—C5—C6—C8	-179.0 (4)	C10—O3—C11—C12	178.9 (4)
Br1—C5—C6—C8	1.5 (5)		

supporting information

Acta Cryst. (2013). E69, o848 [doi:10.1107/S1600536813010611]

Ethyl 3-bromo-4-cyano-5-[(2-ethoxy-2-oxoethyl)sulfanyl]thiophene-2-carboxylate


Xiuping Li, Xiaochuan Jia and Jing Li

S1. Comment


3-Amino-4-cyano-5-ethoxycarbonylmethylsulfanyl-thiophene-2-carboxylic acid ethyl ester is an important intermediate compound for synthesis of thienothienopyrimidines derivatives, which are thought to be potential biological active compounds or pharmaceuticals (Liu, *et al.*, 2008). We obtained the title compound by the Sandmeyer reaction from compound 3-Amino-4-cyano-5-ethoxycarbonylmethylsulfanyl-thiophene-2-carboxylic acid ethyl ester. The crystal for X-ray crystal structure analysis was obtained by recrystallizing the title compound in petroleum ether. In the crystal, the thiophene ring together with its four adjoint groups, *i.e.* CN, Br, S—CH₂ and COO⁻, was located at one perfect plane, which is consistent with the crystal of 3-Amino-4-cyano-5-ethoxycarbonylmethylsulfanyl-thiophene-2-carboxylic acid ethyl ester reported in the literature (Padmavathi *et al.*, 2011). The title compound crystal demonstrated a crystal system of monoclinic and a space group of P2(1)/c. There existed hydrogen bond with length of 2.554 Å between one of —SCH₂ H atom and the O atom of carbonyl adjoined with thiophene ring of neighbor molecule.

S2. Experimental

To a solution of 3-Amino-4-cyano-5-ethoxycarbonylmethylsulfanyl -thiophene-2-carboxylic acid ethyl ester (1.57 g, 5 mmol) in 70% H₂SO₄ (13 mL) was added NaNO₂ (0.4 g, 5.7 mmol) in 5 minutes under ice water temperature. After addition, the solution was stirred for 30 min at room temperature. Then, the reaction mixture was transferred to HBr solution containing CuBr (1 g, 7 mmol). After standing overnight, water (100 mL) was added. The precipitate was collected by filtration and recrystallized from petroleum ether to afford the title compound as colourless crystals, yield 50%.

**Figure 1**

The structure of the title compound showing 30% probability ellipsoids.

**Figure 2**

Packing structure of the title compound.

Ethyl 3-bromo-4-cyano-5-[(2-ethoxy-2-oxoethyl)sulfanyl]thiophene-2-carboxylate

Crystal data

C₁₂H₁₂BrNO₄S₂ $M_r = 378.26$ Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

 $a = 8.5896$ (17) Å $b = 10.837$ (2) Å $c = 16.584$ (3) Å $\beta = 99.44$ (3)° $V = 1522.8$ (5) Å³ $Z = 4$ $F(000) = 760$ $D_x = 1.650$ Mg m⁻³Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3404 reflections

 $\theta = 3.1$ – 27.9 ° $\mu = 2.98$ mm⁻¹ $T = 293$ K

Block, colorless

 $0.20 \times 0.18 \times 0.12$ mm

Data collection

Rigaku Saturn
diffractometer

Radiation source: rotating anode

Confocal monochromator

 ω scansAbsorption correction: multi-scan
(*CrystalClear*; Rigaku, 2005) $T_{\min} = 0.587$, $T_{\max} = 0.716$

15211 measured reflections

3596 independent reflections

2224 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.067$ $\theta_{\text{max}} = 27.9$ °, $\theta_{\text{min}} = 3.1$ ° $h = -11 \rightarrow 11$ $k = -14 \rightarrow 14$ $l = -21 \rightarrow 21$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.052$ $wR(F^2) = 0.157$ $S = 1.00$

3596 reflections

184 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0817P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\text{max}} = 0.003$ $\Delta\rho_{\text{max}} = 0.43$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.71$ e Å⁻³Extinction correction: *SHELXL97* (Sheldrick,
2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.018 (2)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	1.34684 (5)	-0.05401 (5)	0.13435 (3)	0.0723 (3)
S1	0.95059 (11)	0.19092 (9)	0.04312 (6)	0.0421 (3)
S2	0.75972 (12)	0.13383 (10)	0.17935 (7)	0.0505 (3)

O1	1.3519 (4)	0.1042 (3)	-0.0246 (2)	0.0703 (9)
O2	1.1477 (3)	0.2261 (3)	-0.07630 (18)	0.0577 (8)
O3	0.4240 (3)	0.3685 (3)	0.10494 (18)	0.0559 (8)
O4	0.4796 (4)	0.2478 (3)	0.2147 (2)	0.0744 (10)
N1	1.0569 (5)	-0.0869 (4)	0.2904 (3)	0.0784 (13)
C1	1.3411 (7)	0.3617 (5)	-0.1215 (3)	0.0825 (16)
H1A	1.2890	0.4277	-0.0980	0.124*
H1B	1.3794	0.3912	-0.1692	0.124*
H1C	1.4282	0.3325	-0.0823	0.124*
C2	1.2285 (6)	0.2598 (5)	-0.1446 (3)	0.0667 (13)
H2A	1.1510	0.2847	-0.1911	0.080*
H2B	1.2845	0.1885	-0.1607	0.080*
C3	1.2245 (5)	0.1481 (4)	-0.0215 (3)	0.0468 (10)
C4	1.1325 (4)	0.1209 (3)	0.0445 (2)	0.0420 (9)
C5	1.1667 (5)	0.0433 (3)	0.1088 (3)	0.0440 (9)
C6	1.0474 (4)	0.0400 (3)	0.1587 (2)	0.0427 (9)
C7	0.9215 (4)	0.1161 (3)	0.1300 (2)	0.0390 (8)
C8	1.0542 (5)	-0.0323 (4)	0.2309 (3)	0.0551 (11)
C9	0.6413 (4)	0.2381 (4)	0.1105 (2)	0.0473 (10)
H9A	0.7050	0.3072	0.0979	0.057*
H9B	0.6003	0.1960	0.0599	0.057*
C10	0.5069 (4)	0.2838 (4)	0.1508 (3)	0.0464 (10)
C11	0.2964 (6)	0.4242 (5)	0.1395 (3)	0.0632 (13)
H11A	0.3373	0.4630	0.1914	0.076*
H11B	0.2205	0.3617	0.1488	0.076*
C12	0.2186 (6)	0.5190 (5)	0.0800 (4)	0.0903 (18)
H12A	0.2957	0.5782	0.0691	0.135*
H12B	0.1371	0.5603	0.1029	0.135*
H12C	0.1733	0.4791	0.0299	0.135*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0499 (4)	0.0728 (4)	0.0911 (5)	0.0186 (2)	0.0023 (3)	0.0000 (3)
S1	0.0381 (5)	0.0419 (5)	0.0472 (6)	0.0045 (4)	0.0093 (4)	0.0038 (4)
S2	0.0452 (6)	0.0526 (6)	0.0570 (7)	0.0040 (5)	0.0186 (5)	0.0114 (5)
O1	0.0539 (19)	0.088 (2)	0.075 (2)	0.0161 (18)	0.0267 (16)	0.0067 (19)
O2	0.0532 (18)	0.0659 (19)	0.0574 (18)	0.0027 (15)	0.0195 (14)	0.0111 (16)
O3	0.0464 (16)	0.0658 (19)	0.0585 (19)	0.0130 (15)	0.0177 (14)	-0.0037 (15)
O4	0.074 (2)	0.088 (2)	0.070 (2)	0.0212 (18)	0.0363 (18)	0.0164 (19)
N1	0.073 (3)	0.077 (3)	0.087 (3)	0.012 (2)	0.020 (2)	0.035 (3)
C1	0.100 (4)	0.089 (4)	0.064 (3)	-0.022 (3)	0.030 (3)	0.002 (3)
C2	0.076 (3)	0.072 (3)	0.056 (3)	-0.004 (3)	0.022 (2)	0.008 (2)
C3	0.040 (2)	0.049 (2)	0.052 (2)	-0.0011 (19)	0.0102 (18)	-0.005 (2)
C4	0.034 (2)	0.039 (2)	0.054 (2)	0.0016 (16)	0.0101 (17)	-0.0058 (18)
C5	0.036 (2)	0.041 (2)	0.054 (2)	-0.0013 (16)	0.0032 (18)	-0.0033 (18)
C6	0.041 (2)	0.039 (2)	0.047 (2)	-0.0030 (17)	0.0034 (18)	0.0045 (17)
C7	0.036 (2)	0.0346 (19)	0.046 (2)	-0.0006 (16)	0.0064 (16)	-0.0012 (16)

C8	0.045 (2)	0.051 (2)	0.070 (3)	0.006 (2)	0.011 (2)	0.011 (2)
C9	0.036 (2)	0.059 (3)	0.048 (2)	0.0034 (19)	0.0114 (18)	0.003 (2)
C10	0.039 (2)	0.053 (2)	0.049 (2)	0.0003 (19)	0.0099 (18)	-0.003 (2)
C11	0.050 (3)	0.070 (3)	0.074 (3)	0.018 (2)	0.023 (2)	-0.008 (3)
C12	0.068 (3)	0.097 (4)	0.108 (5)	0.036 (3)	0.021 (3)	0.015 (4)

Geometric parameters (Å, °)

Br1—C5	1.864 (4)	C2—H2A	0.9700
S1—C7	1.707 (4)	C2—H2B	0.9700
S1—C4	1.734 (4)	C3—C4	1.480 (6)
S2—C7	1.735 (4)	C4—C5	1.352 (5)
S2—C9	1.798 (4)	C5—C6	1.418 (6)
O1—C3	1.202 (5)	C6—C7	1.381 (5)
O2—C3	1.335 (5)	C6—C8	1.425 (6)
O2—C2	1.468 (5)	C9—C10	1.509 (5)
O3—C10	1.323 (5)	C9—H9A	0.9700
O3—C11	1.449 (5)	C9—H9B	0.9700
O4—C10	1.188 (5)	C11—C12	1.504 (7)
N1—C8	1.147 (6)	C11—H11A	0.9700
C1—C2	1.477 (7)	C11—H11B	0.9700
C1—H1A	0.9600	C12—H12A	0.9600
C1—H1B	0.9600	C12—H12B	0.9600
C1—H1C	0.9600	C12—H12C	0.9600
C7—S1—C4	92.15 (19)	C5—C6—C8	124.7 (4)
C7—S2—C9	100.55 (18)	C6—C7—S1	111.1 (3)
C3—O2—C2	116.1 (3)	C6—C7—S2	123.1 (3)
C10—O3—C11	115.6 (3)	S1—C7—S2	125.8 (2)
C2—C1—H1A	109.5	N1—C8—C6	177.4 (5)
C2—C1—H1B	109.5	C10—C9—S2	108.6 (3)
H1A—C1—H1B	109.5	C10—C9—H9A	110.0
C2—C1—H1C	109.5	S2—C9—H9A	110.0
H1A—C1—H1C	109.5	C10—C9—H9B	110.0
H1B—C1—H1C	109.5	S2—C9—H9B	110.0
O2—C2—C1	111.0 (4)	H9A—C9—H9B	108.4
O2—C2—H2A	109.4	O4—C10—O3	125.0 (4)
C1—C2—H2A	109.4	O4—C10—C9	124.4 (4)
O2—C2—H2B	109.4	O3—C10—C9	110.6 (4)
C1—C2—H2B	109.4	O3—C11—C12	108.0 (4)
H2A—C2—H2B	108.0	O3—C11—H11A	110.1
O1—C3—O2	124.9 (4)	C12—C11—H11A	110.1
O1—C3—C4	123.6 (4)	O3—C11—H11B	110.1
O2—C3—C4	111.4 (3)	C12—C11—H11B	110.1
C5—C4—C3	129.3 (3)	H11A—C11—H11B	108.4
C5—C4—S1	111.1 (3)	C11—C12—H12A	109.5
C3—C4—S1	119.5 (3)	C11—C12—H12B	109.5
C4—C5—C6	113.1 (3)	H12A—C12—H12B	109.5

C4—C5—Br1	126.6 (3)	C11—C12—H12C	109.5
C6—C5—Br1	120.4 (3)	H12A—C12—H12C	109.5
C7—C6—C5	112.6 (3)	H12B—C12—H12C	109.5
C7—C6—C8	122.7 (4)		
C3—O2—C2—C1	-83.9 (5)	C5—C6—C7—S1	-0.2 (4)
C2—O2—C3—O1	-0.4 (6)	C8—C6—C7—S1	179.4 (3)
C2—O2—C3—C4	179.9 (3)	C5—C6—C7—S2	-178.6 (3)
O1—C3—C4—C5	-2.5 (7)	C8—C6—C7—S2	0.9 (6)
O2—C3—C4—C5	177.2 (4)	C4—S1—C7—C6	-0.1 (3)
O1—C3—C4—S1	179.4 (4)	C4—S1—C7—S2	178.3 (3)
O2—C3—C4—S1	-1.0 (5)	C9—S2—C7—C6	-178.3 (3)
C7—S1—C4—C5	0.4 (3)	C9—S2—C7—S1	3.5 (3)
C7—S1—C4—C3	178.9 (3)	C7—C6—C8—N1	-36 (12)
C3—C4—C5—C6	-178.9 (4)	C5—C6—C8—N1	144 (11)
S1—C4—C5—C6	-0.6 (4)	C7—S2—C9—C10	-169.7 (3)
C3—C4—C5—Br1	0.6 (6)	C11—O3—C10—O4	3.5 (6)
S1—C4—C5—Br1	178.9 (2)	C11—O3—C10—C9	-176.8 (4)
C4—C5—C6—C7	0.5 (5)	S2—C9—C10—O4	-5.4 (6)
Br1—C5—C6—C7	-179.0 (3)	S2—C9—C10—O3	174.9 (3)
C4—C5—C6—C8	-179.0 (4)	C10—O3—C11—C12	178.9 (4)
Br1—C5—C6—C8	1.5 (5)		
