

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4-Phenyl-1,2,4-triazaspiro[4.5]dec-1-ene-3-thione

 Mehmet Akkurt,^a Joel T. Mague,^b Shaaban K. Mohamed,^{c,d} Alaa A. Hassan^d and Mustafa R. Albayati^{e*}

^aDepartment of Physics, Faculty of Sciences, Erciyes University, 38039 Kayseri, Turkey, ^bDepartment of Chemistry, Tulane University, New Orleans, LA 70118, USA, ^cChemistry and Environmental Division, Manchester Metropolitan University, Manchester M1 5GD, England, ^dChemistry Department, Faculty of Science, Mini University, 61519 El-Minia, Egypt, and ^eKirkuk University, College of Science, Department of Chemistry, Kirkuk, Iraq
Correspondence e-mail: shaabankamel@yahoo.com


Received 1 July 2013; accepted 10 July 2013

 Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.042; wR factor = 0.104; data-to-parameter ratio = 20.6.

In the title compound, $\text{C}_{13}\text{H}_{15}\text{N}_3\text{S}$, the 4,5-dihydro-3*H*-1,2,4-triazole ring is nearly planar [maximum deviation = 0.020 (1) Å], while the cyclohexane ring adopts a chair conformation. The dihedral angle between the 4,5-dihydro-3*H*-1,2,4-triazole ring and the phenyl ring is 74.68 (7)°. No specific intermolecular interactions are discerned in the crystal packing.

Related literature

For wide-spectrum medicinal applications of spiro compounds incorporating heterocyclic substructures, see: Patil *et al.* (2010); Pawar *et al.* (2009); Thadhaney *et al.* (2010); Chin *et al.* (2008); Wang *et al.* (2007); Chande *et al.* (2005); Obniska *et al.* (2006); Kamiński *et al.* (2008); Sarma *et al.* (2010); Shimakawa *et al.* (2003). For ring-puckering parameters, see: Cremer & Pople (1975).



Experimental

Crystal data

$\text{C}_{13}\text{H}_{15}\text{N}_3\text{S}$	$V = 2465.5$ (4) Å ³
$M_r = 245.34$	$Z = 8$
Orthorhombic, <i>Pbca</i>	Mo $K\alpha$ radiation
$a = 9.4952$ (9) Å	$\mu = 0.24$ mm ⁻¹
$b = 7.4845$ (7) Å	$T = 150$ K
$c = 34.692$ (3) Å	$0.28 \times 0.22 \times 0.17$ mm

Data collection

Bruker SMART APEX CCD diffractometer	41202 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2013)	3168 independent reflections
$T_{\min} = 0.810$, $T_{\max} = 0.960$	2899 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.046$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$	154 parameters
$wR(F^2) = 0.104$	H-atom parameters constrained
$S = 1.10$	$\Delta\rho_{\text{max}} = 0.40$ e Å ⁻³
3168 reflections	$\Delta\rho_{\text{min}} = -0.25$ e Å ⁻³

Data collection: *APEX2* (Bruker, 2013); cell refinement: *S SAINT* (Bruker, 2013); data reduction: *S SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *publCIF* (Westrip, 2010).

Tulane University, Erciyes University and Minia University are gratefully acknowledged for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK5237).

References

- Bruker (2013). *APEX2*, *S SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chande, M. S., Verma, R. S., Barve, P. A., Khanwelkar, R. R., Vaidya, R. B. & Ajaikumar, K. B. (2005). *Eur. J. Med. Chem.* **40**, 1143–1148.
- Chin, Y.-W., Salim, A. A., Su, B.-N., Mi, Q., Chai, H.-B., Riswan, S., Kardono, L. B. S., Ruskandi, A., Farnsworth, N. R., Swanson, S. M. & Kinghorn, A. D. (2008). *J. Nat. Prod.* **3**, 390–395.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Kamiński, K., Obniska, J. & Dybala, M. (2008). *Eur. J. Med. Chem.* **43**, 53–61.
- Obniska, J., Kamiński, K. & Tatarczyńska, E. (2006). *Pharmacol. Rep.* **58**, 207–214.
- Patil, B. S., Krishnamurthy, G., BhojyaNaik, H. S., Lathe, P. R. & Ghate, M. (2010). *Eur. J. Med. Chem.* **45**, 3329–3334.
- Pawar, M. J., Burungale, A. B. & Karale, B. K. (2009). *ARKIVOC*, **XIII**, 97–107.
- Sarma, B. K., Manna, D., Minoura, M. & Muges, G. (2010). *J. Am. Chem. Soc.* **132**, 5364–5374.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Shimakawa, S., Yoshida, Y. & Niki, E. (2003). *Lipids*, **38**, 225–231.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Thadhaney, B., Sain, D., Pernawat, G. & Talesara, G. L. (2010). *Indian J. Chem. Sect. B*, **49**, 368–373.
- Wang, W.-L., Zhu, T.-J., Tao, H.-W., Lu, Z.-Y., Fang, Y.-C., Gu, Q.-Q. & Zhu, W.-M. (2007). *Chem. Biodivers.* **4**, 2913–2919.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2013). E69, o1259 [doi:10.1107/S1600536813019120]

4-Phenyl-1,2,4-triazaspiro[4.5]dec-1-ene-3-thione

Mehmet Akkurt, Joel T. Mague, Shaaban K. Mohamed, Alaa A. Hassan and Mustafa R. Albayati

S1. Comment

Heterocyclic compounds such as 1,2,4-triazoles exhibit a wide range of biological activities (Patil *et al.*, 2010). Several spiro-compounds incorporating different heterocyclic structures have showed significant industrial and pharmaceutical applications such as anti-microbial (Pawar *et al.*, 2009; Thadhaney *et al.*, 2010), anti-cancer (Chin *et al.*, 2008; Wang *et al.*, 2007), anti-tubercular (Chande *et al.*, 2005) and anti-convulsant activities (Obniska *et al.*, 2006; Kamiński *et al.*, 2008) as well as functioning as antioxidants (Sarma *et al.*, 2010; Shimakawa *et al.*, 2003). In view of such facts and as part of our on-going study on the synthesis of bio-active molecules, we herein report the synthesis and crystal structure of the title compound (I).

As shown in Fig. 1, the 4,5-dihydro-3*H*-1,2,4-triazole ring (N1–N3/C1/C2) of (I) is nearly planar with a maximum deviation of 0.020 (1) Å for N1 and it makes a dihedral angle of 74.68 (7)° with the phenyl ring (C8–C13). The cyclohexane ring (C2–C7) adopts a chair conformation with the puckering parameters (Cremer & Pople, 1975) of $Q_T = 0.5610$ (14) Å, $\theta = 1.74$ (14)° and $\varphi = 342$ (4)°.

The stabilization of the molecular packing of (I) is assisted by a number of non-bonded forces including van der Waals.


S2. Experimental

A solution of 2-cyclohexylidene-*N*-phenylhydrazinecarbothioamide (1 mmol) in dry ethyl acetate (15 ml) was added drop wise over 2 h to a stirred solution of 4,5-dichloro-3,6-dioxocyclohexa-1,4-diene-1,2-dicarbonitrile (227 mg, 1 mmol) in dry ethyl acetate (10 ml). The pink coloration of the reaction mixture turned quickly to red brown and the mixture was left to stand at room temperature for 48 h. The precipitated DDQ-H₂ [JM1] was filtered off and washed with few drops of ethyl acetate. The filtrate was collected, concentrated under vacuum and left at room temperature to afford the title compound as red brown crystals suitable for X-ray diffraction.

IR: 3052 (Ar–CH), 2941, 2863 (Alk – CH), 1594 (Ar–C=C), 1350 (C=S); ¹H-NMR (CDCl₃) 1.26, 1.65, 1.94 and 2.20 (10*H*, cyclohexane–CH₂), 7.707, 7.51, 7.55 (5*H*, Ar–H); ¹³C-NMR (CDCl₃) 23.58, 24.52 and 33.90 (cyclohexane–CH₂), 112.19 (spiro C_x b), 127.81, 129.87, 130.22 (Ar–CH), 135.07 (Ar-c), 187.83 (C=S).

S3. Refinement

All H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms with C–H = 0.95 and 0.99 Å, with $U_{iso}(H) = 1.2 U_{iso}(C)$.

**Figure 1**

View of the title compound (I) with the atom numbering scheme. Displacement ellipsoids for non-H atoms are drawn at the 50% probability level.

4-Phenyl-1,2,4-triazaspiro[4.5]dec-1-ene-3-thione

Crystal data

$C_{13}H_{15}N_3S$

$M_r = 245.34$

Orthorhombic, *Pbca*

Hall symbol: -P 2ac 2ab

$a = 9.4952(9) \text{ \AA}$

$b = 7.4845(7) \text{ \AA}$

$c = 34.692(3) \text{ \AA}$

$V = 2465.5(4) \text{ \AA}^3$

$Z = 8$

$F(000) = 1040$

$D_x = 1.322 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 9973 reflections

$\theta = 2.5\text{--}28.6^\circ$

$\mu = 0.24 \text{ mm}^{-1}$

$T = 150 \text{ K}$

Block, red-brown

$0.28 \times 0.22 \times 0.17 \text{ mm}$

Data collection

Bruker SMART APEX CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: $8.3660 \text{ pixels mm}^{-1}$

φ and ω scans

Absorption correction: multi-scan
(*SADABS*; Bruker, 2013)

$T_{\min} = 0.810$, $T_{\max} = 0.960$

41202 measured reflections

3168 independent reflections

2899 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.046$

$\theta_{\max} = 28.7^\circ$, $\theta_{\min} = 2.4^\circ$

$h = -12 \rightarrow 12$

$k = -10 \rightarrow 10$

$l = -46 \rightarrow 46$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.042$	H-atom parameters constrained
$wR(F^2) = 0.104$	$W = 1/[\Sigma^2(FO^2) + (0.0483P)^2 + 1.0447P]$
$S = 1.10$	where $P = (FO^2 + 2FC^2)/3$
3168 reflections	$(\Delta/\sigma)_{\max} = 0.001$
154 parameters	$\Delta\rho_{\max} = 0.40 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.25 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement on F^2 for ALL reflections except those flagged by the user for potential systematic errors. Weighted R -factors wR and all goodnesses of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The observed criterion of $F^2 > \sigma(F^2)$ is used only for calculating $-R$ -factor-obs *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.53002 (4)	1.40579 (5)	0.84564 (2)	0.0310 (1)
N1	0.50442 (11)	1.06645 (14)	0.87249 (3)	0.0199 (3)
N2	0.53364 (11)	1.27385 (15)	0.91860 (3)	0.0252 (3)
N3	0.51796 (11)	1.13018 (15)	0.93661 (3)	0.0233 (3)
C1	0.52216 (12)	1.24250 (17)	0.87698 (4)	0.0219 (3)
C2	0.49469 (12)	0.97820 (16)	0.91027 (3)	0.0188 (3)
C3	0.60868 (13)	0.83732 (18)	0.91727 (4)	0.0241 (3)
C4	0.59589 (15)	0.75735 (19)	0.95773 (4)	0.0282 (4)
C5	0.44880 (15)	0.68348 (18)	0.96539 (4)	0.0280 (4)
C6	0.33484 (14)	0.82248 (17)	0.95781 (3)	0.0240 (3)
C7	0.34706 (13)	0.90101 (16)	0.91725 (3)	0.0209 (3)
C8	0.47802 (13)	0.97938 (17)	0.83642 (3)	0.0208 (3)
C9	0.58094 (15)	0.87142 (18)	0.82019 (4)	0.0273 (4)
C10	0.55240 (17)	0.7858 (2)	0.78557 (4)	0.0346 (4)
C11	0.42447 (19)	0.8099 (2)	0.76734 (4)	0.0377 (5)
C12	0.32316 (18)	0.9190 (2)	0.78367 (4)	0.0369 (4)
C13	0.34905 (15)	1.00429 (19)	0.81847 (4)	0.0283 (4)
H3A	0.70270	0.89270	0.91420	0.0290*
H3B	0.59970	0.74110	0.89790	0.0290*
H4A	0.66580	0.66010	0.96070	0.0340*
H4B	0.61770	0.85060	0.97710	0.0340*
H5A	0.43240	0.57840	0.94860	0.0340*
H5B	0.44270	0.64330	0.99250	0.0340*
H6A	0.34290	0.91980	0.97700	0.0290*
H6B	0.24110	0.76650	0.96090	0.0290*

H7A	0.27600	0.99650	0.91390	0.0250*
H7B	0.32760	0.80660	0.89800	0.0250*
H9	0.66940	0.85640	0.83260	0.0330*
H10	0.62130	0.71000	0.77430	0.0410*
H11	0.40610	0.75150	0.74360	0.0450*
H12	0.23540	0.93560	0.77100	0.0440*
H13	0.27940	1.07870	0.82980	0.0340*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0373 (2)	0.0214 (2)	0.0344 (2)	-0.0050 (1)	0.0002 (1)	0.0067 (1)
N1	0.0234 (5)	0.0179 (5)	0.0183 (5)	-0.0008 (4)	-0.0003 (4)	0.0003 (4)
N2	0.0254 (5)	0.0214 (5)	0.0287 (6)	-0.0028 (4)	0.0010 (4)	-0.0045 (4)
N3	0.0255 (5)	0.0209 (5)	0.0235 (5)	-0.0009 (4)	-0.0004 (4)	-0.0056 (4)
C1	0.0190 (5)	0.0195 (6)	0.0272 (6)	-0.0014 (4)	0.0011 (4)	-0.0011 (5)
C2	0.0230 (5)	0.0172 (6)	0.0162 (5)	-0.0001 (4)	-0.0014 (4)	-0.0013 (4)
C3	0.0240 (6)	0.0248 (6)	0.0235 (6)	0.0054 (5)	-0.0030 (5)	-0.0015 (5)
C4	0.0337 (7)	0.0276 (7)	0.0234 (6)	0.0075 (6)	-0.0077 (5)	0.0000 (5)
C5	0.0406 (7)	0.0218 (6)	0.0215 (6)	0.0034 (5)	-0.0025 (5)	0.0022 (5)
C6	0.0308 (6)	0.0208 (6)	0.0204 (6)	-0.0017 (5)	0.0018 (5)	0.0011 (4)
C7	0.0216 (5)	0.0200 (5)	0.0210 (6)	-0.0004 (5)	-0.0011 (4)	0.0011 (4)
C8	0.0263 (6)	0.0189 (6)	0.0172 (5)	-0.0034 (5)	0.0008 (4)	0.0014 (4)
C9	0.0313 (7)	0.0269 (6)	0.0236 (6)	0.0007 (5)	0.0058 (5)	0.0003 (5)
C10	0.0504 (9)	0.0281 (7)	0.0252 (7)	-0.0031 (6)	0.0146 (6)	-0.0030 (5)
C11	0.0607 (10)	0.0340 (8)	0.0185 (6)	-0.0168 (7)	0.0013 (6)	-0.0025 (5)
C12	0.0434 (8)	0.0404 (8)	0.0268 (7)	-0.0093 (7)	-0.0107 (6)	0.0008 (6)
C13	0.0302 (7)	0.0298 (7)	0.0250 (6)	-0.0013 (5)	-0.0029 (5)	0.0006 (5)

Geometric parameters (Å, °)

S1—C1	1.6375 (14)	C11—C12	1.383 (2)
N1—C1	1.3375 (17)	C12—C13	1.388 (2)
N1—C2	1.4706 (15)	C3—H3A	0.9900
N1—C8	1.4330 (15)	C3—H3B	0.9900
N2—N3	1.2525 (16)	C4—H4A	0.9900
N2—C1	1.4669 (17)	C4—H4B	0.9900
N3—C2	1.4757 (16)	C5—H5A	0.9900
C2—C3	1.5305 (17)	C5—H5B	0.9900
C2—C7	1.5354 (17)	C6—H6A	0.9900
C3—C4	1.531 (2)	C6—H6B	0.9900
C4—C5	1.525 (2)	C7—H7A	0.9900
C5—C6	1.5239 (19)	C7—H7B	0.9900
C6—C7	1.5293 (15)	C9—H9	0.9500
C8—C9	1.3874 (19)	C10—H10	0.9500
C8—C13	1.3864 (19)	C11—H11	0.9500
C9—C10	1.388 (2)	C12—H12	0.9500
C10—C11	1.381 (2)	C13—H13	0.9500

C1—N1—C2	110.28 (10)	H3A—C3—H3B	108.00
C1—N1—C8	124.87 (11)	C3—C4—H4A	109.00
C2—N1—C8	124.27 (10)	C3—C4—H4B	109.00
N3—N2—C1	110.17 (10)	C5—C4—H4A	109.00
N2—N3—C2	111.76 (10)	C5—C4—H4B	109.00
S1—C1—N1	131.58 (11)	H4A—C4—H4B	108.00
S1—C1—N2	122.06 (10)	C4—C5—H5A	109.00
N1—C1—N2	106.36 (11)	C4—C5—H5B	109.00
N1—C2—N3	101.32 (9)	C6—C5—H5A	109.00
N1—C2—C3	113.98 (10)	C6—C5—H5B	109.00
N1—C2—C7	111.53 (9)	H5A—C5—H5B	108.00
N3—C2—C3	109.08 (9)	C5—C6—H6A	109.00
N3—C2—C7	109.21 (9)	C5—C6—H6B	109.00
C3—C2—C7	111.19 (10)	C7—C6—H6A	109.00
C2—C3—C4	111.03 (10)	C7—C6—H6B	109.00
C3—C4—C5	111.97 (11)	H6A—C6—H6B	108.00
C4—C5—C6	111.88 (11)	C2—C7—H7A	109.00
C5—C6—C7	111.55 (10)	C2—C7—H7B	109.00
C2—C7—C6	111.04 (10)	C6—C7—H7A	110.00
N1—C8—C9	119.74 (11)	C6—C7—H7B	109.00
N1—C8—C13	119.05 (11)	H7A—C7—H7B	108.00
C9—C8—C13	121.21 (11)	C8—C9—H9	120.00
C8—C9—C10	118.85 (13)	C10—C9—H9	121.00
C9—C10—C11	120.50 (14)	C9—C10—H10	120.00
C10—C11—C12	120.08 (13)	C11—C10—H10	120.00
C11—C12—C13	120.31 (15)	C10—C11—H11	120.00
C8—C13—C12	119.04 (13)	C12—C11—H11	120.00
C2—C3—H3A	109.00	C11—C12—H12	120.00
C2—C3—H3B	109.00	C13—C12—H12	120.00
C4—C3—H3A	110.00	C8—C13—H13	121.00
C4—C3—H3B	109.00	C12—C13—H13	120.00
C2—N1—C1—S1	-175.88 (10)	N2—N3—C2—C7	-116.28 (11)
C2—N1—C1—N2	3.52 (12)	N1—C2—C3—C4	177.48 (10)
C8—N1—C1—S1	-4.33 (19)	N3—C2—C3—C4	65.07 (13)
C8—N1—C1—N2	175.07 (10)	C7—C2—C3—C4	-55.43 (13)
C1—N1—C2—N3	-3.14 (12)	N1—C2—C7—C6	-175.60 (10)
C1—N1—C2—C3	-120.14 (11)	N3—C2—C7—C6	-64.44 (12)
C1—N1—C2—C7	112.94 (11)	C3—C2—C7—C6	55.98 (13)
C8—N1—C2—N3	-174.76 (10)	C2—C3—C4—C5	54.36 (15)
C8—N1—C2—C3	68.24 (14)	C3—C4—C5—C6	-53.80 (15)
C8—N1—C2—C7	-58.67 (14)	C4—C5—C6—C7	54.12 (14)
C1—N1—C8—C9	110.43 (14)	C5—C6—C7—C2	-55.15 (13)
C1—N1—C8—C13	-69.89 (16)	N1—C8—C9—C10	178.89 (12)
C2—N1—C8—C9	-79.16 (15)	C13—C8—C9—C10	-0.8 (2)
C2—N1—C8—C13	100.51 (14)	N1—C8—C13—C12	-179.57 (12)
C1—N2—N3—C2	0.53 (13)	C9—C8—C13—C12	0.1 (2)

supporting information

N3—N2—C1—S1	176.89 (9)	C8—C9—C10—C11	1.0 (2)
N3—N2—C1—N1	-2.58 (13)	C9—C10—C11—C12	-0.5 (2)
N2—N3—C2—N1	1.50 (12)	C10—C11—C12—C13	-0.2 (2)
N2—N3—C2—C3	122.02 (11)	C11—C12—C13—C8	0.4 (2)
