

1,1',4,5-Tetrahydrotrispiro[1,3,2-diazaphosphole-2,2'-[1,3,5,2,4,6]triazatriphosphinine-4',6''-dibenzo[*d,f*][1,3,2]-dioxaphosphepine-6',6'''-dibenzo[*d,f*]-[1,3,2]dioxaphosphepine] acetone monosolvate

Krystal R. Fontenot,^a Michael W. Easson,^{a*} Frank R. Fronczek^b and Brian D. Condon^a

^aUnited States Department of Agriculture Cotton Chemistry and Utilization, 1100 Robert E. Lee Blvd, New Orleans, LA 70124, USA, and ^bDepartment of Chemistry, Louisiana State University, Baton Rouge, LA 70803, USA
Correspondence e-mail: michael.easson@ars.usda.gov


Received 14 August 2013; accepted 23 August 2013

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.037; wR factor = 0.102; data-to-parameter ratio = 13.1.

The title compound, $\text{C}_{26}\text{H}_{22}\text{N}_5\text{O}_4\text{P}_3 \cdot \text{C}_3\text{H}_6\text{O}$, has been achieved in a two-step synthesis that does not require chromatography. This molecule contains a seven-membered spirocyclic ring at two P-atom positions and a five-membered ring containing new P–N bonds at the other P-atom position. Endocyclic torsion angles about the central biphenyl C–C bonds are -41.5 (3) and -44.4 (3)°, and P–N bonds of the central P_3N_3 ring are within the range 1.5665 (17)–1.6171 (17) Å, while the P–O distances are in the range 1.5940 (14)–1.6041 (14) Å. One N–H group makes an intermolecular N–H···N hydrogen bond, forming centrosymmetric dimers, while the other N–H group makes an N–H···O hydrogen bond to the acetone solvent molecule. The crystal was a two-component non-merohedral twin with ratio 0.811/0.189.

Related literature

For phosphazene-based flame retardants, see: Bakos *et al.* (1982); Drews & Barker (1985). For related structures, bond angles and lengths, see: Allcock (1972); Ciftci *et al.* (2013). For the geometry of phosphazene rings, see: Olthof (1969); Barclay *et al.* (2002). For the synthesis, see: Allen (1991); Carriedo *et al.* (1996). For related structures, see: Chandrasekhar *et al.* (2007; 2011; 2012); Harmjanz *et al.* (2004). For graph-set analysis, see: Etter (1990). For ring asymmetry parameters, see: Duax *et al.* (1976).



Experimental

Crystal data

$\text{C}_{26}\text{H}_{22}\text{N}_5\text{O}_4\text{P}_3 \cdot \text{C}_3\text{H}_6\text{O}$
 $M_r = 619.47$
Monoclinic, $P2_1/c$
 $a = 9.4901$ (9) Å
 $b = 22.9466$ (19) Å
 $c = 13.1776$ (13) Å
 $\beta = 97.978$ (6)°

$V = 2841.9$ (5) Å³
 $Z = 4$
Cu $K\alpha$ radiation
 $\mu = 2.34$ mm⁻¹
 $T = 100$ K
 $0.19 \times 0.12 \times 0.03$ mm

Data collection

Bruker Kappa APEXII DUO CCD diffractometer
Absorption correction: multi-scan (*TWINABS*; Sheldrick, 2004)
 $T_{\min} = 0.664$, $T_{\max} = 0.933$

33208 measured reflections
5065 independent reflections
4276 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.051$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.102$
 $S = 1.08$
5065 reflections
387 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.40$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.39$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> –H··· <i>A</i>	<i>D</i> –H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> –H··· <i>A</i>
N4–H4N···O5	0.82 (3)	2.22 (3)	2.991 (2)	158 (2)
N5–H5N···N3 ⁱ	0.83 (3)	2.53 (3)	3.285 (2)	151 (2)

Symmetry code: (i) $-x + 2, -y + 1, -z + 1$.

Data collection: *APEX2* (Bruker, 2011); cell refinement: *SAINT* (Bruker, 2011); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *SHELXL97*.

We would like to thank Dr Casey Grimm for the MS analysis. Upgrade of the diffractometer was made possible by grant No. LEQSF(2011–12)-ENH-TR-01, administered by the Louisiana Board of Regents.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JJ2173).

References

- Allcock, H. R. (1972). In *Phosphorus-Nitrogen Compounds*. New York: Academic Press, Inc.
- Allen, C. W. (1991). *Chem. Rev.* **91**, 119–135.
- Bakos, D., Kosik, M., Antos, K., Karolyova, M. & Vyskocil, I. (1982). *Fire Mater.* **6**, 10–11.
- Barclay, T. M., Hicks, R. G., Ichimura, A. S. & Patenaude, G. W. (2002). *Can. J. Chem.* **80**, 1501–1506.
- Bruker (2011). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Carriedo, G. A., Fernandez-Catuxo, L., Alonso, F. J. G., Gomez-Elipe, P. & Gonzalez, P. A. (1996). *Macromolecules*, **29**, 5320–5325.
- Chandrasekhar, V., Pandey, M. D., Das, B., Mahanti, B., Gopal, K. & Azhakar, R. (2011). *Tetrahedron*, **67**, 6917–6926.
- Chandrasekhar, V., Senapati, T., Dey, A., Das, S., Kalisz, M. & Clérac, R. (2012). *Inorg. Chem.* **51**, 2031–2038.
- Chandrasekhar, V., Thilagar, P., Krishnan, V., Bickley, J. F. & Steiner, A. (2007). *Cryst. Growth Des.* **7**, 668–675.
- Ciftci, G. Y., Ecik, E. T., Yildirim, T., Bilgin, K., Senkuytu, E., Yuksel, F., Uludag, Y. & Kilic, A. (2013). *Tetrahedron*, **69**, 1454–1461.
- Drewno, M. J. & Barker, R. H. (1985). *Cellulose Chemistry and its applications*, edited by T. P. Nevell & S. H. Zeronian, pp. 423–454. John Wiley & Sons, Inc.
- Duax, W. L., Weeks, C. M. & Rohrer, D. C. (1976). *Topics in Stereochemistry*, Vol. 9, edited by E. L. Eliel & N. Allinger, New York: New York: John Wiley, pp. 271–383.
- Etter, M. C. (1990). *Acc. Chem. Res.* **23**, 120–126.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Harmjanz, M., Piglosiewicz, I. M., Scott, B. L. & Burns, C. J. (2004). *Inorg. Chem.* **43**, 642–650.
- Olthof, R. (1969). *Acta Cryst.* **B25**, 2040–2045.
- Sheldrick, G. (2004). *TWINABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2013). E69, o1491–o1492 [doi:10.1107/S1600536813023830]

1,1',4,5-Tetrahydrotrispiro[1,3,2-diazaphosphole-2,2'-[1,3,5,2,4,6]triazatriphosphinine-4',6''-dibenzo[d,f][1,3,2]dioxaphosphepine-6',6'''-dibenzo[d,f][1,3,2]dioxaphosphepine] acetone monosolvate

Krystal R. Fontenot, Michael W. Easson, Frank R. Fronczek and Brian D. Condon

S1. Comment

The non-halogenated title compound contains two key elements phosphorus and nitrogen, which are incorporated into the backbone moiety of the molecule to achieve flame retardant properties (Bakos *et al.*, 1982). Studies have shown that an increase in the number of phosphorus (P), nitrogen (N), or P—N bonds leads to improved flame retardancy (Drews & Barker, 1985). The substitution of two biphenol groups and ethylenediamine onto phosphazene was achieved through nucleophilic substitution in high yield. This compound was applied to cotton fabric and has shown promising preliminary results as a potential flame retardant.

The proposed efficacy and novelty of this flame retardant is based on the design of phosphorus (P3) surrounded by four nitrogen (N2, N3, N4, N5) atoms, thereby increasing the number of P—N bonds in the molecule. Incorporation of the 2,2'-biphenol moiety at P1 and P2 promotes a seven membered spirocyclic structure with endocyclic torsion angles about the central biphenol C—C bonds being nearly equal, $-44.4(3)^\circ$ for C1/C6/C7/C12 and $-41.5(3)^\circ$ for C13/C18/C19/C24. Both seven-membered rings have twist conformations with the P atom on the local twofold axis; however the deviation from C_2 local symmetry is substantial for both. The asymmetry parameter (Duax *et al.*, 1976) is $14.83(13)^\circ$ for the ring containing P1 and $18.44(14)^\circ$ for that containing P2.

The utilization of ethylenediamine at the P3 position forms a five-membered ring with new P—N bonds, which lies roughly perpendicular (dihedral angle $81.42(4)^\circ$) to the phosphazene ring. The conformation of the 5-membered ring is closest to an envelope with C25 at the flap position. C25 lies $0.554(2) \text{ \AA}$ out of the plane of the other four atoms, and the C_s asymmetry parameter is $3.44(17)^\circ$. The N—P—N bonds have angles of $94.47(9)^\circ$ to $113.64(9)^\circ$, which is lower than typical angles of $\sim 118^\circ$ (Allcock, 1972). The narrowing of the angle may be due to van der Waals repulsions from the four N atoms surrounding P3 (Allcock, 1972). It has been reported that N—P—N bonds with two bulky phenyl groups on P have a narrowed angle of 115° due to mutual repulsions from the bulky phenyl group (Allcock, 1972). The phosphazene ring itself is slightly nonplanar, having a boat distortion, with P2 lying $0.1982(4) \text{ \AA}$ and N3 lying $0.161(2) \text{ \AA}$ on the same side of the plane defined by P1, P3, N1, and N2. The phosphazene ring has P—N bond lengths ranging from $1.5665(17)$ to $1.6171(17) \text{ \AA}$. These values are typical for phosphazene rings (Barclay *et al.* 2002; Olthof, 1969). This molecule has similar bond angles and bond lengths, when substituted with the same or comparable spiro molecules, as found in the literature (Allcock, 1972, Ciftci *et al.*, 2013). Overall, the molecule has approximate C_2 symmetry, as seen in Fig. 1.

Both NH groups donate intermolecular hydrogen bonds, as shown in Fig. 2. The N5—H \cdots N3 (at $2-x, 1-y, 1-z$) bond leads to centrosymmetric hydrogen-bonded dimers, with graph set $R^2_2(8)$ (Etter, 1990), while N4 donates a hydrogen bond to acetone O. The acetone molecule forms two C—H \cdots O hydrogen bonds with the phosphazene molecule, one as a

donor and the other as an acceptor (Table 1).

S2. Experimental

All reagent grade chemicals were purchased from Sigma Aldrich and were used without further purification. ESI Mass Spectra were recorded on an Agilent Technologies 6520 Accurate Mass Q-TOF LC/MS. NMR spectra were recorded using a Varian 400MHZ spectrometer. ^1H and ^{13}C are given in δ relative to TMS and ^{31}P is given δ relative to external 85% aqueous H_3PO_4 .

2,2-Dichloro-4,4,6,6-bis[spiro(2,2'-dioxy-1''-biphenyl)] Cyclotriphosphazene (1) was synthesized as previously reported in the literature (Allen, 1991; Carriedo *et al.* 1996).

A 3-neck round bottom flask was equipped with an argon inlet, an addition funnel, and a stopper. (1) was allowed to stir in CH_2Cl_2 until completely dissolved. A solution of ethylenediamine in CH_2Cl_2 was added drop-wise to the round bottom flask at 0°C and allowed to warm to room temperature. The reaction was allowed to stir for 24 h. Thin Layer Chromatography (TLC) was used to follow the reaction using Hexanes/ CH_2Cl_2 (50:50) and DCM/MeOH (90:10). Purification was achieved by gently extracting the organic layer with warm distilled water. The organic layer was dried by anhydrous sodium sulfate and concentrated. The product (2) obtained was a white solid with a yield of 84%. The compound was tested by ESI-MS and ^1H , ^{13}C , ^{31}P -NMR using Acetone- d_6 . Crystals of (2) were grown in an NMR tube from acetone.

S3. Refinement

H atoms on C were placed in idealized positions with C—H distances 0.95 - 0.99 Å and thereafter treated as riding, with a torsional parameter refined for each methyl group. Coordinates of the NH hydrogen atoms were refined isotropically. U_{iso} for H were assigned as 1.2 times U_{eq} of the attached atoms (1.5 for methyl).





Figure 1

The molecular structure of the title compound with atomic numbering scheme and 50% ellipsoids. The solvent molecule is not shown.

**Figure 2**

Hydrogen bonding, illustrating the centrosymmetric dimer about 1, 1/2, 1/2 and the two acetone acceptors. H atoms on C are not shown.

1,1',4,5-Tetrahydrotrispiro[1,3,2-diazaphosphole-2,2'-[1,3,5,2,4,6]triazatriphosphinine-4',6''-dibenzo[d,f][1,3,2]dioxaphosphepine-6',6'''-dibenzo[d,f][1,3,2]dioxaphosphepine] acetone monosolvate

Crystal data

$C_{26}H_{22}N_5O_4P_3 \cdot C_3H_6O$

$M_r = 619.47$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2ybc$

$a = 9.4901\ (9)\ \text{\AA}$

$b = 22.9466\ (19)\ \text{\AA}$

$c = 13.1776\ (13)\ \text{\AA}$

$\beta = 97.978\ (6)^\circ$

$V = 2841.9\ (5)\ \text{\AA}^3$

$Z = 4$

$F(000) = 1288$

$D_x = 1.448\ \text{Mg m}^{-3}$

Cu $K\alpha$ radiation, $\lambda = 1.54184\ \text{\AA}$

Cell parameters from 3314 reflections

$\theta = 3.9\text{--}67.8^\circ$

$\mu = 2.34\ \text{mm}^{-1}$

$T = 100\ \text{K}$

Plate, colorless

$0.19 \times 0.12 \times 0.03\ \text{mm}$

Data collection

Bruker Kappa APEXII DUO CCD
diffractometer
Radiation source: $I\mu S$ microfocus
QUAZAR multilayer optics monochromator
 φ and ω scans
Absorption correction: multi-scan
(TWINABS; Sheldrick, 2004)
 $T_{\min} = 0.664$, $T_{\max} = 0.933$

33208 measured reflections
5065 independent reflections
4276 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.051$
 $\theta_{\max} = 68.7^\circ$, $\theta_{\min} = 3.9^\circ$
 $h = -11 \rightarrow 11$
 $k = 0 \rightarrow 27$
 $l = 0 \rightarrow 15$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.102$
 $S = 1.08$
5065 reflections
387 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0595P)^2 + 0.7869P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.40 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.39 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. The sample was a multiple crystal ("twin") with two components. The second domain was rotated from first domain by 2.6 degrees about reciprocal axis -0.141 1.000 0.976 and real axis -0.085 0.341 1.000
The twin law to convert hkl from first to this domain (*SHELXL* TWIN matrix): 1.002 - 0.016 0.017, 0.096 0.999 0.015, -0.032 - 0.001 0.996

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections, using a TWIN4 hkl file prepared by TWINABS (Sheldrick, 2004).

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
P1	0.86867 (5)	0.57153 (2)	0.27605 (4)	0.01501 (13)
P2	0.70274 (5)	0.47853 (2)	0.30698 (4)	0.01500 (13)
P3	0.79507 (5)	0.55177 (2)	0.47261 (4)	0.01498 (13)
O1	1.02561 (15)	0.56625 (6)	0.24629 (11)	0.0169 (3)
O2	0.80895 (14)	0.62781 (6)	0.21219 (11)	0.0171 (3)
O3	0.77860 (14)	0.41639 (6)	0.30107 (11)	0.0165 (3)
O4	0.54174 (14)	0.46339 (6)	0.26346 (11)	0.0174 (3)
N1	0.77084 (18)	0.51945 (7)	0.22950 (13)	0.0173 (4)
N2	0.69897 (18)	0.49976 (7)	0.41955 (13)	0.0178 (4)
N3	0.89141 (18)	0.58162 (7)	0.39485 (13)	0.0172 (4)
N4	0.70206 (19)	0.59910 (8)	0.52986 (14)	0.0190 (4)
H4N	0.659 (3)	0.6247 (11)	0.4945 (19)	0.023*
N5	0.89724 (19)	0.53234 (8)	0.57900 (13)	0.0182 (4)
H5N	0.921 (3)	0.4976 (11)	0.5831 (19)	0.022*

C1	1.0620 (2)	0.58150 (9)	0.14985 (16)	0.0175 (4)
C2	1.1120 (2)	0.53795 (9)	0.09149 (16)	0.0195 (4)
H2	1.1120	0.4985	0.1134	0.023*
C3	1.1619 (2)	0.55245 (9)	0.00081 (17)	0.0231 (5)
H3	1.1962	0.5228	-0.0399	0.028*
C4	1.1620 (2)	0.61020 (10)	-0.03067 (16)	0.0233 (5)
H4	1.1973	0.6201	-0.0925	0.028*
C5	1.1105 (2)	0.65351 (9)	0.02810 (16)	0.0205 (4)
H5	1.1096	0.6928	0.0054	0.025*
C6	1.0600 (2)	0.64004 (9)	0.12034 (15)	0.0170 (4)
C7	1.0101 (2)	0.68643 (8)	0.18533 (15)	0.0174 (4)
C8	1.0836 (2)	0.73950 (9)	0.20275 (16)	0.0203 (4)
H8	1.1679	0.7460	0.1730	0.024*
C9	1.0346 (2)	0.78235 (9)	0.26264 (16)	0.0225 (5)
H9	1.0854	0.8180	0.2735	0.027*
C10	0.9119 (2)	0.77390 (9)	0.30718 (16)	0.0218 (4)
H10	0.8794	0.8035	0.3486	0.026*
C11	0.8369 (2)	0.72215 (9)	0.29101 (16)	0.0205 (4)
H11	0.7523	0.7159	0.3206	0.025*
C12	0.8877 (2)	0.67975 (8)	0.23097 (16)	0.0175 (4)
C13	0.7274 (2)	0.37086 (8)	0.35739 (16)	0.0170 (4)
C14	0.8068 (2)	0.35465 (9)	0.44930 (16)	0.0207 (4)
H14	0.8913	0.3751	0.4752	0.025*
C15	0.7602 (3)	0.30792 (9)	0.50272 (17)	0.0265 (5)
H15	0.8124	0.2964	0.5663	0.032*
C16	0.6377 (2)	0.27796 (9)	0.46347 (18)	0.0265 (5)
H16	0.6068	0.2458	0.5000	0.032*
C17	0.5602 (2)	0.29493 (9)	0.37092 (18)	0.0234 (5)
H17	0.4765	0.2741	0.3446	0.028*
C18	0.6038 (2)	0.34235 (8)	0.31581 (16)	0.0182 (4)
C19	0.5226 (2)	0.36114 (9)	0.21647 (16)	0.0182 (4)
C20	0.4629 (2)	0.32016 (9)	0.14425 (18)	0.0246 (5)
H20	0.4762	0.2798	0.1586	0.029*
C21	0.3846 (2)	0.33719 (11)	0.05215 (19)	0.0296 (5)
H21	0.3440	0.3086	0.0047	0.036*
C22	0.3657 (2)	0.39597 (11)	0.02949 (18)	0.0291 (5)
H22	0.3130	0.4077	-0.0338	0.035*
C23	0.4239 (2)	0.43759 (10)	0.09950 (17)	0.0237 (5)
H23	0.4115	0.4779	0.0845	0.028*
C24	0.5001 (2)	0.41980 (9)	0.19118 (16)	0.0178 (4)
C25	0.7808 (2)	0.61684 (9)	0.62821 (16)	0.0233 (5)
H25A	0.7156	0.6317	0.6748	0.028*
H25B	0.8514	0.6475	0.6188	0.028*
C26	0.8542 (2)	0.56094 (9)	0.67000 (16)	0.0213 (4)
H26A	0.9382	0.5698	0.7211	0.026*
H26B	0.7880	0.5360	0.7024	0.026*
O5	0.53486 (16)	0.66469 (7)	0.35745 (12)	0.0258 (3)
C27	0.4623 (2)	0.62727 (9)	0.30959 (17)	0.0220 (5)

C28	0.3829 (2)	0.58239 (9)	0.36180 (18)	0.0255 (5)
H28A	0.4119	0.5846	0.4360	0.038*
H28B	0.4045	0.5435	0.3373	0.038*
H28C	0.2805	0.5897	0.3462	0.038*
C29	0.4499 (3)	0.62433 (11)	0.19530 (18)	0.0307 (5)
H29A	0.4922	0.6594	0.1695	0.046*
H29B	0.3493	0.6219	0.1661	0.046*
H29C	0.5002	0.5898	0.1754	0.046*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
P1	0.0158 (3)	0.0150 (2)	0.0145 (3)	-0.00015 (19)	0.00280 (19)	0.00056 (18)
P2	0.0159 (3)	0.0140 (2)	0.0153 (3)	-0.00058 (19)	0.00309 (19)	-0.00092 (18)
P3	0.0156 (3)	0.0156 (2)	0.0139 (3)	0.00053 (19)	0.00267 (19)	-0.00072 (18)
O1	0.0168 (7)	0.0184 (7)	0.0161 (7)	0.0007 (5)	0.0041 (6)	0.0038 (5)
O2	0.0156 (7)	0.0159 (7)	0.0194 (7)	-0.0013 (5)	0.0007 (6)	0.0012 (5)
O3	0.0165 (7)	0.0152 (6)	0.0183 (7)	-0.0007 (5)	0.0041 (5)	-0.0009 (5)
O4	0.0157 (7)	0.0159 (6)	0.0206 (8)	0.0000 (5)	0.0024 (6)	-0.0040 (6)
N1	0.0196 (9)	0.0177 (8)	0.0151 (9)	-0.0020 (7)	0.0042 (7)	-0.0015 (6)
N2	0.0195 (9)	0.0173 (8)	0.0173 (9)	-0.0020 (7)	0.0051 (7)	0.0005 (7)
N3	0.0181 (9)	0.0174 (8)	0.0158 (9)	-0.0023 (7)	0.0016 (7)	-0.0005 (7)
N4	0.0203 (9)	0.0193 (8)	0.0176 (9)	0.0041 (7)	0.0030 (7)	-0.0017 (7)
N5	0.0204 (9)	0.0187 (8)	0.0154 (9)	0.0028 (7)	0.0019 (7)	0.0007 (7)
C1	0.0141 (10)	0.0234 (10)	0.0151 (10)	-0.0017 (8)	0.0029 (8)	0.0003 (8)
C2	0.0189 (10)	0.0177 (9)	0.0215 (11)	-0.0003 (8)	0.0013 (8)	-0.0007 (8)
C3	0.0216 (11)	0.0263 (11)	0.0212 (11)	0.0020 (9)	0.0025 (9)	-0.0039 (9)
C4	0.0239 (11)	0.0307 (11)	0.0160 (11)	-0.0022 (9)	0.0051 (9)	0.0007 (9)
C5	0.0215 (11)	0.0203 (10)	0.0198 (11)	-0.0004 (8)	0.0031 (8)	0.0031 (8)
C6	0.0151 (10)	0.0193 (10)	0.0167 (10)	-0.0017 (8)	0.0025 (8)	-0.0015 (8)
C7	0.0191 (10)	0.0181 (9)	0.0148 (10)	0.0008 (8)	0.0016 (8)	0.0033 (8)
C8	0.0225 (11)	0.0207 (10)	0.0184 (11)	-0.0024 (8)	0.0054 (8)	0.0034 (8)
C9	0.0307 (12)	0.0159 (9)	0.0209 (11)	-0.0046 (9)	0.0031 (9)	0.0009 (8)
C10	0.0286 (11)	0.0171 (10)	0.0201 (11)	0.0019 (9)	0.0051 (9)	-0.0019 (8)
C11	0.0199 (10)	0.0204 (10)	0.0212 (11)	0.0029 (8)	0.0036 (8)	0.0038 (8)
C12	0.0198 (10)	0.0132 (9)	0.0186 (10)	-0.0013 (8)	-0.0004 (8)	0.0023 (7)
C13	0.0207 (10)	0.0139 (9)	0.0182 (10)	0.0005 (8)	0.0091 (8)	-0.0011 (8)
C14	0.0228 (11)	0.0184 (9)	0.0213 (11)	0.0052 (8)	0.0044 (9)	-0.0030 (8)
C15	0.0379 (13)	0.0220 (10)	0.0208 (11)	0.0116 (10)	0.0087 (10)	0.0028 (9)
C16	0.0331 (13)	0.0194 (10)	0.0308 (13)	0.0044 (9)	0.0181 (10)	0.0044 (9)
C17	0.0236 (11)	0.0172 (10)	0.0316 (13)	0.0004 (8)	0.0113 (9)	-0.0014 (9)
C18	0.0185 (10)	0.0152 (9)	0.0227 (11)	0.0021 (8)	0.0092 (8)	-0.0027 (8)
C19	0.0148 (10)	0.0212 (10)	0.0202 (11)	-0.0021 (8)	0.0073 (8)	-0.0031 (8)
C20	0.0216 (11)	0.0228 (11)	0.0307 (13)	-0.0033 (9)	0.0086 (9)	-0.0087 (9)
C21	0.0229 (11)	0.0378 (13)	0.0276 (13)	-0.0032 (10)	0.0015 (9)	-0.0167 (10)
C22	0.0231 (12)	0.0412 (13)	0.0218 (12)	0.0024 (10)	-0.0015 (9)	-0.0063 (10)
C23	0.0209 (11)	0.0257 (11)	0.0241 (12)	0.0025 (9)	0.0019 (9)	-0.0006 (9)
C24	0.0138 (10)	0.0202 (10)	0.0196 (11)	-0.0008 (8)	0.0038 (8)	-0.0046 (8)

C25	0.0234 (11)	0.0275 (11)	0.0188 (11)	0.0003 (9)	0.0023 (9)	-0.0057 (9)
C26	0.0213 (11)	0.0268 (10)	0.0159 (11)	-0.0005 (9)	0.0029 (8)	-0.0013 (8)
O5	0.0236 (8)	0.0247 (8)	0.0291 (9)	-0.0022 (6)	0.0034 (7)	-0.0016 (6)
C27	0.0162 (10)	0.0234 (10)	0.0268 (12)	0.0065 (9)	0.0039 (9)	-0.0008 (9)
C28	0.0210 (11)	0.0245 (11)	0.0319 (13)	0.0012 (9)	0.0069 (9)	-0.0014 (9)
C29	0.0257 (12)	0.0403 (13)	0.0255 (13)	-0.0010 (10)	0.0017 (10)	0.0027 (10)

Geometric parameters (Å, °)

P1—N3	1.5675 (17)	C10—H10	0.9500
P1—N1	1.5837 (17)	C11—C12	1.382 (3)
P1—O1	1.5965 (14)	C11—H11	0.9500
P1—O2	1.6016 (14)	C13—C14	1.385 (3)
P2—N2	1.5665 (17)	C13—C18	1.388 (3)
P2—N1	1.5890 (17)	C14—C15	1.388 (3)
P2—O4	1.5940 (14)	C14—H14	0.9500
P2—O3	1.6041 (14)	C15—C16	1.387 (3)
P3—N2	1.6024 (17)	C15—H15	0.9500
P3—N3	1.6171 (17)	C16—C17	1.390 (3)
P3—N4	1.6471 (17)	C16—H16	0.9500
P3—N5	1.6514 (18)	C17—C18	1.402 (3)
O1—C1	1.407 (2)	C17—H17	0.9500
O2—C12	1.410 (2)	C18—C19	1.488 (3)
O3—C13	1.407 (2)	C19—C24	1.396 (3)
O4—C24	1.400 (2)	C19—C20	1.401 (3)
N4—C25	1.461 (3)	C20—C21	1.388 (3)
N4—H4N	0.82 (3)	C20—H20	0.9500
N5—C26	1.473 (3)	C21—C22	1.388 (4)
N5—H5N	0.83 (3)	C21—H21	0.9500
C1—C2	1.384 (3)	C22—C23	1.388 (3)
C1—C6	1.398 (3)	C22—H22	0.9500
C2—C3	1.386 (3)	C23—C24	1.381 (3)
C2—H2	0.9500	C23—H23	0.9500
C3—C4	1.389 (3)	C25—C26	1.526 (3)
C3—H3	0.9500	C25—H25A	0.9900
C4—C5	1.390 (3)	C25—H25B	0.9900
C4—H4	0.9500	C26—H26A	0.9900
C5—C6	1.402 (3)	C26—H26B	0.9900
C5—H5	0.9500	O5—C27	1.220 (3)
C6—C7	1.484 (3)	C27—C29	1.496 (3)
C7—C12	1.389 (3)	C27—C28	1.498 (3)
C7—C8	1.406 (3)	C28—H28A	0.9800
C8—C9	1.381 (3)	C28—H28B	0.9800
C8—H8	0.9500	C28—H28C	0.9800
C9—C10	1.388 (3)	C29—H29A	0.9800
C9—H9	0.9500	C29—H29B	0.9800
C10—C11	1.386 (3)	C29—H29C	0.9800

N3—P1—N1	119.36 (9)	C11—C12—O2	118.47 (18)
N3—P1—O1	104.64 (8)	C7—C12—O2	118.17 (17)
N1—P1—O1	111.37 (8)	C14—C13—C18	123.09 (19)
N3—P1—O2	113.28 (8)	C14—C13—O3	118.37 (18)
N1—P1—O2	105.03 (8)	C18—C13—O3	118.46 (18)
O1—P1—O2	101.90 (7)	C13—C14—C15	118.6 (2)
N2—P2—N1	119.30 (9)	C13—C14—H14	120.7
N2—P2—O4	105.18 (8)	C15—C14—H14	120.7
N1—P2—O4	110.65 (8)	C16—C15—C14	120.2 (2)
N2—P2—O3	113.07 (9)	C16—C15—H15	119.9
N1—P2—O3	105.59 (8)	C14—C15—H15	119.9
O4—P2—O3	101.71 (7)	C15—C16—C17	120.2 (2)
N2—P3—N3	112.12 (9)	C15—C16—H16	119.9
N2—P3—N4	112.47 (9)	C17—C16—H16	119.9
N3—P3—N4	113.39 (9)	C16—C17—C18	120.9 (2)
N2—P3—N5	113.64 (9)	C16—C17—H17	119.6
N3—P3—N5	109.59 (9)	C18—C17—H17	119.6
N4—P3—N5	94.47 (9)	C13—C18—C17	117.1 (2)
C1—O1—P1	123.88 (13)	C13—C18—C19	121.14 (18)
C12—O2—P1	116.68 (12)	C17—C18—C19	121.76 (19)
C13—O3—P2	116.39 (12)	C24—C19—C20	116.8 (2)
C24—O4—P2	124.52 (12)	C24—C19—C18	122.21 (18)
P1—N1—P2	117.77 (11)	C20—C19—C18	120.98 (19)
P2—N2—P3	123.95 (11)	C21—C20—C19	121.5 (2)
P1—N3—P3	123.96 (11)	C21—C20—H20	119.3
C25—N4—P3	110.45 (14)	C19—C20—H20	119.3
C25—N4—H4N	117.0 (17)	C22—C21—C20	119.9 (2)
P3—N4—H4N	117.8 (18)	C22—C21—H21	120.0
C26—N5—P3	111.99 (14)	C20—C21—H21	120.0
C26—N5—H5N	118.6 (17)	C21—C22—C23	119.9 (2)
P3—N5—H5N	115.9 (17)	C21—C22—H22	120.1
C2—C1—C6	122.12 (19)	C23—C22—H22	120.1
C2—C1—O1	117.93 (18)	C24—C23—C22	119.3 (2)
C6—C1—O1	119.65 (18)	C24—C23—H23	120.3
C1—C2—C3	119.36 (19)	C22—C23—H23	120.3
C1—C2—H2	120.3	C23—C24—C19	122.56 (19)
C3—C2—H2	120.3	C23—C24—O4	116.64 (18)
C2—C3—C4	120.1 (2)	C19—C24—O4	120.46 (18)
C2—C3—H3	120.0	N4—C25—C26	103.74 (17)
C4—C3—H3	120.0	N4—C25—H25A	111.0
C3—C4—C5	120.1 (2)	C26—C25—H25A	111.0
C3—C4—H4	120.0	N4—C25—H25B	111.0
C5—C4—H4	120.0	C26—C25—H25B	111.0
C4—C5—C6	120.97 (19)	H25A—C25—H25B	109.0
C4—C5—H5	119.5	N5—C26—C25	104.18 (17)
C6—C5—H5	119.5	N5—C26—H26A	110.9
C1—C6—C5	117.38 (19)	C25—C26—H26A	110.9
C1—C6—C7	121.47 (18)	N5—C26—H26B	110.9

C5—C6—C7	121.13 (18)	C25—C26—H26B	110.9
C12—C7—C8	116.76 (18)	H26A—C26—H26B	108.9
C12—C7—C6	121.56 (18)	O5—C27—C29	120.8 (2)
C8—C7—C6	121.68 (18)	O5—C27—C28	122.0 (2)
C9—C8—C7	120.7 (2)	C29—C27—C28	117.21 (19)
C9—C8—H8	119.6	C27—C28—H28A	109.5
C7—C8—H8	119.6	C27—C28—H28B	109.5
C8—C9—C10	120.73 (19)	H28A—C28—H28B	109.5
C8—C9—H9	119.6	C27—C28—H28C	109.5
C10—C9—H9	119.6	H28A—C28—H28C	109.5
C11—C10—C9	119.85 (19)	H28B—C28—H28C	109.5
C11—C10—H10	120.1	C27—C29—H29A	109.5
C9—C10—H10	120.1	C27—C29—H29B	109.5
C12—C11—C10	118.6 (2)	H29A—C29—H29B	109.5
C12—C11—H11	120.7	C27—C29—H29C	109.5
C10—C11—H11	120.7	H29A—C29—H29C	109.5
C11—C12—C7	123.33 (18)	H29B—C29—H29C	109.5
N3—P1—O1—C1	-151.76 (15)	C5—C6—C7—C12	137.4 (2)
N1—P1—O1—C1	77.97 (16)	C1—C6—C7—C8	136.4 (2)
O2—P1—O1—C1	-33.56 (16)	C5—C6—C7—C8	-41.9 (3)
N3—P1—O2—C12	54.75 (16)	C12—C7—C8—C9	0.0 (3)
N1—P1—O2—C12	-173.32 (14)	C6—C7—C8—C9	179.33 (19)
O1—P1—O2—C12	-57.08 (15)	C7—C8—C9—C10	0.2 (3)
N2—P2—O3—C13	52.84 (16)	C8—C9—C10—C11	-0.4 (3)
N1—P2—O3—C13	-175.00 (13)	C9—C10—C11—C12	0.5 (3)
O4—P2—O3—C13	-59.42 (15)	C10—C11—C12—C7	-0.3 (3)
N2—P2—O4—C24	-148.34 (15)	C10—C11—C12—O2	-178.18 (18)
N1—P2—O4—C24	81.56 (17)	C8—C7—C12—C11	0.1 (3)
O3—P2—O4—C24	-30.24 (17)	C6—C7—C12—C11	-179.23 (19)
N3—P1—N1—P2	-1.48 (16)	C8—C7—C12—O2	177.90 (17)
O1—P1—N1—P2	120.63 (11)	C6—C7—C12—O2	-1.4 (3)
O2—P1—N1—P2	-129.84 (11)	P1—O2—C12—C11	-103.17 (18)
N2—P2—N1—P1	16.39 (16)	P1—O2—C12—C7	78.9 (2)
O4—P2—N1—P1	138.55 (10)	P2—O3—C13—C14	-103.07 (18)
O3—P2—N1—P1	-112.16 (11)	P2—O3—C13—C18	80.20 (19)
N1—P2—N2—P3	-17.32 (17)	C18—C13—C14—C15	-0.4 (3)
O4—P2—N2—P3	-142.15 (12)	O3—C13—C14—C15	-176.97 (17)
O3—P2—N2—P3	107.71 (12)	C13—C14—C15—C16	0.8 (3)
N3—P3—N2—P2	2.48 (16)	C14—C15—C16—C17	-0.6 (3)
N4—P3—N2—P2	131.67 (12)	C15—C16—C17—C18	-0.1 (3)
N5—P3—N2—P2	-122.48 (13)	C14—C13—C18—C17	-0.3 (3)
N1—P1—N3—P3	-14.21 (17)	O3—C13—C18—C17	176.31 (17)
O1—P1—N3—P3	-139.60 (11)	C14—C13—C18—C19	-179.62 (18)
O2—P1—N3—P3	110.26 (12)	O3—C13—C18—C19	-3.0 (3)
N2—P3—N3—P1	13.44 (16)	C16—C17—C18—C13	0.5 (3)
N4—P3—N3—P1	-115.26 (13)	C16—C17—C18—C19	179.83 (19)
N5—P3—N3—P1	140.61 (12)	C13—C18—C19—C24	-41.5 (3)

N2—P3—N4—C25	138.76 (14)	C17—C18—C19—C24	139.2 (2)
N3—P3—N4—C25	-92.72 (16)	C13—C18—C19—C20	139.6 (2)
N5—P3—N4—C25	20.88 (16)	C17—C18—C19—C20	-39.7 (3)
N2—P3—N5—C26	-112.97 (15)	C24—C19—C20—C21	-0.2 (3)
N3—P3—N5—C26	120.73 (15)	C18—C19—C20—C21	178.7 (2)
N4—P3—N5—C26	3.95 (16)	C19—C20—C21—C22	0.8 (3)
P1—O1—C1—C2	-117.03 (18)	C20—C21—C22—C23	-0.7 (4)
P1—O1—C1—C6	69.1 (2)	C21—C22—C23—C24	-0.1 (3)
C6—C1—C2—C3	0.1 (3)	C22—C23—C24—C19	0.7 (3)
O1—C1—C2—C3	-173.67 (18)	C22—C23—C24—O4	-172.71 (19)
C1—C2—C3—C4	0.2 (3)	C20—C19—C24—C23	-0.5 (3)
C2—C3—C4—C5	-0.7 (3)	C18—C19—C24—C23	-179.51 (19)
C3—C4—C5—C6	1.0 (3)	C20—C19—C24—O4	172.62 (18)
C2—C1—C6—C5	0.2 (3)	C18—C19—C24—O4	-6.3 (3)
O1—C1—C6—C5	173.83 (17)	P2—O4—C24—C23	-118.86 (18)
C2—C1—C6—C7	-178.11 (19)	P2—O4—C24—C19	67.6 (2)
O1—C1—C6—C7	-4.5 (3)	P3—N4—C25—C26	-37.8 (2)
C4—C5—C6—C1	-0.7 (3)	P3—N5—C26—C25	-25.7 (2)
C4—C5—C6—C7	177.58 (19)	N4—C25—C26—N5	38.3 (2)
C1—C6—C7—C12	-44.4 (3)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N4—H4N...O5	0.82 (3)	2.22 (3)	2.991 (2)	158 (2)
N5—H5N...N3 ⁱ	0.83 (3)	2.53 (3)	3.285 (2)	151 (2)

Symmetry code: (i) $-x+2, -y+1, -z+1$.