

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

**(E)-13-(2-Bromophenyl)micheliolide**

Narsimha Reddy Penthala,<sup>a</sup> Shobanbabu Bommagani,<sup>a</sup>  
Venumadhav Janganati,<sup>a</sup> Sean Parkin<sup>b</sup> and Peter A.  
Crooks<sup>a\*</sup>

<sup>a</sup>Department of Pharmaceutical Sciences, College of Pharmacy, University of Arkansas for Medical Sciences, Little Rock, AR 72205, USA, and <sup>b</sup>Department of Chemistry, University of Kentucky, Lexington, KY 40506, USA  
Correspondence e-mail: pacrooks@uams.edu


Received 20 November 2013; accepted 29 January 2014

Key indicators: single-crystal X-ray study;  $T = 90$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.025;  $wR$  factor = 0.064; data-to-parameter ratio = 18.0.

The title compound,  $\text{C}_{21}\text{H}_{23}\text{BrO}_3$  [systematic name: (3*E*,3*aS*,6*Z*,9*R*,9*aS*,9*bS*)-3-(2-bromobenzylidene)-9-hydroxy-6,9-dimethyl-3,3*a*,4,5,7,8,9,9*a*-octahydroazuleno[4,5-*b*]furan-2(9*bH*)-one] was prepared by the reaction of 1-bromo-2-iodobenzene with micheliolide [systematic name: (3*aS*,*R*,9*aS*,9*bS*,*Z*)-9-hydroxy-6,9-dimethyl-3-methylene-3,3*a*,4,5,7,8,9,9*a*-octahydroazuleno[4,5-*b*]furan-2(9*bH*)-one] under Heck reaction conditions. The title compound exhibits intramolecular O—H...O hydrogen bonding between the hydroxy group and the lactone ring O atom, forming a ring of graph-set motif *S*(6). The 2-bromophenyl group is *trans* to the lactone ring, indicating that this is the *E* isomer (geometry of the exocyclic C=C bond). The dihedral angle between the benzene ring of the 2-bromophenyl moiety and the mean plane of the lactone ring is 51.68 (7)°.

**Related literature**

For the biological activity of micheliolide Michael addition compounds, see: Rodriguez *et al.* (1976); Sethi *et al.* (1984); Neelakantan *et al.* (2009); Zhang *et al.* (2012). For details of the Heck chemistry, see: Han *et al.* (2009). For the crystal structure of micheliolide, see: Acosta *et al.* (1991). For the crystal structure of a similar compound, see: Penthala *et al.* (2013).

**Experimental***Crystal data*

$\text{C}_{21}\text{H}_{23}\text{BrO}_3$   
 $M_r = 403.30$   
Orthorhombic,  $P2_12_12_1$   
 $a = 7.1617$  (1) Å  
 $b = 13.1615$  (2) Å  
 $c = 19.2306$  (3) Å

$V = 1812.65$  (5) Å<sup>3</sup>  
 $Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 2.29$  mm<sup>-1</sup>  
 $T = 90$  K  
0.20 × 0.20 × 0.15 mm

*Data collection*

Nonius KappaCCD diffractometer  
Absorption correction: multi-scan  
(*SADABS*; Sheldrick, 2008*a*)  
 $T_{\min} = 0.550$ ,  $T_{\max} = 0.714$

48139 measured reflections  
4148 independent reflections  
3871 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.047$

*Refinement*

$R[F^2 > 2\sigma(F^2)] = 0.025$   
 $wR(F^2) = 0.064$   
 $S = 1.04$   
4148 reflections  
231 parameters  
H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\text{max}} = 0.41$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.34$  e Å<sup>-3</sup>  
Absolute structure: Flack  
parameter determined using 1569 quotients  $[(I^+) - (I^-)] / [(I^+) + (I^-)]$  (Parsons *et al.*, 2013)  
Absolute structure parameter: 0.032 (3)

**Table 1**

Hydrogen-bond geometry (Å, °).

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O3—H3...O2	0.77 (4)	2.26 (4)	2.883 (3)	139 (4)

Data collection: *COLLECT* (Nonius, 1998); cell refinement: *SCALEPACK* (Otwinowski & Minor, 2006); data reduction: *DENZO-SMN* (Otwinowski & Minor, 2006); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008*b*); program(s) used to refine structure: *SHELXL2013* (Sheldrick, 2008*b*); molecular graphics: *XP in SHELXTL* (Sheldrick, 2008*b*); software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008*b*).

This work was supported by the NIH/NCI [grant No. CA158275].

---

Supporting information for this paper is available from the IUCr electronic archives (Reference: SJ5372).

---

## References

- Acosta, J. C., Fronczek, F. R. & Fischer, N. H. (1991). *Acta Cryst.* **C47**, 2702–2704.
- Han, C., Barrios, F. J., Rioski, M. V. & Colby, D. A. (2009). *J. Org. Chem.* **74**, 7176–7179.
- Neelakantan, S., Nasim, S., Guzman, M. L., Jordan, C. T. & Crooks, P. A. (2009). *Bioorg. Med. Chem. Lett.* **19**, 4346–4349.
- Nonius (1998). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (2006). *International Tables for Crystallography*, Vol. F, ch. 11.4, pp. 226–235. Dordrecht: Kluwer Academic Publishers.
- Parsons, S., Flack, H. D. & Wagner, T. (2013). *Acta Cryst.* **B69**, 249–259.
- Penthala, N. R., Janganati, V., Parkin, S., Varughese, K. I. & Crooks, P. A. (2013). *Acta Cryst.* **E69**, o1709–o1710.
- Rodriguez, E., Towers, G. H. N. & Mitchell, J. C. (1976). *Phytochemistry*, **15**, 1573–1580.
- Sethi, V. K., Thappat, R. K., Dhar, K. L. & Atal, C. K. (1984). *Planta Med.* **50**, 364–?
- Sheldrick, G. M. (2008a). *SADABS* University of Göttingen, Germany.
- Sheldrick, G. M. (2008b). *Acta Cryst.* **A64**, 112–122.
- Zhang, Q., Lu, Y., Ding, Y., Zhai, J., Ji, Q., Ma, W., Yang, M., Fan, H., Long, J., Tong, Z., Shi, Y., Jia, Y., Han, B., Zhang, W., Qiu, C., Ma, X., Li, Q., Shi, Q., Zhang, H., Li, D., Zhang, J., Lin, J., Li, L. Y., Gao, Y. & Chen, Y. (2012). *J. Med. Chem.* **55**, 8757–8769.

## supporting information

*Acta Cryst.* (2014). E70, o251–o252 [doi:10.1107/S1600536814002177]

**(E)-13-(2-Bromophenyl)micheliolide**

**Narsimha Reddy Penthala, Shobanbabu Bommagani, Venumadhav Janganati, Sean Parkin and Peter A. Crooks**

**S1. Comment**

Micheliolide (MCL) belongs to the class of guaianolide sesquiterpene lactones, which are being developed for the treatment of cancer (Sethi *et al.*, 1984 and Zhang *et al.*, 2012). The exocyclic double bond in such sesquiterpenes is believed to be responsible for their biological properties because of its exceptional chemical reactivity with nucleophilic groups (Rodriguez *et al.*, 1976). The MCL crystal structure was described by Acosta *et al.* (1991). Recently, micheliolide Michael addition analogs were reported as potent anti-leukemic agents (Zhang *et al.*, 2012). In a recent communication we reported the crystal structure of (*E*)-13-(4-aminophenyl)parthenolide, a Heck reaction derivative of parthenolide (Penthala *et al.*, 2013). Now, in continuation of our research on sesquiterpene lactones as anti-leukemic agents (Neelakantan *et al.*, 2009), we are focusing on the synthesis of the title *E*-olefinic analogue of micheliolide, which was obtained from the reaction of micheliolide with 1-bromo-2-iodobenzene utilizing Heck chemistry (Han *et al.*, 2009). In order to obtain detailed information on the structural conformation of the title compound and to establish the geometry of the exocyclic double bond, a single-crystal X-ray structure determination has been carried out.

Recrystallization of the title compound from hexanes gave colorless needles that were suitable for X-ray analysis. The title molecule, Fig. 1, contains a central seven-membered carbocyclic ring fused to a 5-membered carbocyclic ring and a *trans* lactone ring. The two five-membered rings are in half-chair conformations, as reported in the literature for the parent compound, micheliolide (Acosta *et al.*, 1991). The X-ray studies revealed that the title compound exhibits intramolecular O—H···O hydrogen bonding to form a ring of graph-set motif *S*(6). The 2-bromophenyl group is oriented *trans* to the lactone ring to form the *E* isomer (geometry of the exocyclic C=C bond). The H atom of the hydroxy group in the molecule forms an intramolecular hydrogen bond with the lactone ring oxygen atom. The dihedral angle between the benzene ring of the 2-bromophenyl group and the mean plane of the lactone ring is 51.68 (7)°.

**S2. Experimental**

A mixture of micheliolide (prepared from parthenolide, Zhang *et al.*, 2012) (50 mg, 0.20 mmol), triethylamine (60 mg, 0.61 mmol), and 1-bromo-2-iodobenzene (63 mg, 0.22 mmol) in dimethylformamide (1 ml) was treated with palladium(II) acetate (0.5 mg, 0.002 mmol) and then stirred at 353 K for 24 h. The reaction mixture was cooled to room temperature, water (8 ml) was added, and the mixture was extracted with ethyl acetate (10 ml x 3). The separated organics were dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated under reduced pressure. The crude residue was purified using silica flash chromatography (7:3, hexanes/EtOAc) to afford the title compound, which was recrystallized from hexanes as colorless needles suitable for X-ray analysis (65 mg, 80% yield; M·P: 421–423 K); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.70–7.70 (d, *J*=2.8 Hz, 1H), 7.62–7.64 (d, *J*=8.0 Hz, 1H), 7.22–7.35 (m, 3H, Ar—H and =CH), 3.89–3.94 (t, *J*=10 Hz, 20.4 Hz, 1H), 3.01–3.06 (t, *J*=8.8 Hz, 19.6 Hz, 1H), 2.73–2.76 (d, *J*=11.2 Hz, 1H), 2.71 (s, 1H), 2.35–2.41 (m, 1H), 2.01–2.21 (m, 1H), 1.76–1.89 (m, 3H), 1.61 (s, 3H), 1.32 (s, 3H), 1.00–1.04 (m, 1H) *p.p.m.*. <sup>13</sup>C NMR (100 MHz,

$\text{CDCl}_3$ ):  $\delta$  22.57, 23.69, 24.91, 29.86, 35.36, 38.23, 48.93, 58.99, 80.79, 124.14, 126.93, 129.86, 130.47, 130.55, 131.03, 131.50, 132.86, 134.54, 136.70, 170.68 *p.p.m.*.

### S3. Refinement

H atoms were found in difference Fourier maps. Carbon-bound H atoms were subsequently placed at idealized positions with constrained distances of 0.98 Å ( $R\text{CH}_3$ ), 0.99 Å ( $R_2\text{CH}_2$ ), 1.00 Å ( $R_3\text{CH}$ ) and 0.95 Å ( $C_{\text{sp}^2}\text{H}$ ). The hydroxy hydrogen coordinates were refined.  $U_{\text{iso}}(\text{H})$  values were set to either  $1.2U_{\text{eq}}$  or  $1.5U_{\text{eq}}$  ( $R\text{CH}_3$ , OH) of the attached atom.




Figure 1

A view of the molecule with displacement ellipsoids drawn at the 50% probability level.

(3*E*,3*aS*,6*Z*,9*R*,9*aS*,9*bS*)-3-(2-Bromobenzylidene)-9-hydroxy-6,9-dimethyl-3,3*a*,4,5,7,8,9,9*a*-octahydroazuleno[4,5-*b*]furan-2(9*bH*)-one

#### Crystal data

$\text{C}_{21}\text{H}_{23}\text{BrO}_3$

$M_r = 403.30$

Orthorhombic,  $P2_12_12_1$

$a = 7.1617$  (1) Å

$b = 13.1615$  (2) Å

$c = 19.2306$  (3) Å

$V = 1812.65$  (5) Å<sup>3</sup>

$Z = 4$

$F(000) = 832$

$D_x = 1.478$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 127083 reflections

$\theta = 1.0\text{--}27.5^\circ$

$\mu = 2.29$  mm<sup>-1</sup>

$T = 90$  K

Block, cut from needle, colourless

0.20 × 0.20 × 0.15 mm

#### Data collection

Nonius KappaCCD

diffractometer

Radiation source: fine-focus sealed-tube

Detector resolution: 9.1 pixels mm<sup>-1</sup>

$\varphi$  and  $\omega$  scans at fixed  $\chi = 55^\circ$

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 2008*a*)

$T_{\text{min}} = 0.550$ ,  $T_{\text{max}} = 0.714$

48139 measured reflections

4148 independent reflections

3871 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.047$   
 $\theta_{\text{max}} = 27.5^\circ$ ,  $\theta_{\text{min}} = 1.9^\circ$   
 $h = -9 \rightarrow 9$

$k = -17 \rightarrow 17$   
 $l = -24 \rightarrow 24$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.025$   
 $wR(F^2) = 0.064$   
 $S = 1.04$   
 4148 reflections  
 231 parameters  
 0 restraints  
 Primary atom site location: structure-invariant  
 direct methods  
 Secondary atom site location: difference Fourier  
 map

Hydrogen site location: mixed  
 H atoms treated by a mixture of independent  
 and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.0352P)^2 + 0.4253P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\text{max}} = 0.001$   
 $\Delta\rho_{\text{max}} = 0.41 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.34 \text{ e } \text{\AA}^{-3}$   
 Absolute structure: Flack parameter determined  
 using 1569 quotients  $[(I^-)-(I^+)]/[(I^-)+(I^+)]$   
 (Parsons *et al.*, 2013)  
 Absolute structure parameter: 0.032 (3)

*Special details*

**Experimental.** The crystal was mounted with polyisobutene oil on the tip of a fine glass fibre, fastened in a copper mounting pin with electrical solder. It was placed directly into the cold stream of a liquid nitrogen based cryostat. Diffraction data were collected with the crystal at 90 K, which is standard practice in this laboratory for the majority of flash-cooled crystals.

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.79983 (4)	-0.11654 (2)	0.35920 (2)	0.03215 (9)
O1	0.7715 (3)	0.26157 (14)	0.45446 (10)	0.0248 (4)
O2	0.5251 (3)	0.34175 (14)	0.40803 (9)	0.0215 (4)
O3	0.3499 (3)	0.53339 (15)	0.43985 (11)	0.0285 (5)
H3	0.434 (6)	0.498 (3)	0.444 (2)	0.043*
C1	0.0338 (3)	0.3743 (2)	0.33548 (13)	0.0202 (5)
C2	-0.0575 (4)	0.4794 (2)	0.33076 (16)	0.0252 (6)
H2A	-0.0553	0.5046	0.2823	0.030*
H2B	-0.1887	0.4765	0.3468	0.030*
C3	0.0577 (4)	0.5486 (2)	0.37792 (14)	0.0243 (6)
H3A	0.0633	0.6184	0.3589	0.029*
H3B	0.0037	0.5512	0.4253	0.029*
C4	0.2513 (4)	0.50008 (19)	0.37900 (13)	0.0208 (5)
C5	0.2022 (3)	0.38520 (18)	0.38337 (12)	0.0185 (4)
H5	0.1602	0.3704	0.4319	0.022*
C6	0.3602 (3)	0.31433 (18)	0.36651 (14)	0.0192 (5)
H6	0.3917	0.3211	0.3161	0.023*
C7	0.3212 (4)	0.20208 (18)	0.38266 (13)	0.0198 (5)
H7	0.2314	0.1984	0.4224	0.024*
C8	0.2325 (4)	0.1511 (2)	0.31966 (14)	0.0237 (6)

H8A	0.2408	0.0764	0.3251	0.028*
H8B	0.3031	0.1702	0.2774	0.028*
C9	0.0275 (4)	0.1814 (2)	0.31045 (16)	0.0259 (6)
H9A	-0.0413	0.1538	0.3509	0.031*
H9B	-0.0193	0.1449	0.2690	0.031*
C10	-0.0321 (4)	0.2927 (2)	0.30252 (14)	0.0212 (5)
C11	0.5102 (4)	0.1654 (2)	0.40740 (13)	0.0201 (5)
C12	0.6223 (4)	0.2566 (2)	0.42595 (13)	0.0204 (5)
C13	0.5791 (4)	0.07309 (19)	0.42009 (13)	0.0210 (5)
H13	0.7092	0.0697	0.4295	0.025*
C14	-0.2025 (4)	0.3016 (2)	0.25629 (14)	0.0257 (5)
H14A	-0.2471	0.3720	0.2565	0.039*
H14B	-0.3011	0.2567	0.2738	0.039*
H14C	-0.1695	0.2817	0.2087	0.039*
C15	0.3630 (4)	0.5270 (2)	0.31404 (15)	0.0255 (6)
H15A	0.3776	0.6010	0.3111	0.038*
H15B	0.2969	0.5024	0.2727	0.038*
H15C	0.4864	0.4951	0.3165	0.038*
C16	0.4774 (4)	-0.0247 (2)	0.42131 (13)	0.0204 (5)
C17	0.5616 (4)	-0.1163 (2)	0.40254 (13)	0.0220 (5)
C18	0.4756 (4)	-0.2095 (2)	0.41364 (14)	0.0252 (6)
H18	0.5356	-0.2707	0.3999	0.030*
C19	0.3015 (5)	-0.21243 (19)	0.44500 (14)	0.0261 (6)
H19	0.2434	-0.2759	0.4542	0.031*
C20	0.2116 (4)	-0.1224 (2)	0.46308 (13)	0.0251 (5)
H20	0.0909	-0.1244	0.4835	0.030*
C21	0.2988 (4)	-0.02945 (19)	0.45117 (13)	0.0228 (5)
H21	0.2364	0.0317	0.4634	0.027*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Br1	0.03169 (14)	0.02821 (14)	0.03656 (15)	0.00934 (12)	0.00802 (12)	0.00322 (13)
O1	0.0264 (10)	0.0225 (9)	0.0254 (9)	-0.0003 (8)	-0.0026 (8)	0.0020 (8)
O2	0.0249 (9)	0.0140 (8)	0.0256 (9)	-0.0014 (7)	-0.0045 (7)	0.0001 (7)
O3	0.0383 (12)	0.0152 (9)	0.0319 (11)	0.0016 (8)	-0.0109 (9)	-0.0040 (8)
C1	0.0231 (11)	0.0163 (12)	0.0212 (11)	-0.0009 (10)	0.0027 (9)	0.0025 (10)
C2	0.0257 (13)	0.0145 (12)	0.0355 (15)	0.0004 (10)	-0.0034 (11)	0.0028 (11)
C3	0.0301 (13)	0.0148 (12)	0.0281 (14)	0.0020 (10)	0.0017 (11)	0.0004 (10)
C4	0.0289 (12)	0.0112 (11)	0.0224 (13)	-0.0009 (9)	-0.0046 (10)	-0.0001 (9)
C5	0.0231 (11)	0.0120 (10)	0.0204 (10)	-0.0016 (12)	0.0008 (9)	0.0005 (9)
C6	0.0242 (11)	0.0148 (11)	0.0185 (12)	-0.0012 (9)	-0.0028 (10)	-0.0006 (10)
C7	0.0258 (12)	0.0119 (11)	0.0218 (12)	-0.0010 (10)	0.0007 (10)	-0.0009 (9)
C8	0.0304 (14)	0.0162 (12)	0.0246 (13)	0.0016 (10)	-0.0023 (11)	-0.0038 (10)
C9	0.0290 (13)	0.0172 (13)	0.0316 (14)	0.0001 (11)	-0.0044 (11)	-0.0055 (11)
C10	0.0229 (12)	0.0187 (13)	0.0221 (13)	0.0003 (10)	0.0010 (10)	0.0000 (10)
C11	0.0261 (13)	0.0175 (13)	0.0168 (12)	0.0002 (10)	0.0008 (10)	0.0004 (10)
C12	0.0254 (13)	0.0176 (12)	0.0182 (12)	-0.0011 (10)	0.0006 (10)	0.0013 (10)

C13	0.0255 (13)	0.0184 (12)	0.0191 (12)	-0.0010 (10)	0.0022 (10)	0.0006 (10)
C14	0.0236 (12)	0.0222 (13)	0.0312 (14)	-0.0014 (12)	-0.0005 (12)	-0.0016 (11)
C15	0.0262 (13)	0.0179 (13)	0.0323 (15)	-0.0013 (10)	-0.0030 (11)	0.0043 (11)
C16	0.0277 (13)	0.0157 (12)	0.0179 (12)	0.0002 (10)	-0.0028 (11)	0.0009 (10)
C17	0.0243 (12)	0.0205 (12)	0.0211 (12)	0.0033 (12)	-0.0017 (9)	0.0009 (11)
C18	0.0360 (15)	0.0148 (13)	0.0247 (14)	0.0045 (11)	-0.0032 (12)	-0.0021 (11)
C19	0.0361 (14)	0.0163 (12)	0.0259 (13)	-0.0040 (13)	-0.0051 (14)	0.0014 (10)
C20	0.0275 (12)	0.0214 (12)	0.0264 (12)	-0.0042 (13)	-0.0001 (11)	0.0025 (11)
C21	0.0279 (12)	0.0155 (12)	0.0251 (12)	0.0024 (12)	-0.0004 (13)	0.0019 (10)

*Geometric parameters (Å, °)*

Br1—C17	1.899 (3)	C8—H8B	0.9900
O1—C12	1.203 (3)	C9—C10	1.533 (4)
O2—C12	1.364 (3)	C9—H9A	0.9900
O2—C6	1.470 (3)	C9—H9B	0.9900
O3—C4	1.435 (3)	C10—C14	1.514 (4)
O3—H3	0.77 (4)	C11—C13	1.335 (4)
C1—C10	1.333 (4)	C11—C12	1.487 (4)
C1—C5	1.525 (3)	C13—C16	1.479 (4)
C1—C2	1.532 (4)	C13—H13	0.9500
C2—C3	1.527 (4)	C14—H14A	0.9800
C2—H2A	0.9900	C14—H14B	0.9800
C2—H2B	0.9900	C14—H14C	0.9800
C3—C4	1.527 (4)	C15—H15A	0.9800
C3—H3A	0.9900	C15—H15B	0.9800
C3—H3B	0.9900	C15—H15C	0.9800
C4—C15	1.525 (4)	C16—C17	1.396 (4)
C4—C5	1.555 (3)	C16—C21	1.403 (4)
C5—C6	1.502 (3)	C17—C18	1.389 (4)
C5—H5	1.0000	C18—C19	1.386 (4)
C6—C7	1.535 (3)	C18—H18	0.9500
C6—H6	1.0000	C19—C20	1.393 (4)
C7—C11	1.514 (4)	C19—H19	0.9500
C7—C8	1.524 (4)	C20—C21	1.392 (4)
C7—H7	1.0000	C20—H20	0.9500
C8—C9	1.532 (4)	C21—H21	0.9500
C8—H8A	0.9900		
C12—O2—C6	110.22 (19)	C8—C9—H9A	106.9
C4—O3—H3	106 (3)	C10—C9—H9A	106.9
C10—C1—C5	130.0 (2)	C8—C9—H9B	106.9
C10—C1—C2	123.2 (2)	C10—C9—H9B	106.9
C5—C1—C2	106.8 (2)	H9A—C9—H9B	106.7
C3—C2—C1	105.8 (2)	C1—C10—C14	120.2 (2)
C3—C2—H2A	110.6	C1—C10—C9	128.6 (2)
C1—C2—H2A	110.6	C14—C10—C9	110.8 (2)
C3—C2—H2B	110.6	C13—C11—C12	119.4 (2)

C1—C2—H2B	110.6	C13—C11—C7	132.7 (2)
H2A—C2—H2B	108.7	C12—C11—C7	107.5 (2)
C4—C3—C2	104.4 (2)	O1—C12—O2	121.6 (2)
C4—C3—H3A	110.9	O1—C12—C11	129.3 (2)
C2—C3—H3A	110.9	O2—C12—C11	109.1 (2)
C4—C3—H3B	110.9	C11—C13—C16	127.8 (2)
C2—C3—H3B	110.9	C11—C13—H13	116.1
H3A—C3—H3B	108.9	C16—C13—H13	116.1
O3—C4—C15	109.8 (2)	C10—C14—H14A	109.5
O3—C4—C3	109.3 (2)	C10—C14—H14B	109.5
C15—C4—C3	111.6 (2)	H14A—C14—H14B	109.5
O3—C4—C5	111.4 (2)	C10—C14—H14C	109.5
C15—C4—C5	112.9 (2)	H14A—C14—H14C	109.5
C3—C4—C5	101.7 (2)	H14B—C14—H14C	109.5
C6—C5—C1	114.0 (2)	C4—C15—H15A	109.5
C6—C5—C4	115.0 (2)	C4—C15—H15B	109.5
C1—C5—C4	103.7 (2)	H15A—C15—H15B	109.5
C6—C5—H5	107.9	C4—C15—H15C	109.5
C1—C5—H5	107.9	H15A—C15—H15C	109.5
C4—C5—H5	107.9	H15B—C15—H15C	109.5
O2—C6—C5	109.59 (19)	C17—C16—C21	117.4 (2)
O2—C6—C7	105.82 (19)	C17—C16—C13	122.3 (2)
C5—C6—C7	114.6 (2)	C21—C16—C13	119.7 (2)
O2—C6—H6	108.9	C18—C17—C16	122.1 (2)
C5—C6—H6	108.9	C18—C17—Br1	117.7 (2)
C7—C6—H6	108.9	C16—C17—Br1	120.2 (2)
C11—C7—C8	118.8 (2)	C19—C18—C17	119.3 (3)
C11—C7—C6	102.0 (2)	C19—C18—H18	120.3
C8—C7—C6	109.8 (2)	C17—C18—H18	120.3
C11—C7—H7	108.6	C18—C19—C20	120.1 (2)
C8—C7—H7	108.6	C18—C19—H19	120.0
C6—C7—H7	108.6	C20—C19—H19	120.0
C7—C8—C9	112.1 (2)	C21—C20—C19	119.9 (2)
C7—C8—H8A	109.2	C21—C20—H20	120.0
C9—C8—H8A	109.2	C19—C20—H20	120.0
C7—C8—H8B	109.2	C20—C21—C16	121.0 (3)
C9—C8—H8B	109.2	C20—C21—H21	119.5
H8A—C8—H8B	107.9	C16—C21—H21	119.5
C8—C9—C10	121.8 (2)		
C10—C1—C2—C3	178.0 (2)	C2—C1—C10—C14	1.5 (4)
C5—C1—C2—C3	-1.5 (3)	C5—C1—C10—C9	8.9 (5)
C1—C2—C3—C4	26.5 (3)	C2—C1—C10—C9	-170.6 (3)
C2—C3—C4—O3	-158.4 (2)	C8—C9—C10—C1	-39.4 (4)
C2—C3—C4—C15	80.0 (3)	C8—C9—C10—C14	147.9 (3)
C2—C3—C4—C5	-40.6 (3)	C8—C7—C11—C13	-48.2 (4)
C10—C1—C5—C6	31.2 (4)	C6—C7—C11—C13	-169.0 (3)
C2—C1—C5—C6	-149.3 (2)	C8—C7—C11—C12	138.5 (2)



C10—C1—C5—C4	157.0 (3)	C6—C7—C11—C12	17.7 (3)
C2—C1—C5—C4	-23.5 (3)	C6—O2—C12—O1	173.0 (2)
O3—C4—C5—C6	-79.2 (3)	C6—O2—C12—C11	-9.5 (3)
C15—C4—C5—C6	44.9 (3)	C13—C11—C12—O1	-3.0 (4)
C3—C4—C5—C6	164.5 (2)	C7—C11—C12—O1	171.3 (3)
O3—C4—C5—C1	155.6 (2)	C13—C11—C12—O2	179.7 (2)
C15—C4—C5—C1	-80.3 (2)	C7—C11—C12—O2	-6.0 (3)
C3—C4—C5—C1	39.3 (2)	C12—C11—C13—C16	164.5 (2)
C12—O2—C6—C5	145.0 (2)	C7—C11—C13—C16	-8.2 (5)
C12—O2—C6—C7	21.0 (3)	C11—C13—C16—C17	148.9 (3)
C1—C5—C6—O2	171.08 (19)	C11—C13—C16—C21	-40.2 (4)
C4—C5—C6—O2	51.4 (3)	C21—C16—C17—C18	-1.1 (4)
C1—C5—C6—C7	-70.1 (3)	C13—C16—C17—C18	170.0 (3)
C4—C5—C6—C7	170.2 (2)	C21—C16—C17—Br1	178.06 (18)
O2—C6—C7—C11	-22.9 (2)	C13—C16—C17—Br1	-10.8 (3)
C5—C6—C7—C11	-143.8 (2)	C16—C17—C18—C19	-0.7 (4)
O2—C6—C7—C8	-149.8 (2)	Br1—C17—C18—C19	-179.9 (2)
C5—C6—C7—C8	89.3 (3)	C17—C18—C19—C20	2.0 (4)
C11—C7—C8—C9	168.4 (2)	C18—C19—C20—C21	-1.6 (4)
C6—C7—C8—C9	-74.9 (3)	C19—C20—C21—C16	-0.3 (4)
C7—C8—C9—C10	57.4 (4)	C17—C16—C21—C20	1.6 (4)
C5—C1—C10—C14	-179.1 (2)	C13—C16—C21—C20	-169.8 (2)

*Hydrogen-bond geometry (Å, °)*

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
O3—H3...O2	0.77 (4)	2.26 (4)	2.883 (3)	139 (4)