

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1-(Furan-2-yl)-2-(2H-indazol-2-yl)ethanone

Özden Özel Güven,^a Gökhan Türk,^a Philip D. F. Adler,^b Simon J. Coles^b and Tuncer Hökelek^{c*}^aDepartment of Chemistry, Bülent Ecevit University, 67100 Zonguldak, Turkey,^bDepartment of Chemistry, Southampton University, SO17 1BJ Southampton, England, and ^cDepartment of Physics, Hacettepe University, 06800 Beytepe, Ankara, Turkey

Correspondence e-mail: merzifon@hacettepe.edu.tr


Received 19 March 2014; accepted 25 March 2014

Key indicators: single-crystal X-ray study; $T = 294$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.044; wR factor = 0.127; data-to-parameter ratio = 17.1.

The asymmetric unit of the title compound, $\text{C}_{13}\text{H}_{10}\text{N}_2\text{O}_2$, contains two crystallographically independent molecules (*A* and *B*). The indazole ring systems are approximately planar [maximum deviations = 0.0037 (15) and -0.0198 (15) Å], and their mean planes are oriented at 80.10 (5) and 65.97 (4)° with respect to the furan rings in molecules *A* and *B*, respectively. In the crystal, pairs of $\text{C}-\text{H}\cdots\text{N}$ hydrogen bonds link the *B* molecules, forming inversion dimers. These dimers are bridged by the *A* molecules via $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds, forming sheets parallel to (011). There are also $\text{C}-\text{H}\cdots\pi$ interactions present, and $\pi-\pi$ interactions between neighbouring furan and the indazole rings [centroid-centroid distance = 3.8708 (9) Å] of inversion-related molecules, forming a three-dimensional structure.

Related literature

For related structures, see: Peeters *et al.* (1979); Freer *et al.* (1986); Özel Güven *et al.* (2008*a,b*, 2013, 2014).



Experimental

Crystal data

$\text{C}_{13}\text{H}_{10}\text{N}_2\text{O}_2$
 $M_r = 226.23$
 Triclinic, $P\bar{1}$
 $a = 9.2899$ (3) Å
 $b = 10.6863$ (4) Å

$c = 11.7826$ (5) Å
 $\alpha = 77.046$ (3)°
 $\beta = 70.780$ (3)°
 $\gamma = 88.930$ (4)°
 $V = 1074.47$ (7) Å³

$Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹

$T = 294$ K
 $0.17 \times 0.15 \times 0.10$ mm

Data collection

Rigaku Saturn724+ diffractometer
 Absorption correction: multi-scan
 (*CrystalClear*; Rigaku, 2011)
 $T_{\min} = 0.984$, $T_{\max} = 0.990$

10517 measured reflections
 5256 independent reflections
 4130 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.030$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.127$
 $S = 1.14$
 5256 reflections

307 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.35$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.33$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$\text{Cg}1$, $\text{Cg}2$, $\text{Cg}5$ and $\text{Cg}6$ are the centroids of the $\text{O}2/\text{C}2-\text{C}5$, $\text{N}1/\text{N}2/\text{C}7/\text{C}8/\text{C}13$, $\text{N}1'/\text{N}2'/\text{C}7'/\text{C}8'/\text{C}13'$, and $\text{C}8'-\text{C}13'$ rings, respectively.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}3'-\text{H}3'\cdots\text{N}2^{\text{iv}}$	0.93	2.57	3.4031 (18)	149
$\text{C}7-\text{H}7\cdots\text{O}1^{\text{iii}}$	0.93	2.48	3.2544 (17)	141
$\text{C}10'-\text{H}10'\cdots\text{O}1^{\text{iii}}$	0.93	2.44	3.2719 (17)	148
$\text{C}7-\text{H}7\cdots\text{Cg}5^{\text{iv}}$	0.93	2.97	3.6000 (16)	126
$\text{C}9-\text{H}9\cdots\text{Cg}6^{\text{iv}}$	0.93	2.81	3.4312 (17)	125
$\text{C}9'-\text{H}9'\cdots\text{Cg}2^{\text{v}}$	0.93	2.94	3.6743 (15)	137
$\text{C}12'-\text{H}12'\cdots\text{Cg}1^{\text{i}}$	0.93	2.63	3.4480 (16)	147

Symmetry codes: (i) $-x+1, -y+1, -z$; (ii) $-x, -y+1, -z+1$; (iii) $x, y+1, z-1$; (iv) $x+1, y, z$; (v) $-x+1, -y+2, -z$.

Data collection: *CrystalClear* (Rigaku, 2011); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *WinGX* (Farrugia, 2012) and *PLATON* (Spek, 2009).

The authors acknowledge the Zonguldak Karaelmas University Research Fund (project No. 2012-10-03-12).

Supporting information for this paper is available from the IUCr electronic archives (Reference: SU2716).

References

- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
 Freer, A. A., Pearson, A. & Salole, E. G. (1986). *Acta Cryst.* **C42**, 1350–1352.
 Özel Güven, Ö., Erdoğan, T., Coles, S. J. & Hökelek, T. (2008*a*). *Acta Cryst.* **E64**, o1358.
 Özel Güven, Ö., Tahtacı, H., Coles, S. J. & Hökelek, T. (2008*b*). *Acta Cryst.* **E64**, o1604.
 Özel Güven, Ö., Türk, G., Adler, P. D. F., Coles, S. J. & Hökelek, T. (2013). *Acta Cryst.* **E69**, o184.
 Özel Güven, Ö., Türk, G., Adler, P. D. F., Coles, S. J. & Hökelek, T. (2014). *Acta Cryst.* **E70**, o410.
 Peeters, O. M., Blaton, N. M. & De Ranter, C. J. (1979). *Acta Cryst.* **B35**, 2461–2464.
 Rigaku (2011). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supporting information

Acta Cryst. (2014). E70, o505 [doi:10.1107/S1600536814006606]

1-(Furan-2-yl)-2-(2H-indazol-2-yl)ethanone

Özden Özel Güven, Gökhan Türk, Philip D. F. Adler, Simon J. Coles and Tuncer Hökelek

S1. Comment

Azole group containing ketones are starting materials of important antifungal compounds like miconazole (Peeters *et al.*, 1979) and econazole (Freer *et al.*, 1986) having an ether structure. The crystal structures of some phenyl ketones having a benzimidazole ring (Özel Güven *et al.*, 2008a), a 1,2,4-triazole ring (Özel Güven *et al.*, 2008b) and an indazole ring (Özel Güven *et al.*, 2013) have been reported. The crystal structure of indazole ring containing an ether structure has also been reported (Özel Güven *et al.*, 2014). Herein we report on the crystal structure of the title indazole derivative.

The asymmetric unit of the title compound contains two crystallographically independent molecules (A and B), shown in Fig. 1. The bond lengths and angles are generally within normal ranges. The indazole ring systems in the two molecules [N1/N2/C7—C13 and N1'/N2'/C7'—C13'] are approximately planar [maximum deviations of 0.0037 (15) Å for atom C12 and -0.0198 (15) Å for atom C13']. Their mean planes are oriented with respect to the furan rings [O2/C2—C5 and O2'/C2'—C5'] at dihedral angles of 80.10 (5) and 65.97 (4) °, in molecules A and B, respectively. Atoms C6 and C6' are displaced by -0.0508 (14) and 0.1004 (15) Å from their respective indazole ring plane, while atoms O1, C1, C6 and O1', C1', C6' are displaced by 0.0239 (11), -0.0104 (14), -0.1155 (14) and -0.0504 (10), -0.0166 (14), 0.0371 (15) Å from their respective furan ring mean plane.

In the crystal, C—H...N hydrogen bonds link the B molecules forming inversion dimers. These dimers are bridged by the A molecules, via C—H...O hydrogen bonds, forming sheets parallel to (011), see Table 1 and Fig. 2. Weak C—H... π interactions (Table 1) are also present together with π ... π interactions between furan and the indazole rings of inversion related molecules Cg1—Cg4ⁱ, [centroid-centroid distance = 3.8708 (9) Å; symmetry code: (i) -x+I, -y+I, -z+I; Cg1 and Cg4 are the centroids of the rings O2/C2—C5 and N1'/N2'/C7'—C13'] forming a three-dimensional structure.

S2. Experimental

2-Acetylfuran (3.303 g, 30.01 mmol) was dissolved in 18 ml dioxane/ether (1:2) mixture. This solution was cooled in ice-water, and then bromine (1.96 ml, 38.14 mmol) was added over a period of 30 min. The reaction mixture was stirred for 12 h at room temperature. At the end of this period, saturated ammonium chloride solution (25 ml) was added and the solution extracted with ether. After evaporation of the ether, the residue obtained was purified by column chromatography using a hexane-ethylacetate (60:1) mixture. The resulting furacyl bromide (2.500 g, 13.23 mmol) was dissolved in toluene (30 ml). 1H-indazole (3.125 g, 26.45 mmol) was added in small quantities and the reaction mixture was refluxed for 12 h. The solvent was evaporated and the resulting residue was purified by column chromatography using chloroform as eluent. The title ketone was crystallized from 2-propanol to obtain colourless crystals suitable for X-ray analysis [yield 0.9 g; 30%].

S3. Refinement

H atoms were positioned geometrically with C—H = 0.93 and 0.97 Å for aromatic and methylene H, respectively, and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.




Figure 1

The molecular structure of the two independent molecules (A and B) of the title compound, with atom labelling. Displacement ellipsoids are drawn at the 50% probability level.




Figure 2

A view of the crystal packing of the title compound with the hydrogen bonds shown as dashed lines [see Table 1 for details; H-atoms not involved in hydrogen bonding have been omitted for clarity].

1-(Furan-2-yl)-2-(2H-indazol-2-yl)ethanone

Crystal data

$C_{13}H_{10}N_2O_2$

$M_r = 226.23$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 9.2899\ (3)\ \text{\AA}$

$b = 10.6863\ (4)\ \text{\AA}$

$c = 11.7826\ (5)\ \text{\AA}$

$\alpha = 77.046\ (3)^\circ$

$\beta = 70.780\ (3)^\circ$

$\gamma = 88.930\ (4)^\circ$

$V = 1074.47\ (7)\ \text{\AA}^3$

$Z = 4$

$F(000) = 472$

$D_x = 1.399\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 9141 reflections

$\theta = 3.0\text{--}28.7^\circ$

$\mu = 0.10\ \text{mm}^{-1}$

$T = 294$ K $0.17 \times 0.15 \times 0.10$ mm
 Block, colourless

Data collection

Rigaku Saturn724+ diffractometer	10517 measured reflections 5256 independent reflections
Radiation source: fine-focus sealed tube	4130 reflections with $I > 2\sigma(I)$
Graphite monochromator	$R_{\text{int}} = 0.030$
ω scans	$\theta_{\text{max}} = 28.7^\circ$, $\theta_{\text{min}} = 3.0^\circ$
Absorption correction: multi-scan (<i>CrystalClear</i> ; Rigaku, 2011)	$h = -12 \rightarrow 11$ $k = -14 \rightarrow 14$ $l = -15 \rightarrow 15$
$T_{\text{min}} = 0.984$, $T_{\text{max}} = 0.990$	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.044$	H-atom parameters constrained
$wR(F^2) = 0.127$	$w = 1/[\sigma^2(F_o^2) + (0.0629P)^2 + 0.1259P]$
$S = 1.14$	where $P = (F_o^2 + 2F_c^2)/3$
5256 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
307 parameters	$\Delta\rho_{\text{max}} = 0.35 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.33 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.34655 (12)	0.40513 (9)	0.72286 (9)	0.0301 (2)
O2	0.12038 (12)	0.66910 (9)	0.67289 (10)	0.0306 (2)
N1	0.13004 (13)	0.35998 (10)	0.95859 (10)	0.0222 (2)
N2	0.22341 (13)	0.35403 (11)	1.02685 (10)	0.0240 (3)
C1	0.24816 (16)	0.48234 (12)	0.74176 (12)	0.0227 (3)
C2	0.23765 (16)	0.58932 (12)	0.64362 (13)	0.0238 (3)
C3	0.32561 (18)	0.63034 (14)	0.52231 (13)	0.0295 (3)
H3	0.4122	0.5936	0.4795	0.035*
C4	0.2570 (2)	0.74121 (15)	0.47503 (14)	0.0356 (4)
H4	0.2902	0.7910	0.3945	0.043*
C5	0.13524 (19)	0.76044 (14)	0.56853 (15)	0.0358 (4)
H5	0.0703	0.8270	0.5625	0.043*
C6	0.13153 (16)	0.47746 (12)	0.86879 (12)	0.0236 (3)
H6A	0.1536	0.5500	0.8986	0.028*

H6B	0.0307	0.4862	0.8613	0.028*
C7	0.03917 (15)	0.25281 (12)	0.98598 (12)	0.0228 (3)
H7	-0.0318	0.2389	0.9492	0.027*
C8	0.07307 (15)	0.16724 (12)	1.08045 (12)	0.0220 (3)
C9	0.02026 (16)	0.04070 (13)	1.15061 (13)	0.0267 (3)
H9	-0.0550	-0.0043	1.1372	0.032*
C10	0.08333 (17)	-0.01324 (13)	1.23873 (14)	0.0297 (3)
H10	0.0501	-0.0961	1.2859	0.036*
C11	0.19855 (18)	0.05475 (14)	1.25960 (14)	0.0302 (3)
H11	0.2388	0.0151	1.3203	0.036*
C12	0.25205 (17)	0.17660 (13)	1.19336 (13)	0.0263 (3)
H12	0.3278	0.2199	1.2079	0.032*
C13	0.18851 (15)	0.23470 (12)	1.10204 (12)	0.0221 (3)
O1'	0.12771 (11)	0.65903 (9)	0.22019 (9)	0.0277 (2)
O2'	0.12290 (11)	0.46552 (9)	0.41588 (8)	0.0256 (2)
N1'	0.36805 (13)	0.72809 (11)	0.00180 (10)	0.0221 (2)
N2'	0.36774 (13)	0.71328 (10)	-0.10991 (10)	0.0215 (2)
C1'	0.23839 (15)	0.59371 (12)	0.21100 (12)	0.0212 (3)
C2'	0.24799 (15)	0.48921 (12)	0.31045 (11)	0.0210 (3)
C3'	0.35788 (16)	0.40672 (12)	0.32360 (12)	0.0238 (3)
H3'	0.4531	0.4032	0.2654	0.029*
C4'	0.29694 (18)	0.32704 (13)	0.44471 (13)	0.0281 (3)
H4'	0.3444	0.2603	0.4814	0.034*
C5'	0.15650 (17)	0.36727 (13)	0.49609 (13)	0.0270 (3)
H5'	0.0917	0.3321	0.5759	0.032*
C6'	0.37597 (16)	0.61447 (13)	0.09316 (12)	0.0252 (3)
H6C	0.4680	0.6225	0.1133	0.030*
H6D	0.3827	0.5397	0.0582	0.030*
C7'	0.36543 (16)	0.85057 (13)	0.01246 (12)	0.0238 (3)
H7'	0.3641	0.8792	0.0817	0.029*
C8'	0.36502 (15)	0.92660 (12)	-0.10033 (12)	0.0214 (3)
C9'	0.36551 (15)	1.05980 (13)	-0.15096 (12)	0.0242 (3)
H9'	0.3624	1.1197	-0.1039	0.029*
C10'	0.37064 (16)	1.09798 (13)	-0.27098 (13)	0.0264 (3)
H10'	0.3718	1.1852	-0.3060	0.032*
C11'	0.37424 (17)	1.00751 (13)	-0.34351 (12)	0.0267 (3)
H11'	0.3779	1.0372	-0.4249	0.032*
C12'	0.37248 (16)	0.87798 (12)	-0.29719 (12)	0.0236 (3)
H12'	0.3747	0.8196	-0.3455	0.028*
C13'	0.36711 (15)	0.83599 (12)	-0.17304 (12)	0.0201 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0299 (6)	0.0234 (5)	0.0321 (6)	0.0032 (4)	-0.0042 (4)	-0.0062 (4)
O2	0.0290 (6)	0.0273 (5)	0.0327 (6)	0.0029 (4)	-0.0084 (4)	-0.0044 (4)
N1	0.0198 (6)	0.0240 (5)	0.0223 (6)	-0.0006 (4)	-0.0066 (4)	-0.0047 (4)
N2	0.0224 (6)	0.0257 (6)	0.0248 (6)	-0.0011 (5)	-0.0091 (5)	-0.0057 (5)

C1	0.0226 (7)	0.0200 (6)	0.0248 (7)	-0.0029 (5)	-0.0065 (5)	-0.0056 (5)
C2	0.0238 (7)	0.0217 (6)	0.0266 (7)	0.0010 (5)	-0.0080 (6)	-0.0076 (5)
C3	0.0294 (8)	0.0318 (7)	0.0267 (7)	-0.0066 (6)	-0.0045 (6)	-0.0118 (6)
C4	0.0491 (10)	0.0320 (8)	0.0268 (8)	-0.0105 (7)	-0.0180 (7)	0.0002 (6)
C5	0.0393 (9)	0.0277 (7)	0.0426 (9)	0.0008 (6)	-0.0216 (8)	-0.0003 (6)
C6	0.0228 (7)	0.0220 (6)	0.0238 (7)	0.0001 (5)	-0.0059 (5)	-0.0038 (5)
C7	0.0186 (6)	0.0255 (7)	0.0242 (7)	-0.0013 (5)	-0.0061 (5)	-0.0068 (5)
C8	0.0184 (6)	0.0239 (6)	0.0225 (6)	-0.0008 (5)	-0.0044 (5)	-0.0065 (5)
C9	0.0232 (7)	0.0247 (7)	0.0297 (7)	-0.0031 (5)	-0.0055 (6)	-0.0058 (5)
C10	0.0306 (8)	0.0229 (7)	0.0304 (7)	-0.0006 (6)	-0.0066 (6)	-0.0011 (5)
C11	0.0320 (8)	0.0302 (7)	0.0293 (8)	0.0051 (6)	-0.0128 (6)	-0.0049 (6)
C12	0.0246 (7)	0.0281 (7)	0.0283 (7)	0.0006 (5)	-0.0113 (6)	-0.0069 (6)
C13	0.0207 (6)	0.0219 (6)	0.0220 (6)	-0.0002 (5)	-0.0049 (5)	-0.0049 (5)
O1'	0.0231 (5)	0.0331 (5)	0.0226 (5)	0.0041 (4)	-0.0050 (4)	-0.0017 (4)
O2'	0.0238 (5)	0.0299 (5)	0.0177 (5)	-0.0005 (4)	-0.0031 (4)	-0.0005 (4)
N1'	0.0214 (6)	0.0248 (6)	0.0168 (5)	-0.0008 (4)	-0.0037 (4)	-0.0021 (4)
N2'	0.0216 (6)	0.0241 (5)	0.0169 (5)	-0.0010 (4)	-0.0044 (4)	-0.0040 (4)
C1'	0.0207 (7)	0.0247 (6)	0.0190 (6)	-0.0004 (5)	-0.0072 (5)	-0.0053 (5)
C2'	0.0213 (6)	0.0244 (6)	0.0157 (6)	-0.0019 (5)	-0.0044 (5)	-0.0039 (5)
C3'	0.0264 (7)	0.0249 (6)	0.0201 (6)	0.0009 (5)	-0.0075 (5)	-0.0057 (5)
C4'	0.0378 (8)	0.0242 (7)	0.0237 (7)	0.0020 (6)	-0.0138 (6)	-0.0025 (5)
C5'	0.0319 (8)	0.0259 (7)	0.0200 (6)	-0.0035 (6)	-0.0083 (6)	0.0008 (5)
C6'	0.0225 (7)	0.0276 (7)	0.0204 (6)	0.0028 (5)	-0.0041 (5)	-0.0004 (5)
C7'	0.0232 (7)	0.0282 (7)	0.0201 (6)	-0.0022 (5)	-0.0058 (5)	-0.0075 (5)
C8'	0.0187 (6)	0.0238 (6)	0.0206 (6)	-0.0016 (5)	-0.0040 (5)	-0.0064 (5)
C9'	0.0219 (7)	0.0242 (6)	0.0255 (7)	-0.0017 (5)	-0.0038 (5)	-0.0099 (5)
C10'	0.0260 (7)	0.0209 (6)	0.0271 (7)	-0.0010 (5)	-0.0040 (6)	-0.0025 (5)
C11'	0.0303 (7)	0.0265 (7)	0.0199 (7)	0.0007 (6)	-0.0061 (6)	-0.0018 (5)
C12'	0.0272 (7)	0.0236 (6)	0.0200 (6)	-0.0008 (5)	-0.0069 (5)	-0.0060 (5)
C13'	0.0180 (6)	0.0210 (6)	0.0198 (6)	-0.0012 (5)	-0.0045 (5)	-0.0045 (5)

Geometric parameters (Å, °)

O1—C1	1.2190 (17)	O1'—C1'	1.2207 (16)
O2—C2	1.3720 (17)	O2'—C2'	1.3701 (16)
O2—C5	1.3562 (18)	O2'—C5'	1.3528 (16)
N1—N2	1.3556 (15)	N1'—C6'	1.4496 (16)
N1—C6	1.4462 (16)	N1'—C7'	1.3417 (17)
N1—C7	1.3488 (16)	N2'—N1'	1.3614 (15)
N2—C13	1.3550 (16)	N2'—C13'	1.3559 (16)
C1—C2	1.4598 (18)	C1'—C6'	1.5214 (18)
C1—C6	1.5219 (19)	C2'—C1'	1.4514 (18)
C2—C3	1.366 (2)	C2'—C3'	1.3610 (19)
C3—C4	1.425 (2)	C3'—C4'	1.4223 (19)
C3—H3	0.9300	C3'—H3'	0.9300
C4—C5	1.344 (2)	C4'—H4'	0.9300
C4—H4	0.9300	C5'—C4'	1.353 (2)
C5—H5	0.9300	C5'—H5'	0.9300

C6—H6A	0.9700	C6'—H6C	0.9700
C6—H6B	0.9700	C6'—H6D	0.9700
C7—H7	0.9300	C7'—H7'	0.9300
C8—C7	1.3922 (19)	C8'—C7'	1.3948 (18)
C8—C9	1.4174 (18)	C9'—C8'	1.4141 (18)
C9—H9	0.9300	C9'—C10'	1.3660 (19)
C10—C9	1.369 (2)	C9'—H9'	0.9300
C10—C11	1.420 (2)	C10'—C11'	1.4202 (19)
C10—H10	0.9300	C10'—H10'	0.9300
C11—H11	0.9300	C11'—H11'	0.9300
C12—C11	1.3655 (19)	C12'—C11'	1.3675 (18)
C12—H12	0.9300	C12'—C13'	1.4150 (18)
C13—C8	1.4231 (18)	C12'—H12'	0.9300
C13—C12	1.4130 (19)	C13'—C8'	1.4255 (18)
C5—O2—C2	106.78 (12)	C5'—O2'—C2'	106.24 (11)
N2—N1—C6	119.17 (10)	N2'—N1'—C6'	118.38 (11)
C7—N1—N2	114.56 (11)	C7'—N1'—N2'	114.33 (11)
C7—N1—C6	126.24 (11)	C7'—N1'—C6'	127.26 (11)
C13—N2—N1	103.11 (10)	C13'—N2'—N1'	102.97 (10)
O1—C1—C2	121.81 (13)	O1'—C1'—C2'	122.77 (12)
O1—C1—C6	122.96 (12)	O1'—C1'—C6'	122.35 (11)
C2—C1—C6	115.20 (12)	C2'—C1'—C6'	114.88 (12)
O2—C2—C1	117.61 (12)	O2'—C2'—C1'	115.92 (12)
C3—C2—O2	109.94 (12)	C3'—C2'—O2'	110.48 (11)
C3—C2—C1	132.44 (14)	C3'—C2'—C1'	133.60 (12)
C2—C3—C4	105.59 (14)	C2'—C3'—C4'	105.77 (13)
C2—C3—H3	127.2	C2'—C3'—H3'	127.1
C4—C3—H3	127.2	C4'—C3'—H3'	127.1
C3—C4—H4	126.4	O2'—C5'—H5'	124.5
C5—C4—C3	107.20 (13)	C4'—C5'—O2'	110.93 (12)
C5—C4—H4	126.4	C4'—C5'—H5'	124.5
O2—C5—H5	124.8	C3'—C4'—H4'	126.7
C4—C5—O2	110.48 (14)	C5'—C4'—C3'	106.58 (13)
C4—C5—H5	124.8	C5'—C4'—H4'	126.7
N1—C6—C1	113.33 (11)	N1'—C6'—C1'	112.61 (11)
N1—C6—H6A	108.9	N1'—C6'—H6C	109.1
N1—C6—H6B	108.9	N1'—C6'—H6D	109.1
C1—C6—H6A	108.9	C1'—C6'—H6C	109.1
C1—C6—H6B	108.9	C1'—C6'—H6D	109.1
H6A—C6—H6B	107.7	H6C—C6'—H6D	107.8
N1—C7—C8	106.09 (11)	N1'—C7'—C8'	106.76 (11)
N1—C7—H7	127.0	N1'—C7'—H7'	126.6
C8—C7—H7	127.0	C8'—C7'—H7'	126.6
C7—C8—C9	135.26 (13)	C7'—C8'—C9'	135.85 (12)
C7—C8—C13	104.46 (11)	C7'—C8'—C13'	103.89 (11)
C9—C8—C13	120.28 (12)	C9'—C8'—C13'	120.25 (12)
C8—C9—H9	121.0	C8'—C9'—H9'	121.0

C10—C9—C8	118.01 (13)	C10'—C9'—C8'	118.06 (12)
C10—C9—H9	121.0	C10'—C9'—H9'	121.0
C9—C10—C11	121.33 (13)	C9'—C10'—C11'	121.52 (12)
C9—C10—H10	119.3	C9'—C10'—H10'	119.2
C11—C10—H10	119.3	C11'—C10'—H10'	119.2
C10—C11—H11	119.0	C10'—C11'—H11'	119.0
C12—C11—C10	122.04 (13)	C12'—C11'—C10'	121.97 (12)
C12—C11—H11	119.0	C12'—C11'—H11'	119.0
C11—C12—C13	117.78 (13)	C11'—C12'—C13'	117.58 (12)
C11—C12—H12	121.1	C11'—C12'—H12'	121.2
C13—C12—H12	121.1	C13'—C12'—H12'	121.2
N2—C13—C8	111.78 (12)	N2'—C13'—C8'	112.04 (11)
N2—C13—C12	127.66 (12)	N2'—C13'—C12'	127.32 (12)
C12—C13—C8	120.56 (12)	C12'—C13'—C8'	120.61 (11)
C5—O2—C2—C1	179.45 (12)	C5'—O2'—C2'—C1'	-179.04 (11)
C5—O2—C2—C3	0.22 (15)	C5'—O2'—C2'—C3'	0.50 (14)
C2—O2—C5—C4	0.01 (16)	C2'—O2'—C5'—C4'	-0.84 (15)
C6—N1—N2—C13	177.98 (11)	N2'—N1'—C6'—C1'	-119.87 (13)
C7—N1—N2—C13	0.09 (15)	C7'—N1'—C6'—C1'	62.35 (18)
N2—N1—C6—C1	89.29 (14)	N2'—N1'—C7'—C8'	-0.71 (16)
C7—N1—C6—C1	-93.08 (15)	C6'—N1'—C7'—C8'	177.15 (12)
N2—N1—C7—C8	0.11 (16)	C13'—N2'—N1'—C6'	-177.16 (11)
C6—N1—C7—C8	-177.61 (12)	C13'—N2'—N1'—C7'	0.91 (15)
N1—N2—C13—C8	-0.25 (15)	N1'—N2'—C13'—C8'	-0.76 (14)
N1—N2—C13—C12	-179.97 (14)	N1'—N2'—C13'—C12'	177.36 (13)
O1—C1—C2—O2	178.16 (12)	O1'—C1'—C6'—N1'	7.19 (18)
O1—C1—C2—C3	-2.8 (2)	C2'—C1'—C6'—N1'	-173.78 (11)
C6—C1—C2—O2	-3.81 (17)	O2'—C2'—C1'—O1'	1.42 (19)
C6—C1—C2—C3	175.21 (14)	O2'—C2'—C1'—C6'	-177.60 (11)
O1—C1—C6—N1	-11.17 (18)	C3'—C2'—C1'—O1'	-177.99 (14)
C2—C1—C6—N1	170.83 (11)	C3'—C2'—C1'—C6'	3.0 (2)
O2—C2—C3—C4	-0.35 (15)	O2'—C2'—C3'—C4'	0.00 (15)
C1—C2—C3—C4	-179.43 (14)	C1'—C2'—C3'—C4'	179.43 (14)
C2—C3—C4—C5	0.35 (16)	C2'—C3'—C4'—C5'	-0.50 (15)
C3—C4—C5—O2	-0.22 (17)	O2'—C5'—C4'—C3'	0.85 (16)
C9—C8—C7—N1	179.74 (15)	C9'—C8'—C7'—N1'	-178.62 (15)
C13—C8—C7—N1	-0.25 (15)	C13'—C8'—C7'—N1'	0.19 (15)
C7—C8—C9—C10	-179.87 (16)	C10'—C9'—C8'—C7'	177.57 (15)
C13—C8—C9—C10	0.1 (2)	C10'—C9'—C8'—C13'	-1.1 (2)
C11—C10—C9—C8	-0.2 (2)	C8'—C9'—C10'—C11'	0.5 (2)
C9—C10—C11—C12	0.0 (2)	C9'—C10'—C11'—C12'	0.1 (2)
C13—C12—C11—C10	0.2 (2)	C13'—C12'—C11'—C10'	-0.1 (2)
N2—C13—C8—C7	0.32 (16)	C11'—C12'—C13'—N2'	-178.52 (13)
N2—C13—C8—C9	-179.67 (12)	C11'—C12'—C13'—C8'	-0.5 (2)
C12—C13—C8—C7	-179.94 (13)	N2'—C13'—C8'—C7'	0.38 (15)
C12—C13—C8—C9	0.1 (2)	N2'—C13'—C8'—C9'	179.41 (12)
N2—C13—C12—C11	179.48 (14)	C12'—C13'—C8'—C7'	-177.89 (12)

C8—C13—C12—C11	-0.2 (2)	C12'—C13'—C8'—C9'	1.2 (2)
----------------	----------	-------------------	---------

Hydrogen-bond geometry (Å, °)

Cg1, Cg2, Cg5 and Cg6 are the centroids of the O2/C2—C5, N1/N2/C7/C8/C13, N1'/N2'/C7'/C8'/C13', and C8'—C13' rings, respectively.

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
C3'—H3'···N2' ⁱ	0.93	2.57	3.4031 (18)	149
C7—H7···O1' ⁱⁱ	0.93	2.48	3.2544 (17)	141
C10'—H10'···O1' ⁱⁱⁱ	0.93	2.44	3.2719 (17)	148
C7—H7···Cg5 ^{iv}	0.93	2.97	3.6000 (16)	126
C9—H9···Cg6 ^{iv}	0.93	2.81	3.4312 (17)	125
C9'—H9'···Cg2 ^v	0.93	2.94	3.6743 (15)	137
C12'—H12'···Cg1 ⁱ	0.93	2.63	3.4480 (16)	147

Symmetry codes: (i) $-x+1, -y+1, -z$; (ii) $-x, -y+1, -z+1$; (iii) $x, y+1, z-1$; (iv) $x+1, y, z$; (v) $-x+1, -y+2, -z$.