

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 2-Amino-6-methylpyridinium 4-methylbenzenesulfonate

K. Syed Suresh Babu,<sup>a,b</sup> M. Dhavamurthy,<sup>a</sup>  
M. NizamMohideen,<sup>b\*</sup> G. Peramaiyan<sup>a</sup> and R. Mohan<sup>a\*</sup><sup>a</sup>Department of Physics, Presidency College (Autonomous), Chennai 600 005, Tamil Nadu, India, and <sup>b</sup>Department of Physics, The New College (Autonomous), Chennai 600 014, Tamil Nadu, India

Correspondence e-mail: mnizam\_new@yahoo.in, professorsomohan@yahoo.co.in


Received 10 April 2014; accepted 15 April 2014

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.040;  $wR$  factor = 0.119; data-to-parameter ratio = 16.8.

In the asymmetric unit of the title salt,  $\text{C}_6\text{H}_9\text{N}_2^+ \cdot \text{C}_7\text{H}_7\text{O}_3\text{S}^-$ , there are two independent 2-amino-6-methylpyridinium cations and two independent 4-methylbenzenesulfonate anions. Both cations are protonated at their pyridine N atoms and their geometries reveal amine–imine tautomerism. In the 4-methylbenzenesulfonate anions, the carboxylate groups are twisted out of the benzene ring planes by 88.4 (1) and 86.2 (2)°. In the crystal, the sulfonate O atoms of an anion interact with the protonated N atoms and the 2-amino groups of a cation *via* a pair of  $\text{N}-\text{H} \cdots \text{O}$  hydrogen bonds, forming an  $R_2^2(8)$  ring motif. These motifs are connected *via*  $\text{N}-\text{H} \cdots \text{O}$  hydrogen bonds, forming chains running along the  $a$ -axis direction. Within the chains there are weak  $\text{C}-\text{H} \cdots \text{O}$  hydrogen bonds present. In addition, aromatic  $\pi-\pi$  stacking interactions [centroid–centroid distances = 3.771 (2), 3.599 (2), 3.599 (2) and 3.497 (2) Å] involving neighbouring chains are also observed.

## Related literature

For crystal structures of related pyridine derivatives and their applications, see: Babu *et al.* (2014); Rajkumar *et al.* (2014); Jin *et al.* (2005). For unprotonated aminopyridine derivatives, see: Anderson *et al.* (2005). For the structure of amino-methylpyridinium, see: Nahringerbauer & Kvik (1977). For details of sulfonates, see: Onoda *et al.* (2001); Baskar Raj *et al.* (2003). For applications of benzenesulfonic acid, see: Wang & Wei (2007). For simple organic–inorganic salts containing strong intermolecular hydrogen bonds, see: Sethuram *et al.* (2013*a,b*); Shihabuddeen Syed *et al.* (2013); Showrilu *et al.* (2013); Huq *et al.* (2013). For bond-length data, see: Allen *et al.* (1987). For studies on the tautomeric forms of 2-aminopyridine systems, see: Ishikawa *et al.* (2002). For graph-set analysis, see: Etter (1990); Bernstein *et al.* (1995).



## Experimental

## Crystal data

$\text{C}_6\text{H}_9\text{N}_2^+ \cdot \text{C}_7\text{H}_7\text{O}_3\text{S}^-$   
 $M_r = 280.35$   
 Triclinic,  $P\bar{1}$   
 $a = 7.5343$  (2) Å  
 $b = 13.6212$  (5) Å  
 $c = 13.9887$  (5) Å  
 $\alpha = 106.307$  (2)°  
 $\beta = 97.946$  (1)°

$\gamma = 92.103$  (2)°  
 $V = 1360.31$  (8) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.24$  mm<sup>-1</sup>  
 $T = 293$  K  
 $0.35 \times 0.25 \times 0.20$  mm

## Data collection

Bruker Kappa APEXII CCD diffractometer  
 Absorption correction: multi-scan (SADABS; Sheldrick, 2004)  
 $T_{\min} = 0.920$ ,  $T_{\max} = 0.953$

32534 measured reflections  
 6237 independent reflections  
 4709 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.026$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$   
 $wR(F^2) = 0.119$   
 $S = 1.06$   
 6237 reflections  
 372 parameters  
 6 restraints

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.33$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.37$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{N1}-\text{H1A} \cdots \text{O2}$	0.90 (1)	1.88 (1)	2.772 (2)	171 (2)
$\text{N2}-\text{H2A} \cdots \text{O3}^i$	0.87 (1)	2.01 (1)	2.880 (2)	174 (2)
$\text{N2}-\text{H2B} \cdots \text{O1}$	0.88 (1)	2.07 (1)	2.919 (2)	162 (2)
$\text{N3}-\text{H3A} \cdots \text{O5}$	0.89 (1)	1.90 (1)	2.789 (2)	174 (2)
$\text{N4}-\text{H4A} \cdots \text{O4}^{ii}$	0.88 (1)	2.02 (1)	2.882 (2)	167 (2)
$\text{N4}-\text{H4B} \cdots \text{O6}$	0.88 (1)	2.04 (1)	2.883 (2)	162 (2)
$\text{C22}-\text{H22} \cdots \text{O5}^{ii}$	0.93	2.58	3.455 (2)	157

Symmetry codes: (i)  $x - 1, y, z$ ; (ii)  $x + 1, y, z$ .

Data collection: APEX2 (Bruker, 2004); cell refinement: APEX2 and SAINT (Bruker, 2004); data reduction: SAINT and XPREP (Bruker, 2004); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 2012) and Mercury (Macrae *et al.*, 2008); software used to prepare material for publication: WinGX (Farrugia, 2012) and PLATON (Spek, 2009).

The authors are thankful to the SAIF, IIT Madras, for the data collection.

Supporting information for this paper is available from the IUCr electronic archives (Reference: SU2726).

## References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Anderson, F. P., Gallagher, J. F., Kenny, P. T. M. & Lough, A. J. (2005). *Acta Cryst. E61*, o1350–o1353.
- Babu, K. S. S., Peramaiyan, G., NizamMohideen, M. & Mohan, R. (2014). *Acta Cryst. E70*, o391–o392.
- Baskar Raj, S., Sethuraman, V., Francis, S., Hemamalini, M., Muthiah, P. T., Bocelli, G., Cantoni, A., Rychlewska, U. & Warzajtis, B. (2003). *CrystEngComm*, **5**, 70–76.
- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2004). *APEX2, SAINT and XPREP*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Etter, M. C. (1990). *Acc. Chem. Res.* **23**, 120–126.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Huq, C. A. M. A., Fouzia, S. & NizamMohideen, M. (2013). *Acta Cryst. E69*, o1766–o1767.
- Ishikawa, H., Iwata, K. & Hamaguchi, H. (2002). *J. Phys. Chem. A*, **106**, 2305–2312.
- Jin, Z.-M., Shun, N., Lü, Y.-P., Hu, M.-L. & Shen, L. (2005). *Acta Cryst. C61*, m43–m45.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Nahringbauer, I. & Kvik, Å. (1977). *Acta Cryst. B33*, 2902–2905.
- Onoda, A., Yamada, Y., Doi, M., Okamura, T. & Ueyama, N. (2001). *Inorg. Chem.* **40**, 516–521.
- Rajkumar, M. A., Xavier, S. S. J., Anbarasu, S., Devarajan, P. A. & NizamMohideen, M. (2014). *Acta Cryst. E70*, o473–o474.
- Sethuram, M., Bhargavi, G., Dhandapani, M., Amirthaganesan, G. & NizamMohideen, M. (2013a). *Acta Cryst. E69*, o1301–o1302.
- Sethuram, M., Rajasekharan, M. V., Dhandapani, M., Amirthaganesan, G. & NizamMohideen, M. (2013b). *Acta Cryst. E69*, o957–o958.
- Sheldrick, G. M. (2004). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Shihabuddeen Syed, A., Rajarajan, K. & NizamMohideen, M. (2013). *Acta Cryst. E69*, i33.
- Showrilu, K., Rajarajan, K. & NizamMohideen, M. (2013). *Acta Cryst. E69*, m469–m470.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
- Wang, Z.-L. & Wei, L.-H. (2007). *Acta Cryst. E63*, o1448–o1449.

## supporting information

*Acta Cryst.* (2014). E70, o600–o601 [doi:10.1107/S1600536814008587]

## 2-Amino-6-methylpyridinium 4-methylbenzenesulfonate

K. Syed Suresh Babu, M. Dhavamurthy, M. NizamMohideen, G. Peramaiyan and R. Mohan

### S1. Introduction

2-Aminopyridine and its derivatives play an important role in heterocyclic chemistry. Pyridine heterocycles and their derivatives are present in many large molecules having photo-chemical, electro-chemical and catalytic applications (Babu *et al.*, 2014). Simple organic-inorganic salts containing strong intermolecular hydrogen bonds have attracted an attention as materials which display ferroelectric-paraelectric phase transitions (Sethuram, *et al.*, 2013*a,b*; Huq *et al.*, 2013; Shihabuddeen Syed *et al.*, 2013; Showrilu *et al.*, 2013). Hydrogen-bonding patterns involving sulfonate groups in biological systems and metal complexes are of current interest (Onoda *et al.*, 2001). Such interactions can be utilized for designing supramolecular architectures (Baskar Raj *et al.*, 2003). Benzenesulfonic acid, is a particularly strong organic acid which is capable of protonating N-containing heterocycles and other Lewis bases (Wang & Wei, 2007). We have recently reported the crystal structures of 2-amino-6-methylpyridinium 2,2,2-trichloroacetate (Babu *et al.*, 2014) and 2-Amino-5-nitropyridinium hydrogen oxalate (Rajkumar *et al.*, 2014). In continuation of our studies of pyridinium derivatives, the crystal structure determination of the title compound has been undertaken.

### S2. Comment / Result and Discussion

The asymmetric unit of title salt, Fig. 1, consists of two crystallographically independent protonated 2-amino-6-methylpyridinium cation and two crystallographically independent 4-methyl benzenesulfonate anions. The normal probability plot analyses (International Tables for X-ray Crystallography, 1974, Vol. IV, pp. 293–309) for both bond lengths and angles show that the differences between the two symmetry independent molecules are of a statistical nature. All bond lengths (Allen *et al.*, 1987) and angles are within normal ranges and comparable with those in closely related structures (Babu *et al.*, 2014; Rajkumar *et al.*, 2014). A proton transfer from the carboxyl group of *p*-toluenesulfonic acid to atom N1 and N3 of 2-amino-6-methyl pyridine resulted in the formation of a salt. This protonation lead to the widening of the C8—N1—C12 and C21—N3—C25 angles of the pyridine rings to 124.0 (2) ° and 123.8 (2) °, compared to 115.3 (2) ° in the unprotonated aminopyridine (Anderson *et al.*, 2005). This type of protonation is observed in various aminopyridine acid complexes (Babu *et al.*, 2014; Rajkumar *et al.*, 2014).

In the cation, the N2—C8 [1.325 (2) Å] N4—C21 bonds [1.325 (2) Å] is shorter than the N1—C8 [1.347 (2) Å], N1—C12 [1.360 (2) Å], N3—C21 [1.352 (2) Å] and N3—C25 [1.362 (2) Å] bonds, and the C8—C9 [1.406 (3) Å], C10—C11 [1.398 (3) Å], C21—C22 [1.405 (3) Å] and C23—C24 [1.401 (3) Å] bonds are significantly longer than C9—C10 [1.357 (3) Å], C11—C12 [1.356 (3) Å], C22—C23 [1.357 (3) Å] and C24—C25 [1.353 (3) Å] bonds, are similar to those in the amino-methylpyridinium cation (Babu *et al.*, 2014; Rajkumar *et al.*, 2014). In contrast, in the solid state structure of amino-methylpyridinium, the N—C bond out of ring is clearly longer than that in the ring (Nahringbauer *et al.*, 1977). The geometrical features of the amino-methylpyridinium cation (N1/N2/C1/C6 and N3/N4/C9—C13) resemble those observed in other 2-aminopyridinium structures (Babu *et al.*, 2014; Rajkumar *et al.*, 2014) that are believed to be involved in amine-imine tautomerism (Ishikawa *et al.*, 2002). Similar features are also provided by cation amino-methyl-

pyridinium (N3/N4/C7/C12). However, previous study show that a pyridinium cation always possesses an expanded angle of C—N—C in comparison with the parent pyridine (Jin *et al.*, 2005).

The examination of pyridinium rings shows that these units are planar with mean deviation of -0.006 (2) and 0.005 (2) Å for atoms C8 and C21, from the mean planes defined by the six constituent atoms. The dihedral angle between the 2-amino-6-methylpyridinium cation and 4-methylbenzenesulfonate anion group is 88.4 (2) and 86.2 (2)° for the both molecules, respectively. In both the molecules, the protonated 2-amino-6-methylpyridinium cation is essentially planar, with maximum deviations of -0.012 (2) for atom C13 and -0.006 (2) Å for atom C25.

### S3. Hydrogen bonding interaction / Intermolecular N—H···O and C—H···O interaction

In the crystal (Fig. 2), the protonated atoms (N1 and N3) and a nitrogen atom of the 2-amino groups (N2 and N4) of the 2-amino-6-methylpyridinium cations are hydrogen bonded to the carboxylate oxygen atoms (O1, O2, O3 and O4) of the sulfonate groups of the *p*-toluenesulfonate anions *via* a pair of intermolecular N—H···O hydrogen bonds (Table 1), forming a ring motif with a graph-set notation of  $R^2_2(8)$  [Etter, 1990; Bernstein *et al.*, 1995]. The sulfonate group mimics the carboxylate anion's mode of association, which is more commonly seen when binding with 2-aminopyrimidines. It is well known that sulfonates imitate carboxylates in forming such bidentate motifs (Baskar Raj *et al.*, 2003).

Furthermore, these motifs are connected *via* N—H···O hydrogen bonds (Fig. 2 and Table 1), involving the 2-amino group of the 2-amino-6-methyl pyridinium cation and atoms O3 and O4 of an anion, to form a supramolecular chains along the *a* axis direction. Weak C—H···O hydrogen bonds, involving a pyridine group of the cation and an O atom of a sulfonate anion, within the chains are also observed (Fig. 2 and Table 1).

### S4. Aromatic interaction

In addition, the cations of neighbouring chains are linked through aromatic  $\pi$ - $\pi$  interactions with centroid distances  $Cg1\cdots Cg1^{iii} = 3.771$  (2),  $Cg1\cdots Cg2^{iv} = 3.599$  (2),  $Cg2\cdots Cg1^v = 3.599$  (2) and  $Cg2\cdots Cg2^{vi} = 3.497$  (2) Å [symmetry codes are as in Table 1 and (iii) = -x+1, -y+1, -z+2; (iv) = x, y, z+ (v) = x, y, z+1; (vi) = -x+1, -y, -z; Cg1 and Cg2 are the centroids of the N1/C8—C12 and N3/C21—C25 rings, respectively].

### S5. Uses


The identification of such supramolecular patterns will help us design and construct preferred hydrogen bonding patterns of drug like molecules.

### S6. Synthesis and crystallization


Crystals of the title compound were obtained by slow evaporation of a 1:1 equimolar mixture of 2-amino-6-methylpyridine and benzenesulfonic acid in methanol at room temperature.

### S7. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 1. N-bound H atoms were located in a difference Fourier map and refined with distance restraints: N—H = 0.88 (1) and 0.90 (1) Å for NH<sub>2</sub> and NH H atoms, respectively. The C-bound H atoms were positioned geometrically and refined using a riding model: C—H = 0.93–0.96 Å with  $U_{iso}(H) = 1.5U_{eq}(C\text{-methyl})$  and  $= 1.2U_{eq}(C)$  for other H atoms. A rotating group model was used for the methyl group.

**Figure 1**

A view of the molecular structure of the two independent benzenesulfonate anions and the two independent 2-amino-6-methylpyridinium cations of the title salt. Displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**

The crystal packing of the title compound, viewed along the *b* axis. The N—H···O and C—H···O hydrogen bonds are shown as dashed lines (see Table 1 for details; H atoms not involved in hydrogen bonding have been omitted for clarity).

## 2-Amino-6-methylpyridinium 4-methylbenzenesulfonate

## Crystal data

C<sub>6</sub>H<sub>9</sub>N<sub>2</sub><sup>+</sup>·C<sub>7</sub>H<sub>7</sub>O<sub>3</sub>S<sup>-</sup> $M_r = 280.35$ Triclinic,  $P\bar{1}$ 

Hall symbol: -P 1

 $a = 7.5343$  (2) Å $b = 13.6212$  (5) Å $c = 13.9887$  (5) Å $\alpha = 106.307$  (2)° $\beta = 97.946$  (1)° $\gamma = 92.103$  (2)° $V = 1360.31$  (8) Å<sup>3</sup> $Z = 4$  $F(000) = 592$  $D_x = 1.369$  Mg m<sup>-3</sup>Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 6237 reflections

 $\theta = 2.0$ – $28.1$ ° $\mu = 0.24$  mm<sup>-1</sup> $T = 293$  K

Block, colourless

 $0.35 \times 0.25 \times 0.20$  mm

## Data collection

Bruker Kappa APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 $\omega$  and  $\varphi$  scans

Absorption correction: multi-scan

(SADABS; Sheldrick, 2004)

 $T_{\min} = 0.920$ ,  $T_{\max} = 0.953$ 

32534 measured reflections

6237 independent reflections

4709 reflections with  $I > 2\sigma(I)$  $R_{\text{int}} = 0.026$  $\theta_{\text{max}} = 27.5$ °,  $\theta_{\text{min}} = 1.6$ ° $h = -9 \rightarrow 9$  $k = -17 \rightarrow 17$  $l = -18 \rightarrow 18$ 

## Refinement

Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.040$  $wR(F^2) = 0.119$  $S = 1.06$ 

6237 reflections

372 parameters

6 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier

map

Hydrogen site location: inferred from  
neighbouring sitesH atoms treated by a mixture of independent  
and constrained refinement $w = 1/[\sigma^2(F_o^2) + (0.0502P)^2 + 0.655P]$ where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\text{max}} = 0.001$  $\Delta\rho_{\text{max}} = 0.33$  e Å<sup>-3</sup> $\Delta\rho_{\text{min}} = -0.37$  e Å<sup>-3</sup>Extinction correction: SHELXL97 (Sheldrick,  
2008),  $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$ 

Extinction coefficient: 0.0067 (10)

## Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>)

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.7573 (2)	0.33148 (14)	0.65370 (12)	0.0340 (4)
C2	0.6441 (3)	0.28156 (17)	0.56563 (15)	0.0510 (5)

---

H2	0.5822	0.2192	0.5594	0.061*
C3	0.6232 (3)	0.32433 (19)	0.48708 (16)	0.0568 (6)
H3	0.5472	0.2897	0.4279	0.068*
C4	0.7111 (3)	0.41650 (17)	0.49332 (15)	0.0450 (5)
C5	0.8219 (3)	0.46630 (18)	0.58213 (16)	0.0525 (5)
H5	0.8818	0.5293	0.5886	0.063*
C6	0.8455 (3)	0.42432 (17)	0.66170 (15)	0.0498 (5)
H6	0.9213	0.4590	0.7209	0.060*
C7	0.6848 (3)	0.4621 (2)	0.40658 (18)	0.0658 (7)
H7A	0.7653	0.4341	0.3602	0.099*
H7B	0.7093	0.5352	0.4315	0.099*
H7C	0.5630	0.4461	0.3727	0.099*
O1	0.63184 (18)	0.20570 (11)	0.73960 (10)	0.0473 (3)
O2	0.80180 (19)	0.36019 (11)	0.84663 (10)	0.0482 (3)
O3	0.95409 (17)	0.22525 (12)	0.74752 (10)	0.0487 (4)
S1	0.78890 (6)	0.27601 (4)	0.75373 (3)	0.03657 (13)
C14	0.1612 (2)	0.16545 (14)	0.35185 (12)	0.0335 (4)
C15	0.2427 (3)	0.07546 (17)	0.34413 (15)	0.0497 (5)
H15	0.2824	0.0416	0.2843	0.060*
C16	0.2656 (3)	0.03531 (17)	0.42480 (16)	0.0532 (5)
H16	0.3205	-0.0258	0.4186	0.064*
C17	0.2085 (3)	0.08433 (16)	0.51461 (14)	0.0427 (4)
C18	0.1255 (3)	0.17361 (18)	0.52060 (15)	0.0522 (5)
H18	0.0851	0.2074	0.5802	0.063*
C19	0.1007 (3)	0.21434 (16)	0.44031 (15)	0.0471 (5)
H19	0.0434	0.2746	0.4459	0.057*
C20	0.2366 (3)	0.0415 (2)	0.60338 (17)	0.0610 (6)
H20A	0.3489	0.0705	0.6450	0.092*
H20B	0.2384	-0.0317	0.5797	0.092*
H20C	0.1403	0.0584	0.6420	0.092*
O4	-0.01417 (17)	0.27792 (12)	0.25616 (11)	0.0496 (4)
O5	0.13593 (18)	0.13852 (11)	0.15948 (9)	0.0471 (3)
O6	0.30945 (17)	0.28838 (11)	0.27035 (10)	0.0453 (3)
S2	0.14592 (6)	0.22210 (4)	0.25238 (3)	0.03629 (13)
C21	0.5816 (2)	0.16145 (13)	0.08860 (13)	0.0327 (4)
C22	0.7160 (2)	0.14424 (14)	0.02675 (14)	0.0368 (4)
H22	0.8366	0.1588	0.0550	0.044*
C23	0.6682 (2)	0.10619 (15)	-0.07461 (14)	0.0407 (4)
H23	0.7569	0.0954	-0.1157	0.049*
C24	0.4871 (2)	0.08288 (14)	-0.11818 (14)	0.0392 (4)
H24	0.4556	0.0568	-0.1877	0.047*
C25	0.3584 (2)	0.09863 (13)	-0.05818 (13)	0.0340 (4)
C26	0.1613 (2)	0.07772 (16)	-0.09541 (15)	0.0452 (5)
H26A	0.1106	0.1403	-0.0996	0.068*
H26B	0.1047	0.0509	-0.0496	0.068*
H26C	0.1419	0.0285	-0.1609	0.068*
N3	0.40875 (19)	0.13737 (11)	0.04333 (11)	0.0325 (3)
N4	0.6161 (2)	0.19844 (15)	0.18808 (12)	0.0445 (4)

C8	0.3602 (2)	0.33977 (13)	0.91967 (13)	0.0334 (4)
C9	0.2264 (2)	0.35815 (15)	0.98204 (14)	0.0393 (4)
H9	0.1055	0.3440	0.9543	0.047*
C10	0.2761 (3)	0.39686 (16)	1.08322 (15)	0.0449 (5)
H10	0.1883	0.4085	1.1248	0.054*
C11	0.4573 (3)	0.41950 (16)	1.12590 (14)	0.0432 (4)
H11	0.4896	0.4463	1.1953	0.052*
C12	0.5854 (2)	0.40210 (13)	1.06520 (13)	0.0348 (4)
C13	0.7826 (3)	0.42082 (16)	1.10162 (15)	0.0460 (5)
H13A	0.8040	0.4713	1.1664	0.069*
H13B	0.8402	0.4453	1.0547	0.069*
H13C	0.8305	0.3580	1.1072	0.069*
N1	0.53289 (19)	0.36343 (11)	0.96397 (11)	0.0326 (3)
N2	0.3236 (2)	0.30260 (14)	0.82019 (13)	0.0449 (4)
H2A	0.2112 (15)	0.2825 (16)	0.7953 (15)	0.050 (6)*
H4B	0.531 (2)	0.2199 (16)	0.2235 (14)	0.050 (6)*
H4A	0.7276 (16)	0.2181 (17)	0.2166 (16)	0.058 (7)*
H2B	0.409 (2)	0.2798 (17)	0.7850 (15)	0.055 (7)*
H1A	0.619 (2)	0.3551 (17)	0.9244 (14)	0.051 (6)*
H3A	0.322 (2)	0.1430 (16)	0.0815 (13)	0.045 (6)*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0265 (8)	0.0429 (10)	0.0285 (8)	0.0011 (7)	0.0034 (6)	0.0043 (7)
C2	0.0497 (12)	0.0530 (12)	0.0423 (11)	-0.0155 (9)	-0.0094 (9)	0.0114 (10)
C3	0.0541 (13)	0.0694 (15)	0.0389 (11)	-0.0148 (11)	-0.0147 (9)	0.0157 (10)
C4	0.0383 (10)	0.0593 (12)	0.0387 (10)	0.0060 (9)	0.0052 (8)	0.0166 (9)
C5	0.0553 (13)	0.0522 (12)	0.0472 (12)	-0.0124 (10)	0.0042 (10)	0.0140 (10)
C6	0.0486 (11)	0.0569 (13)	0.0341 (10)	-0.0154 (10)	-0.0048 (8)	0.0050 (9)
C7	0.0627 (15)	0.0858 (18)	0.0588 (15)	0.0065 (13)	0.0054 (11)	0.0386 (14)
O1	0.0392 (7)	0.0579 (9)	0.0471 (8)	-0.0029 (6)	0.0109 (6)	0.0176 (7)
O2	0.0492 (8)	0.0618 (9)	0.0292 (7)	0.0042 (7)	0.0089 (6)	0.0047 (6)
O3	0.0341 (7)	0.0696 (10)	0.0431 (8)	0.0140 (6)	0.0050 (6)	0.0164 (7)
S1	0.0285 (2)	0.0514 (3)	0.0289 (2)	0.00414 (18)	0.00579 (16)	0.00929 (19)
C14	0.0279 (8)	0.0418 (10)	0.0281 (8)	0.0000 (7)	0.0043 (6)	0.0060 (7)
C15	0.0591 (13)	0.0552 (13)	0.0343 (10)	0.0181 (10)	0.0157 (9)	0.0065 (9)
C16	0.0624 (13)	0.0512 (12)	0.0461 (12)	0.0192 (10)	0.0076 (10)	0.0129 (10)
C17	0.0374 (10)	0.0538 (12)	0.0359 (10)	-0.0053 (8)	0.0003 (8)	0.0151 (9)
C18	0.0590 (13)	0.0651 (14)	0.0342 (10)	0.0124 (11)	0.0193 (9)	0.0106 (10)
C19	0.0527 (12)	0.0499 (12)	0.0413 (11)	0.0156 (9)	0.0171 (9)	0.0111 (9)
C20	0.0603 (14)	0.0771 (17)	0.0494 (13)	-0.0044 (12)	-0.0013 (10)	0.0302 (12)
O4	0.0337 (7)	0.0652 (9)	0.0513 (8)	0.0065 (6)	-0.0015 (6)	0.0226 (7)
O5	0.0446 (7)	0.0630 (9)	0.0281 (7)	-0.0099 (6)	0.0044 (5)	0.0064 (6)
O6	0.0335 (7)	0.0574 (9)	0.0437 (7)	-0.0099 (6)	0.0020 (6)	0.0163 (6)
S2	0.0269 (2)	0.0506 (3)	0.0298 (2)	-0.00331 (18)	0.00094 (16)	0.01147 (19)
C21	0.0292 (8)	0.0340 (9)	0.0360 (9)	0.0009 (7)	0.0005 (7)	0.0140 (7)
C22	0.0264 (8)	0.0400 (10)	0.0446 (10)	0.0042 (7)	0.0037 (7)	0.0140 (8)



C23	0.0367 (9)	0.0460 (11)	0.0420 (10)	0.0081 (8)	0.0109 (8)	0.0144 (9)
C24	0.0405 (10)	0.0438 (10)	0.0328 (9)	0.0034 (8)	0.0024 (7)	0.0117 (8)
C25	0.0330 (9)	0.0334 (9)	0.0367 (9)	-0.0004 (7)	-0.0012 (7)	0.0154 (7)
C26	0.0346 (10)	0.0527 (12)	0.0464 (11)	-0.0040 (8)	-0.0051 (8)	0.0177 (9)
N3	0.0269 (7)	0.0378 (8)	0.0345 (8)	0.0008 (6)	0.0036 (6)	0.0140 (6)
N4	0.0319 (8)	0.0622 (11)	0.0351 (9)	-0.0002 (8)	0.0003 (7)	0.0098 (8)
C8	0.0309 (8)	0.0348 (9)	0.0366 (9)	0.0069 (7)	0.0038 (7)	0.0139 (7)
C9	0.0303 (9)	0.0456 (11)	0.0445 (10)	0.0088 (7)	0.0074 (7)	0.0154 (8)
C10	0.0436 (10)	0.0532 (12)	0.0445 (11)	0.0133 (9)	0.0175 (9)	0.0185 (9)
C11	0.0497 (11)	0.0494 (11)	0.0316 (9)	0.0083 (9)	0.0052 (8)	0.0133 (8)
C12	0.0379 (9)	0.0333 (9)	0.0353 (9)	0.0038 (7)	0.0016 (7)	0.0149 (7)
C13	0.0397 (10)	0.0508 (12)	0.0457 (11)	-0.0001 (9)	-0.0040 (8)	0.0167 (9)
N1	0.0301 (7)	0.0365 (8)	0.0325 (8)	0.0055 (6)	0.0055 (6)	0.0116 (6)
N2	0.0328 (8)	0.0611 (11)	0.0365 (9)	0.0045 (8)	0.0028 (7)	0.0081 (8)

*Geometric parameters (Å, °)*

C1—C6	1.375 (3)	O6—S2	1.4497 (13)
C1—C2	1.380 (2)	C21—N4	1.325 (2)
C1—S1	1.7609 (18)	C21—N3	1.352 (2)
C2—C3	1.375 (3)	C21—C22	1.405 (3)
C2—H2	0.9300	C22—C23	1.357 (3)
C3—C4	1.372 (3)	C22—H22	0.9300
C3—H3	0.9300	C23—C24	1.401 (3)
C4—C5	1.378 (3)	C23—H23	0.9300
C4—C7	1.504 (3)	C24—C25	1.353 (3)
C5—C6	1.381 (3)	C24—H24	0.9300
C5—H5	0.9300	C25—N3	1.362 (2)
C6—H6	0.9300	C25—C26	1.493 (2)
C7—H7A	0.9600	C26—H26A	0.9600
C7—H7B	0.9600	C26—H26B	0.9600
C7—H7C	0.9600	C26—H26C	0.9600
O1—S1	1.4499 (14)	N3—H3A	0.894 (9)
O2—S1	1.4605 (14)	N4—H4B	0.876 (9)
O3—S1	1.4469 (13)	N4—H4A	0.876 (10)
C14—C15	1.375 (3)	C8—N2	1.325 (2)
C14—C19	1.378 (3)	C8—N1	1.347 (2)
C14—S2	1.7636 (18)	C8—C9	1.406 (3)
C15—C16	1.379 (3)	C9—C10	1.357 (3)
C15—H15	0.9300	C9—H9	0.9300
C16—C17	1.382 (3)	C10—C11	1.398 (3)
C16—H16	0.9300	C10—H10	0.9300
C17—C18	1.375 (3)	C11—C12	1.356 (3)
C17—C20	1.507 (3)	C11—H11	0.9300
C18—C19	1.380 (3)	C12—N1	1.360 (2)
C18—H18	0.9300	C12—C13	1.491 (3)
C19—H19	0.9300	C13—H13A	0.9600
C20—H20A	0.9600	C13—H13B	0.9600

C20—H20B	0.9600	C13—H13C	0.9600
C20—H20C	0.9600	N1—H1A	0.900 (9)
O4—S2	1.4485 (14)	N2—H2A	0.873 (10)
O5—S2	1.4582 (14)	N2—H2B	0.877 (10)
C6—C1—C2	119.11 (18)	O4—S2—C14	106.93 (8)
C6—C1—S1	120.76 (13)	O6—S2—C14	106.19 (8)
C2—C1—S1	120.13 (15)	O5—S2—C14	106.66 (8)
C3—C2—C1	119.76 (19)	N4—C21—N3	118.92 (16)
C3—C2—H2	120.1	N4—C21—C22	123.44 (16)
C1—C2—H2	120.1	N3—C21—C22	117.63 (16)
C4—C3—C2	122.01 (18)	C23—C22—C21	119.40 (16)
C4—C3—H3	119.0	C23—C22—H22	120.3
C2—C3—H3	119.0	C21—C22—H22	120.3
C3—C4—C5	117.70 (19)	C22—C23—C24	120.92 (17)
C3—C4—C7	120.99 (19)	C22—C23—H23	119.5
C5—C4—C7	121.3 (2)	C24—C23—H23	119.5
C4—C5—C6	121.2 (2)	C25—C24—C23	119.41 (17)
C4—C5—H5	119.4	C25—C24—H24	120.3
C6—C5—H5	119.4	C23—C24—H24	120.3
C1—C6—C5	120.25 (18)	C24—C25—N3	118.87 (16)
C1—C6—H6	119.9	C24—C25—C26	124.50 (17)
C5—C6—H6	119.9	N3—C25—C26	116.63 (16)
C4—C7—H7A	109.5	C25—C26—H26A	109.5
C4—C7—H7B	109.5	C25—C26—H26B	109.5
H7A—C7—H7B	109.5	H26A—C26—H26B	109.5
C4—C7—H7C	109.5	C25—C26—H26C	109.5
H7A—C7—H7C	109.5	H26A—C26—H26C	109.5
H7B—C7—H7C	109.5	H26B—C26—H26C	109.5
O3—S1—O1	113.06 (9)	C21—N3—C25	123.75 (15)
O3—S1—O2	111.56 (8)	C21—N3—H3A	119.2 (13)
O1—S1—O2	111.86 (8)	C25—N3—H3A	117.0 (13)
O3—S1—C1	107.19 (8)	C21—N4—H4B	121.1 (15)
O1—S1—C1	106.16 (8)	C21—N4—H4A	118.4 (16)
O2—S1—C1	106.52 (9)	H4B—N4—H4A	118 (2)
C15—C14—C19	119.43 (18)	N2—C8—N1	119.09 (16)
C15—C14—S2	120.43 (14)	N2—C8—C9	123.12 (16)
C19—C14—S2	120.03 (15)	N1—C8—C9	117.77 (16)
C14—C15—C16	120.19 (18)	C10—C9—C8	119.13 (17)
C14—C15—H15	119.9	C10—C9—H9	120.4
C16—C15—H15	119.9	C8—C9—H9	120.4
C15—C16—C17	121.08 (19)	C9—C10—C11	121.08 (18)
C15—C16—H16	119.5	C9—C10—H10	119.5
C17—C16—H16	119.5	C11—C10—H10	119.5
C18—C17—C16	117.94 (18)	C12—C11—C10	119.44 (18)
C18—C17—C20	120.96 (19)	C12—C11—H11	120.3
C16—C17—C20	121.1 (2)	C10—C11—H11	120.3
C17—C18—C19	121.60 (18)	C11—C12—N1	118.61 (17)

C17—C18—H18	119.2	C11—C12—C13	124.54 (17)
C19—C18—H18	119.2	N1—C12—C13	116.85 (16)
C14—C19—C18	119.74 (19)	C12—C13—H13A	109.5
C14—C19—H19	120.1	C12—C13—H13B	109.5
C18—C19—H19	120.1	H13A—C13—H13B	109.5
C17—C20—H20A	109.5	C12—C13—H13C	109.5
C17—C20—H20B	109.5	H13A—C13—H13C	109.5
H20A—C20—H20B	109.5	H13B—C13—H13C	109.5
C17—C20—H20C	109.5	C8—N1—C12	123.96 (15)
H20A—C20—H20C	109.5	C8—N1—H1A	118.4 (14)
H20B—C20—H20C	109.5	C12—N1—H1A	117.6 (14)
O4—S2—O6	112.84 (9)	C8—N2—H2A	115.9 (15)
O4—S2—O5	112.66 (8)	C8—N2—H2B	120.4 (15)
O6—S2—O5	111.06 (8)	H2A—N2—H2B	120 (2)
C6—C1—C2—C3	0.9 (3)	C15—C14—S2—O4	151.01 (16)
S1—C1—C2—C3	-178.58 (18)	C19—C14—S2—O4	-32.74 (18)
C1—C2—C3—C4	-0.5 (4)	C15—C14—S2—O6	-88.27 (17)
C2—C3—C4—C5	-0.4 (4)	C19—C14—S2—O6	87.97 (17)
C2—C3—C4—C7	-179.5 (2)	C15—C14—S2—O5	30.25 (18)
C3—C4—C5—C6	0.8 (3)	C19—C14—S2—O5	-153.50 (15)
C7—C4—C5—C6	179.9 (2)	N4—C21—C22—C23	-179.93 (18)
C2—C1—C6—C5	-0.5 (3)	N3—C21—C22—C23	1.0 (3)
S1—C1—C6—C5	178.96 (17)	C21—C22—C23—C24	-0.7 (3)
C4—C5—C6—C1	-0.3 (3)	C22—C23—C24—C25	0.0 (3)
C6—C1—S1—O3	-81.39 (18)	C23—C24—C25—N3	0.3 (3)
C2—C1—S1—O3	98.11 (17)	C23—C24—C25—C26	179.47 (17)
C6—C1—S1—O1	157.51 (16)	N4—C21—N3—C25	-179.79 (17)
C2—C1—S1—O1	-22.99 (19)	C22—C21—N3—C25	-0.7 (3)
C6—C1—S1—O2	38.15 (18)	C24—C25—N3—C21	0.0 (3)
C2—C1—S1—O2	-142.35 (17)	C26—C25—N3—C21	-179.21 (16)
C19—C14—C15—C16	-0.9 (3)	N2—C8—C9—C10	-179.67 (19)
S2—C14—C15—C16	175.40 (17)	N1—C8—C9—C10	-1.1 (3)
C14—C15—C16—C17	-0.3 (3)	C8—C9—C10—C11	0.6 (3)
C15—C16—C17—C18	1.0 (3)	C9—C10—C11—C12	-0.3 (3)
C15—C16—C17—C20	-178.9 (2)	C10—C11—C12—N1	0.4 (3)
C16—C17—C18—C19	-0.6 (3)	C10—C11—C12—C13	-178.58 (18)
C20—C17—C18—C19	179.3 (2)	N2—C8—N1—C12	179.90 (17)
C15—C14—C19—C18	1.2 (3)	C9—C8—N1—C12	1.3 (3)
S2—C14—C19—C18	-175.05 (16)	C11—C12—N1—C8	-0.9 (3)
C17—C18—C19—C14	-0.5 (3)	C13—C12—N1—C8	178.12 (16)

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1A $\cdots$ O2	0.90 (1)	1.88 (1)	2.772 (2)	171 (2)
N2—H2A $\cdots$ O3 <sup>i</sup>	0.87 (1)	2.01 (1)	2.880 (2)	174 (2)
N2—H2B $\cdots$ O1	0.88 (1)	2.07 (1)	2.919 (2)	162 (2)

---

N3—H3A···O5	0.89 (1)	1.90 (1)	2.789 (2)	174 (2)
N4—H4A···O4 <sup>ii</sup>	0.88 (1)	2.02 (1)	2.882 (2)	167 (2)
N4—H4B···O6	0.88 (1)	2.04 (1)	2.883 (2)	162 (2)
C22—H22···O5 <sup>ii</sup>	0.93	2.58	3.455 (2)	157

---

Symmetry codes: (i)  $x-1, y, z$ ; (ii)  $x+1, y, z$ .