

# Crystal structures of 1,4-diazabicyclo[2.2.2]octan-1-ium 4-nitrobenzoate dihydrate and 1,4-diazabicyclo[2.2.2]octane-1,4-diium bis(4-nitrobenzoate): the influence of solvent upon the stoichiometry of the formed salt


Aina Mardia Akhmad Aznan, Zanariah Abdullah\* and Edward R. T. Tiekink\*


Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia. \*Correspondence e-mail: zana@um.edu.my, edward.tiekink@gmail.com

The 1:1 co-crystallization of 1,4-diazabicyclo[2.2.2]octane (DABCO) with 4-nitrobenzoic acid in ethanol–water (3/1) gave the salt dihydrate  $C_6H_{13}N_2^+ \cdot C_7H_4NO_4^- \cdot 2H_2O$ , (1), whereas from methanol, the salt  $C_6H_{14}N_2^{2+} \cdot 2C_7H_4NO_4^-$ , (2), was isolated. In (1), the cation and anion are linked by a strong  $N-H \cdots O$  hydrogen bond, and the carboxylate anion is close to planar [dihedral angle between terminal residues =  $6.83(9)^\circ$ ]. In (2), a three-ion aggregate is assembled by two  $N-H \cdots O$  hydrogen bonds, and the carboxylate anions are again close to planar [dihedral angles between terminal residues =  $1.7(3)$  and  $5.9(3)^\circ$ ]. Through the intervention of solvent water molecules, which self-assemble into helical supramolecular chains along the  $b$  axis, the three-dimensional architecture in (1) is stabilized by water–DABCO  $O-H \cdots N$  and water–carboxylate  $O-H \cdots O$  hydrogen bonds, with additional stability afforded by  $C-H \cdots O$  interactions. The global crystal structure comprises alternating layers of water molecules and ion pairs stacked along the  $c$  axis. In the crystal of (2), the three-ion aggregates are assembled into a three-dimensional architecture by a large number of methylene–carboxylate/nitro  $C-H \cdots O$  interactions as well as  $\pi-\pi$  contacts between inversion-related benzene rings [inter-centroid distances =  $3.5644(16)$  and  $3.6527(16)$  Å]. The cations and anions assemble into alternating layers along the  $c$  axis.

## 1. Chemical context

The formation of co-crystals or salts is dependent on the difference in  $pK_a$  of the interacting species (Childs *et al.*, 2007). Thus, when the  $\Delta(pK_a)$  [=  $pK_a(\text{base}) - pK_a(\text{acid})$ ] value is greater than three, a salt is anticipated. In this context, it is not surprising that a search of the Cambridge Structural Database (CSD, version 53.5, last update November 2013; Allen, 2002) showed that nearly 90% of the 57 multi-component crystals, containing species derived from highly basic 1,4-diazabicyclo[2.2.2]octane (DABCO) and a carboxylic acid, contained at least a mono-protonated form of DABCO. It was in the context of on-going studies of co-crystallization experiments (Broker & Tiekink, 2007; Arman & Tiekink, 2013; Arman *et al.*, 2014) between nitrogen-containing molecules and carboxylic acids, that the title salts were isolated. The co-crystallization experiments yielding the title salts produced unexpected outcomes in that while (1) formed as a 1:1 salt dihydrate from the 1:1 co-crystallization of DABCO and 4-nitrobenzoic acid in ethanol/water (3/1) solution, a 1:2 salt (2) was isolated from the 1:1 co-crystallization of DABCO and 4-nitrobenzoic acid in methanol solution. The molecular and crystal structures of (1) and (2) are described herein.






## 2. Structural commentary

The asymmetric unit of (1) comprises a 1,4-diazabicyclo[2.2.2]octan-1-ium mono-cation, a 4-nitrobenzoate anion and two water molecules of hydration (Fig. 1). The most notable feature in the cation is the elongation of the N2—C bond lengths [1.4951 (16)–1.5007 (15) Å] compared to the N3—C bond lengths [1.4635 (17)–1.4773 (17) Å], consistent with protonation at the N2 atom. In the anion, the near equivalence of the C1—O1,O2 bond lengths of 1.2625 (15) and 1.2495 (16) Å, respectively, is again consistent with proton transfer; the longer bond involves atom O1 which forms a strong N—H···O hydrogen bond (Table 1). The dihedral angles between the central ring and the carboxylate and nitro groups are 8.68 (8) and 3.80 (5)°, respectively, and the dihedral angle between the terminal groups is 6.83 (9)°, consistent with an approximately planar molecule.

The asymmetric unit of (2) comprises a 1,4-diazabicyclo[2.2.2]octane-1,4-diium di-cation and two 4-nitrobenzoate anions (Fig. 2). In the dication, the N3—C [1.483 (3)–1.487 (3) Å] and N4—C [1.486 (3)–1.487 (3) Å] bond lengths are experimentally equivalent and consistent with diprotonation. In the anions, the disparity of the C1—O1, O2 bond lengths, *i.e.* 1.281 (3) and 1.228 (3) Å, is slightly greater than that in C8—O5, O6 of 1.273 (3) and 1.231 (3) Å, respectively. In each case the longer bond forms a strong N—H···O hydrogen bond (Table 2). In order to confirm the location of the acidic hydrogen atoms, an unrestrained refinement was conducted, see *Refinement* for details. While there was some elongation in the N—H bond lengths, unrestrained refinement confirmed protonation at both nitrogen atoms. In the O1-containing anion, the dihedral angles between the central ring and the carboxylate and nitro



**Figure 1**

The molecular structures of the four independent constituents of (1), with atom labelling. Displacement ellipsoids are drawn at the 50% probability level. Hydrogen bonds are shown as dashed lines (see Table 1 for details).

**Table 1**

Hydrogen-bond geometry (Å, °) for (1).

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N2—H2N···O1	0.89 (1)	1.76 (1)	2.6431 (14)	173 (1)
O1W—H1W···N3	0.86 (2)	1.95 (2)	2.7974 (15)	172 (2)
O1W—H2W···O2W <sup>i</sup>	0.86 (2)	1.91 (1)	2.7500 (15)	165 (2)
O2W—H3W···O1W	0.85 (2)	1.87 (2)	2.7218 (15)	180 (2)
O2W—H4W···O2 <sup>ii</sup>	0.86 (1)	1.87 (1)	2.7182 (15)	171 (2)
C10—H10A···O3 <sup>iii</sup>	0.99	2.49	3.4253 (17)	158
C12—H12A···O2W <sup>iv</sup>	0.99	2.49	3.3711 (16)	147
C12—H12B···O2 <sup>v</sup>	0.99	2.57	3.4818 (16)	153

Symmetry codes: (i)  $-x, y - \frac{1}{2}, -z + \frac{3}{2}$ ; (ii)  $x - 1, y + 1, z$ ; (iii)  $-x + 2, -y + 1, -z + 1$ ; (iv)  $x + 1, y - 1, z$ ; (v)  $x - 1, y, z$ .

**Table 2**

Hydrogen-bond geometry (Å, °) for (2).


<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N3—H3N···O1	0.88 (2)	1.66 (2)	2.539 (3)	173 (2)
N4—H4N···O5	0.89 (2)	1.65 (2)	2.542 (3)	175 (3)
C15—H15A···O6 <sup>i</sup>	0.99	2.42	3.193 (3)	134
C16—H16A···O4 <sup>ii</sup>	0.99	2.42	3.375 (3)	161
C17—H17A···O7 <sup>iii</sup>	0.99	2.42	3.338 (3)	153
C17—H17B···O6 <sup>i</sup>	0.99	2.42	3.313 (3)	149
C20—H20A···O2 <sup>iv</sup>	0.99	2.41	3.043 (3)	121
C20—H20B···O8 <sup>v</sup>	0.99	2.42	3.339 (3)	154

Symmetry codes: (i)  $-x + 2, -y + 1, -z + 1$ ; (ii)  $x + 1, y, z - 1$ ; (iii)  $x - 1, y, z + 1$ ; (iv)  $-x + 1, -y + 2, -z + 1$ ; (v)  $-x + 2, -y + 2, -z$ .

groups are 7.0 (3) and 8.7 (2)°, respectively, and the dihedral angle between the terminal groups is 1.7 (3)°. The comparable angles for the O5-containing anion are 2.2 (3), 7.4 (2) and 5.9 (3)°, respectively. As discussed below in *Supramolecular features*, the ions participate in strong N—H···O hydrogen bonds, forming a three-ion aggregate (Fig. 2) in which the dihedral angle between the benzene rings is 9.26 (14)°.


## 3. Supramolecular features

In (1), the cation and anion are linked by a strong N2—H···O1 hydrogen bond (Fig. 1 and Table 1). The two ion aggregates inter-digitate in columns aligned along the *b* axis. Adjacent columns stack along the *a* axis to form layers in the *ab* plane. The layers are interspersed by layers of water molecules which self-assemble into helical chains along the *b* axis, where each independent water molecule donates and accepts a water—O—H···O(water) hydrogen bond (Table 1). This leads to the formation of a three-dimensional structure (Fig. 3).



**Figure 2**


The molecular structures of the three independent constituents of (2), with atom labelling. Displacement ellipsoids are drawn at the 50% probability level. Hydrogen bonds are shown as dashed lines (see Table 2 for details).




**Figure 3**  
Unit-cell contents shown in projection down the *a* axis for (1). The O—H···O, O—H···N and C—H···O hydrogen bonds are shown as orange, blue and green dashed lines, respectively (see Table 1 for details).

Thus, links between layers are of the type water-O1W—H···N3 and water-OW2—H···O2(carboxylate). Additional stability to the supramolecular assembly is afforded by methylene-C—H···O2(carboxylate) and O2W(water) interactions; it is noteworthy that both of the former interactions involve hydrogen atoms derived from the same methylene-C12 atom (Table 1). A methylene-C—H···O3(nitro) interaction is also formed; the nitro-O4 atom does not form a significant interaction in this scenario. Although there is an alignment of benzene rings, the closest  $\pi$ — $\pi$  contact is 3.7376 (7) Å, occurring between centrosymmetrically related rings [symmetry operation:  $2 - x, 1 - y, 1 - z$ ].

In (2), the di-cation is linked to two anions *via* strong N—H···O hydrogen bonds (Fig. 4 and Table 2). Globally, the three ion aggregates assemble into layers in the *ab* plane that stack along the *c* axis. A large number of C—H···O interactions occur, remarkably featuring a narrow range of H···O



**Figure 4**  
Unit-cell contents shown in projection down the *a* axis for (2). The N—H···O and C—H···O hydrogen bonds are shown as blue and green dashed lines, respectively (see Table 2 for details).



**Figure 5**  
Unit-cell contents shown in projection down the *b* axis for (2). The N—H···O and C—H···O hydrogen bonds are shown as blue and green dashed lines, respectively (see Table 2 for details).

separations, *i.e.* 2.41–2.42 Å (Table 2). All interactions involve methylene-H atoms as donors. The carboxylate-O2 and O4 atoms and all nitro but O3 atoms are acceptors; both methylene-H atoms of methylene-C17 and C20 participate in C—H···O interactions. The result of these interactions is the formation of a three-dimensional architecture (Fig. 4). Additional stability to the supramolecular assembly is afforded by  $\pi$ — $\pi$  interactions between inversion-related rings, *i.e.* inter-centroid distances = 3.5644 (16) Å for interactions between the C2—C8 rings (symmetry code:  $-x + 1, -y + 2, -z + 2$ ) and 3.6527 (16) Å between C9—C14 rings (symmetry code:  $-x + 2, -y + 1, -z$ ). An alternate description of the global crystal packing is based on alternating of layers of cations and layers of anions along the *c* axis (Fig. 5).

#### 4. Database survey

As mentioned in the *Chemical context*, there are 57 species in the crystallographic literature containing DABCO or its mono- or diprotonated forms and a carboxylic acid or carboxylate anion. In fact, co-crystals are rare, being around 10% of all structures. Co-crystals are formed with several dicarboxylic acids where the functional groups are separated by long chains of over four carbon atoms (Braga *et al.*, 2003; Moon & Park, 2012), with phosphonoacetic acid (Bowes *et al.*, 2003) and with isophthalic acid (Marivel *et al.*, 2010). While the majority of the remaining structures contain species derived from a dicarboxylic acid, there are 13 examples of structures containing species derived from a mono-carboxylic acid which are more directly suitable for comparison with (1) and (2). Further, in each case the original carboxylic acid was connected to an aromatic ring. Of the sub-set of 13 structures,

**Table 3**  
Experimental details.

	(1)	(2)
Crystal data		
Chemical formula	$C_6H_{13}N_2^+ \cdot C_7H_4NO_4^- \cdot 2H_2O$	$C_6H_{14}N_2^{2+} \cdot 2C_7H_4NO_4^-$
$M_r$	315.33	446.42
Crystal system, space group	Monoclinic, $P2_1/c$	Triclinic, $P\bar{1}$
Temperature (K)	100	100
$a, b, c$ (Å)	6.5982 (1), 6.6074 (1), 34.4574 (6)	9.1036 (4), 9.5027 (3), 12.0736 (3)
$\alpha, \beta, \gamma$ (°)	90, 94.809 (1), 90	73.982 (3), 83.624 (3), 88.661 (3)
$V$ (Å <sup>3</sup> )	1496.95 (4)	997.68 (6)
$Z$	4	2
Radiation type	Cu $K\alpha$	Cu $K\alpha$
$\mu$ (mm <sup>-1</sup> )	0.94	0.99
Crystal size (mm)	0.30 × 0.30 × 0.20	0.40 × 0.40 × 0.20
Data collection		
Diffractometer	Agilent SuperNova Dual with an Atlas detector	Agilent SuperNova Dual with an Atlas detector
Absorption correction	Multi-scan ( <i>CrysAlis PRO</i> ; Agilent, 2013)	Multi-scan ( <i>CrysAlis PRO</i> ; Agilent, 2013)
$T_{min}, T_{max}$	0.888, 1.000	0.991, 1.000
No. of measured, independent and observed [ $I > 2\sigma(I)$ ] reflections	11203, 3072, 2949	17849, 4101, 3775
$R_{int}$	0.015	0.046
$(\sin \theta/\lambda)_{max}$ (Å <sup>-1</sup> )	0.626	0.626
Refinement		
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.039, 0.099, 1.04	0.074, 0.229, 1.12
No. of reflections	3072	4101
No. of parameters	215	295
No. of restraints	7	2
H-atom treatment	H atoms treated by a mixture of independent and constrained refinement	H atoms treated by a mixture of independent and constrained refinement
$\Delta\rho_{max}, \Delta\rho_{min}$ (e Å <sup>-3</sup> )	0.40, -0.30	0.60, -0.58

Computer programs: *CrysAlis PRO* (Agilent, 2013), *SHELXS97* and *SHELXL97* (Sheldrick, 2008), *ORTEP-3 for Windows* (Farrugia, 2012), *DIAMOND* (Brandenburg, 2006) and *publCIF* (Westrip, 2010).

three are similar to (1), having the mono-protonated form of DABCO. The carboxylate counter-ions are 2,4-dinitrobenzoate (Rosli *et al.*, 2006), 3,5-dihydroxybenzoate (Burchell *et al.*, 2001a) and 6-hydroxy-2-naphthoate (Jacobs *et al.*, 2010); the latter two structures were characterized as mono- and sesqui-hydrates, respectively. Analogues of (2) were found in seven examples, namely in both polymorphs of benzoate, and in 2-hydroxybenzoate and 2-acetoxybenzoate (Skovsgaard & Bond, 2009), 2-chlorobenzoate (Skovsgaard & Bond, 2008), 2-hydroxybenzoate (Skovsgaard & Bond, 2008), and in polymorphic hydrates of 3,5-dinitrobenzoate (Burchell *et al.*, 2001b; Chantrapromma & Fun, 2004). Finally, there are three intriguing examples where a mono-protonated DABCO cation is present along with a carboxylate anion and the neutral form of the original carboxylic acid. These contain the following carboxylic acids: 1-hydroxy-2-naphthoic acid and 3-hydroxy-2-naphthoic acid (Jacobs *et al.*, 2010) and 2-aminobenzoic acid (Arman *et al.*, 2011). In light of the foregoing structural diversity, in retrospect perhaps it is not so surprising that solvent can influence product formation, especially when water is involved.

### 5. Synthesis and crystallization

1,4-Diazabicyclo[2.2.2]octane (Merck; 0.10 g, 0.0009 mol), was mixed with 4-nitrobenzoic acid (Merck; 0.15 g, 0.0009 mol) in

a solution containing ethanol (30 ml) and water (10 ml). The solution was heated for 2 h at 350 K. The mixture was then left for slow evaporation and colourless crystals of (1) formed after 4 days. In a similar experiment, 1,4-diazabicyclo[2.2.2]octane (0.298 g, 0.00265 mol) was mixed with 4-nitrobenzoic acid (0.444 g, 0.00265 mol) in a solution of methanol (50 ml). The solution was heated for 2 h at 345 K. The mixture was then left for slow evaporation and colourless crystals of (2) formed after 4 days.

### 6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 3. Carbon-bound H-atoms were placed in calculated positions ( $C-H = 0.95-0.99$  Å) and were included in the refinement in the riding-model approximation, with  $U_{iso}(H) = 1.2U_{eq}(C)$ . The N-bound H-atoms were located in a difference Fourier map but were refined with a distance restraint:  $N-H = 0.88$  (1) Å with  $U_{iso}(H) = 1.2U_{eq}(N)$ . For (1), the water-bound H atoms were refined with distance restraints:  $O-H = 0.84$  (1) and  $H \cdots H = 1.39$  (2) Å with  $U_{iso}(H) = 1.5U_{eq}(O)$ . For (2), the maximum and minimum residual electron density peaks of 0.60 and 0.58 e Å<sup>-3</sup>, respectively, were located 0.81 and 0.10 Å from atoms O5 and H4N, respectively. In order to confirm the location of the N-bound H atoms, in a separate refinement these were refined

without distance restraints. For (1), the N2–H2N bond length was 0.948 (17) Å. For (2), the N3–H3N and N4–H4N bond lengths were 0.93 (4) and 1.08 (3) Å, respectively. In the refinement of (1), one reflection, *i.e.* (180), was omitted from the refinement owing to poor agreement. For the same reasons, the following reflections were omitted from the final refinement of (2): (550), ( $\bar{2}06$ ), (136), (139), (2410), (331), (224) and (662).

### Acknowledgements

Intensity data were provided by the University of Malaya Crystallographic Laboratory. This research is supported by the University of Malaya Research Grant Scheme (RG125/AFC10R).

### References

- Agilent (2013). *CrysAlis PRO*. Agilent Technologies Inc., Santa Clara, CA, USA.
- Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
- Arman, H. D., Kaulgud, T., Miller, T. & Tiekink, E. R. T. (2014). *Z. Kristallogr.* **229**, 295–302.
- Arman, H. D., Kaulgud, T. & Tiekink, E. R. T. (2011). *Acta Cryst.* **E67**, o2933.
- Arman, H. D. & Tiekink, E. R. T. (2013). *Z. Kristallogr.* **228**, 289–294.
- Bowes, K. F., Ferguson, G., Lough, A. J., Zakaria, C. M. & Glidewell, C. (2003). *Acta Cryst.* **B59**, 87–99.
- Braga, D., Maini, L., de Sanctis, G., Rubini, K., Grepioni, F., Chierotti, M. R. & Gobetto, R. (2003). *Chem. Eur. J.* **9**, 5538–5548.
- Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Broker, G. A. & Tiekink, E. R. T. (2007). *CrystEngComm*, **9**, 1096–1109.
- Burchell, C. J., Ferguson, G., Lough, A. J., Gregson, R. M. & Glidewell, C. (2001a). *Acta Cryst.* **B57**, 329–338.
- Burchell, C. J., Glidewell, C., Lough, A. J. & Ferguson, G. (2001b). *Acta Cryst.* **B57**, 201–212.
- Chantrapromma, S. & Fun, H.-K. (2004). *Acta Cryst.* **E60**, o1250–o1252.
- Childs, S. L., Stahly, G. P. & Park, A. (2007). *Mol. Pharm.* **4**, 323–338.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Jacobs, A., Nassimbeni, L. R., Ramon, G. & Sebogisi, B. K. (2010). *CrystEngComm*, **12**, 3065–3070.
- Marivel, S., Braga, D., Grepioni, F. & Lampronti, G. I. (2010). *CrystEngComm*, **12**, 2107–2112.
- Moon, S.-H. & Park, K.-M. (2012). *Acta Cryst.* **E68**, o2344.
- Rosli, M. M., Fun, H.-K., Lee, B. S. & Chantrapromma, S. (2006). *Acta Cryst.* **E62**, o4575–o4577.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Skovsgaard, S. & Bond, A. D. (2008). *Acta Cryst.* **E64**, o1416.
- Skovsgaard, S. & Bond, A. D. (2009). *CrystEngComm*, **11**, 444–453.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

## supporting information

*Acta Cryst.* (2014). E70, 31-35 [https://doi.org/10.1107/S1600536814011532]

## Crystal structures of 1,4-diazabicyclo[2.2.2]octan-1-ium 4-nitrobenzoate dihydrate and 1,4-diazabicyclo[2.2.2]octane-1,4-dium bis(4-nitrobenzoate): the influence of solvent upon the stoichiometry of the formed salt

Aina Mardia Akhmad Aznan, Zanariah Abdullah and Edward R. T. Tiekink

### Computing details

For both compounds, data collection: *CrysAlis PRO* (Agilent, 2013); cell refinement: *CrysAlis PRO* (Agilent, 2013); data reduction: *CrysAlis PRO* (Agilent, 2013); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).

### (1) 1,4-Diazabicyclo[2.2.2]octan-1-ium 4-nitrobenzoate dihydrate

#### Crystal data

$C_6H_{13}N_2^+ \cdot C_7H_4NO_4^- \cdot 2H_2O$

$M_r = 315.33$

Monoclinic,  $P2_1/c$

Hall symbol: -P 2ybc

$a = 6.5982$  (1) Å

$b = 6.6074$  (1) Å

$c = 34.4574$  (6) Å

$\beta = 94.809$  (1)°

$V = 1496.95$  (4) Å<sup>3</sup>

$Z = 4$

$F(000) = 672$

$D_x = 1.399$  Mg m<sup>-3</sup>

Cu  $K\alpha$  radiation,  $\lambda = 1.54184$  Å

Cell parameters from 7046 reflections

$\theta = 3.9$ – $76.3$ °

$\mu = 0.94$  mm<sup>-1</sup>

$T = 100$  K

Prism, colourless

$0.30 \times 0.30 \times 0.20$  mm

#### Data collection

Agilent SuperNova Dual

diffractometer with an Atlas detector

Radiation source: SuperNova (Cu) X-ray

Source

Mirror monochromator

Detector resolution: 10.4041 pixels mm<sup>-1</sup>

$\omega$  scan

Absorption correction: multi-scan

(*CrysAlis PRO*; Agilent, 2013)

$T_{\min} = 0.888$ ,  $T_{\max} = 1.000$

11203 measured reflections

3072 independent reflections

2949 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.015$

$\theta_{\max} = 75.0$ °,  $\theta_{\min} = 5.2$ °

$h = -8 \rightarrow 8$

$k = -8 \rightarrow 5$

$l = -43 \rightarrow 42$

#### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.039$

$wR(F^2) = 0.099$

$S = 1.04$

3072 reflections

215 parameters

7 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent

and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0467P)^2 + 0.8811P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.40 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.30 \text{ e } \text{\AA}^{-3}$$

Extinction correction: SHELXL97 (Sheldrick, 2008),  $F_c^* = kFc[1 + 0.001x Fc^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0076 (6)

### Special details

**Geometry.** All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.63866 (13)	0.34327 (15)	0.59514 (3)	0.0233 (2)
O2	0.94425 (14)	0.33794 (16)	0.62902 (3)	0.0261 (2)
O3	1.11387 (14)	0.20664 (15)	0.42606 (3)	0.0234 (2)
O4	1.40402 (14)	0.17440 (16)	0.45978 (3)	0.0262 (2)
N1	1.21908 (16)	0.20014 (15)	0.45723 (3)	0.0175 (2)
C1	0.82971 (19)	0.32425 (18)	0.59836 (4)	0.0178 (3)
C2	0.93105 (18)	0.28490 (18)	0.56096 (3)	0.0153 (2)
C3	1.13841 (18)	0.24078 (18)	0.56251 (3)	0.0162 (3)
H3	1.2149	0.2318	0.5871	0.019*
C4	1.23420 (18)	0.20989 (18)	0.52871 (4)	0.0163 (2)
H4	1.3751	0.1786	0.5297	0.020*
C5	1.11798 (18)	0.22606 (17)	0.49335 (3)	0.0149 (2)
C6	0.91145 (18)	0.26736 (17)	0.49062 (3)	0.0155 (2)
H6	0.8354	0.2752	0.4660	0.019*
C7	0.81863 (17)	0.29702 (18)	0.52492 (3)	0.0156 (2)
H7	0.6772	0.3258	0.5238	0.019*
N2	0.48292 (15)	0.59677 (17)	0.64400 (3)	0.0178 (2)
H2N	0.535 (2)	0.503 (2)	0.6291 (4)	0.021*
N3	0.33337 (16)	0.86385 (17)	0.68589 (3)	0.0207 (2)
C8	0.31552 (19)	0.7058 (2)	0.62050 (4)	0.0213 (3)
H8A	0.2168	0.6074	0.6082	0.026*
H8B	0.3720	0.7863	0.5997	0.026*
C9	0.2099 (2)	0.8456 (2)	0.64826 (4)	0.0309 (3)
H9A	0.1903	0.9811	0.6363	0.037*
H9B	0.0744	0.7900	0.6527	0.037*
C10	0.64491 (19)	0.7447 (2)	0.65768 (4)	0.0212 (3)
H10A	0.7121	0.7988	0.6352	0.025*
H10B	0.7491	0.6777	0.6756	0.025*
C11	0.5423 (2)	0.9174 (2)	0.67881 (5)	0.0325 (3)
H11A	0.6212	0.9456	0.7039	0.039*

H11B	0.5416	1.0418	0.6628	0.039*
C12	0.39871 (18)	0.49677 (19)	0.67830 (4)	0.0199 (3)
H12A	0.5038	0.4108	0.6923	0.024*
H12B	0.2810	0.4105	0.6695	0.024*
C13	0.3323 (2)	0.6654 (2)	0.70520 (4)	0.0257 (3)
H13A	0.1938	0.6361	0.7127	0.031*
H13B	0.4257	0.6688	0.7292	0.031*
O1W	0.19593 (16)	1.12932 (16)	0.74123 (3)	0.0268 (2)
H1W	0.243 (3)	1.058 (3)	0.7232 (4)	0.040*
H2W	0.151 (3)	1.039 (2)	0.7565 (4)	0.040*
O2W	-0.12706 (15)	1.33213 (17)	0.70557 (3)	0.0289 (2)
H3W	-0.026 (2)	1.268 (3)	0.7167 (5)	0.043*
H4W	-0.117 (3)	1.328 (3)	0.6810 (3)	0.043*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0175 (4)	0.0299 (5)	0.0231 (5)	0.0006 (4)	0.0055 (3)	-0.0078 (4)
O2	0.0235 (5)	0.0391 (6)	0.0158 (4)	0.0014 (4)	0.0025 (4)	-0.0027 (4)
O3	0.0271 (5)	0.0281 (5)	0.0153 (4)	0.0000 (4)	0.0033 (4)	0.0001 (4)
O4	0.0181 (5)	0.0355 (6)	0.0262 (5)	0.0027 (4)	0.0087 (4)	0.0009 (4)
N1	0.0200 (5)	0.0147 (5)	0.0185 (5)	-0.0011 (4)	0.0058 (4)	0.0004 (4)
C1	0.0190 (6)	0.0167 (6)	0.0181 (6)	-0.0009 (5)	0.0042 (5)	-0.0018 (5)
C2	0.0170 (6)	0.0122 (5)	0.0171 (6)	-0.0017 (4)	0.0034 (4)	-0.0006 (4)
C3	0.0165 (6)	0.0154 (6)	0.0165 (6)	-0.0012 (4)	-0.0004 (4)	0.0003 (4)
C4	0.0137 (5)	0.0144 (5)	0.0209 (6)	-0.0006 (4)	0.0026 (4)	0.0007 (4)
C5	0.0178 (6)	0.0116 (5)	0.0159 (6)	-0.0020 (4)	0.0053 (4)	-0.0006 (4)
C6	0.0174 (6)	0.0124 (5)	0.0164 (5)	-0.0022 (4)	-0.0002 (4)	0.0002 (4)
C7	0.0135 (5)	0.0133 (5)	0.0200 (6)	-0.0016 (4)	0.0021 (4)	-0.0005 (4)
N2	0.0148 (5)	0.0219 (5)	0.0171 (5)	0.0002 (4)	0.0033 (4)	-0.0034 (4)
N3	0.0212 (5)	0.0228 (6)	0.0187 (5)	0.0016 (4)	0.0044 (4)	-0.0023 (4)
C8	0.0200 (6)	0.0270 (7)	0.0167 (6)	0.0018 (5)	0.0005 (5)	-0.0011 (5)
C9	0.0331 (7)	0.0375 (8)	0.0217 (7)	0.0148 (6)	-0.0002 (6)	-0.0017 (6)
C10	0.0153 (6)	0.0274 (7)	0.0215 (6)	-0.0047 (5)	0.0041 (5)	-0.0018 (5)
C11	0.0276 (7)	0.0300 (7)	0.0416 (8)	-0.0100 (6)	0.0126 (6)	-0.0138 (7)
C12	0.0174 (6)	0.0207 (6)	0.0219 (6)	-0.0009 (5)	0.0039 (5)	0.0027 (5)
C13	0.0299 (7)	0.0274 (7)	0.0210 (6)	0.0011 (5)	0.0104 (5)	0.0017 (5)
O1W	0.0311 (5)	0.0285 (5)	0.0210 (5)	0.0053 (4)	0.0026 (4)	-0.0024 (4)
O2W	0.0258 (5)	0.0420 (6)	0.0191 (5)	0.0084 (4)	0.0041 (4)	0.0005 (4)

*Geometric parameters (Å, °)*

O1—C1	1.2625 (15)	N3—C13	1.4706 (17)
O2—C1	1.2495 (16)	N3—C9	1.4773 (17)
O3—N1	1.2298 (14)	C8—C9	1.5381 (18)
O4—N1	1.2279 (14)	C8—H8A	0.9900
N1—C5	1.4704 (15)	C8—H8B	0.9900
C1—C2	1.5231 (16)	C9—H9A	0.9900



C2—C7	1.3943 (17)	C9—H9B	0.9900
C2—C3	1.3956 (16)	C10—C11	1.5408 (19)
C3—C4	1.3857 (17)	C10—H10A	0.9900
C3—H3	0.9500	C10—H10B	0.9900
C4—C5	1.3884 (17)	C11—H11A	0.9900
C4—H4	0.9500	C11—H11B	0.9900
C5—C6	1.3852 (17)	C12—C13	1.5365 (18)
C6—C7	1.3900 (16)	C12—H12A	0.9900
C6—H6	0.9500	C12—H12B	0.9900
C7—H7	0.9500	C13—H13A	0.9900
N2—C10	1.4951 (16)	C13—H13B	0.9900
N2—C8	1.4982 (16)	O1W—H1W	0.861 (9)
N2—C12	1.5007 (15)	O1W—H2W	0.866 (9)
N2—H2N	0.892 (9)	O2W—H3W	0.852 (9)
N3—C11	1.4635 (17)	O2W—H4W	0.856 (9)
O4—N1—O3	123.50 (10)	N2—C8—H8B	110.2
O4—N1—C5	118.32 (10)	C9—C8—H8B	110.2
O3—N1—C5	118.18 (10)	H8A—C8—H8B	108.5
O2—C1—O1	126.58 (11)	N3—C9—C8	110.47 (11)
O2—C1—C2	116.76 (11)	N3—C9—H9A	109.6
O1—C1—C2	116.65 (11)	C8—C9—H9A	109.6
C7—C2—C3	119.51 (11)	N3—C9—H9B	109.6
C7—C2—C1	120.35 (11)	C8—C9—H9B	109.6
C3—C2—C1	120.13 (11)	H9A—C9—H9B	108.1
C4—C3—C2	120.90 (11)	N2—C10—C11	107.60 (10)
C4—C3—H3	119.6	N2—C10—H10A	110.2
C2—C3—H3	119.6	C11—C10—H10A	110.2
C3—C4—C5	117.96 (11)	N2—C10—H10B	110.2
C3—C4—H4	121.0	C11—C10—H10B	110.2
C5—C4—H4	121.0	H10A—C10—H10B	108.5
C6—C5—C4	122.86 (11)	N3—C11—C10	110.91 (11)
C6—C5—N1	118.62 (11)	N3—C11—H11A	109.5
C4—C5—N1	118.52 (10)	C10—C11—H11A	109.5
C5—C6—C7	118.10 (11)	N3—C11—H11B	109.5
C5—C6—H6	120.9	C10—C11—H11B	109.5
C7—C6—H6	120.9	H11A—C11—H11B	108.0
C6—C7—C2	120.66 (11)	N2—C12—C13	107.39 (10)
C6—C7—H7	119.7	N2—C12—H12A	110.2
C2—C7—H7	119.7	C13—C12—H12A	110.2
C10—N2—C8	109.35 (10)	N2—C12—H12B	110.2
C10—N2—C12	109.99 (9)	C13—C12—H12B	110.2
C8—N2—C12	109.42 (9)	H12A—C12—H12B	108.5
C10—N2—H2N	109.7 (10)	N3—C13—C12	111.17 (10)
C8—N2—H2N	109.0 (10)	N3—C13—H13A	109.4
C12—N2—H2N	109.4 (11)	C12—C13—H13A	109.4
C11—N3—C13	109.35 (11)	N3—C13—H13B	109.4
C11—N3—C9	109.36 (11)	C12—C13—H13B	109.4

C13—N3—C9	107.53 (11)	H13A—C13—H13B	108.0
N2—C8—C9	107.72 (10)	H1W—O1W—H2W	103.0 (15)
N2—C8—H8A	110.2	H3W—O2W—H4W	107.8 (16)
C9—C8—H8A	110.2		
O2—C1—C2—C7	-170.56 (11)	C1—C2—C7—C6	178.03 (11)
O1—C1—C2—C7	8.23 (17)	C10—N2—C8—C9	-67.76 (13)
O2—C1—C2—C3	8.02 (17)	C12—N2—C8—C9	52.77 (14)
O1—C1—C2—C3	-173.19 (11)	C11—N3—C9—C8	50.98 (16)
C7—C2—C3—C4	0.32 (18)	C13—N3—C9—C8	-67.65 (14)
C1—C2—C3—C4	-178.27 (11)	N2—C8—C9—N3	13.17 (16)
C2—C3—C4—C5	0.59 (18)	C8—N2—C10—C11	53.58 (13)
C3—C4—C5—C6	-1.34 (18)	C12—N2—C10—C11	-66.60 (13)
C3—C4—C5—N1	178.05 (10)	C13—N3—C11—C10	51.74 (15)
O4—N1—C5—C6	176.15 (11)	C9—N3—C11—C10	-65.75 (15)
O3—N1—C5—C6	-3.43 (16)	N2—C10—C11—N3	11.63 (16)
O4—N1—C5—C4	-3.27 (16)	C10—N2—C12—C13	53.46 (13)
O3—N1—C5—C4	177.15 (10)	C8—N2—C12—C13	-66.68 (13)
C4—C5—C6—C7	1.12 (18)	C11—N3—C13—C12	-65.51 (14)
N1—C5—C6—C7	-178.28 (10)	C9—N3—C13—C12	53.13 (14)
C5—C6—C7—C2	-0.14 (17)	N2—C12—C13—N3	11.20 (14)
C3—C2—C7—C6	-0.56 (18)		

## Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N2—H2N...O1	0.89 (1)	1.76 (1)	2.6431 (14)	173 (1)
O1W—H1W...N3	0.86 (2)	1.95 (2)	2.7974 (15)	172 (2)
O1W—H2W...O2W <sup>iv</sup>	0.86 (2)	1.91 (1)	2.7500 (15)	165 (2)
O2W—H3W...O1W	0.85 (2)	1.87 (2)	2.7218 (15)	180 (2)
O2W—H4W...O2 <sup>ii</sup>	0.86 (1)	1.87 (1)	2.7182 (15)	171 (2)
C10—H10A...O3 <sup>iii</sup>	0.99	2.49	3.4253 (17)	158
C12—H12A...O2W <sup>iv</sup>	0.99	2.49	3.3711 (16)	147
C12—H12B...O2 <sup>v</sup>	0.99	2.57	3.4818 (16)	153

Symmetry codes: (i)  $-x, y-1/2, -z+3/2$ ; (ii)  $x-1, y+1, z$ ; (iii)  $-x+2, -y+1, -z+1$ ; (iv)  $x+1, y-1, z$ ; (v)  $x-1, y, z$ .

## (2) 1,4-Diazabicyclo[2.2.2]octane-1,4-dium bis(4-nitrobenzoate)

## Crystal data

 $C_6H_{14}N_2^{2+} \cdot 2C_7H_4NO_4^-$  $M_r = 446.42$ Triclinic,  $P\bar{1}$ Hall symbol:  $-P 1$  $a = 9.1036$  (4) Å $b = 9.5027$  (3) Å $c = 12.0736$  (3) Å $\alpha = 73.982$  (3)° $\beta = 83.624$  (3)° $\gamma = 88.661$  (3)° $V = 997.68$  (6) Å<sup>3</sup> $Z = 2$  $F(000) = 468$  $D_x = 1.486$  Mg m<sup>-3</sup>Cu  $K\alpha$  radiation,  $\lambda = 1.54184$  Å

Cell parameters from 10186 reflections

 $\theta = 3.8$ – $76.8$ ° $\mu = 0.99$  mm<sup>-1</sup> $T = 100$  K

Prism, colourless

 $0.40 \times 0.40 \times 0.20$  mm

*Data collection*

Agilent SuperNova Dual  
 diffractometer with an Atlas detector  
 Radiation source: SuperNova (Cu) X-ray  
 Source  
 Mirror monochromator  
 Detector resolution: 10.4041 pixels mm<sup>-1</sup>  
 $\omega$  scan  
 Absorption correction: multi-scan  
 (CrysAlis PRO; Agilent, 2013)

$T_{\min} = 0.991$ ,  $T_{\max} = 1.000$   
 17849 measured reflections  
 4101 independent reflections  
 3775 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.046$   
 $\theta_{\max} = 75.0^\circ$ ,  $\theta_{\min} = 3.8^\circ$   
 $h = -11 \rightarrow 11$   
 $k = -11 \rightarrow 11$   
 $l = -15 \rightarrow 15$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.074$   
 $wR(F^2) = 0.229$   
 $S = 1.12$   
 4101 reflections  
 295 parameters  
 2 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H atoms treated by a mixture of independent  
 and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.1192P)^2 + 1.3867P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.60 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.58 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.6213 (2)	0.7377 (2)	0.82692 (16)	0.0320 (5)
O2	0.4577 (2)	0.9073 (2)	0.75460 (15)	0.0301 (4)
O3	0.3598 (2)	0.7520 (2)	1.38056 (16)	0.0365 (5)
O4	0.2049 (2)	0.9218 (2)	1.31087 (17)	0.0343 (5)
N1	0.3033 (3)	0.8347 (2)	1.29939 (18)	0.0278 (5)
C1	0.5159 (3)	0.8269 (3)	0.8358 (2)	0.0230 (5)
C2	0.4601 (3)	0.8288 (3)	0.9588 (2)	0.0225 (5)
C3	0.3361 (3)	0.9096 (3)	0.9783 (2)	0.0239 (5)
H3A	0.2873	0.9646	0.9148	0.029*
C4	0.2829 (3)	0.9105 (3)	1.0906 (2)	0.0245 (5)
H4A	0.1976	0.9649	1.1051	0.029*
C5	0.3578 (3)	0.8300 (3)	1.1807 (2)	0.0244 (5)
C6	0.4829 (3)	0.7496 (3)	1.1642 (2)	0.0269 (5)
H6	0.5320	0.6958	1.2280	0.032*
C7	0.5343 (3)	0.7498 (3)	1.0515 (2)	0.0250 (5)

H7	0.6202	0.6961	1.0374	0.030*
O5	0.8758 (2)	0.7357 (2)	0.20757 (15)	0.0345 (5)
O6	1.0354 (2)	0.5601 (2)	0.27470 (17)	0.0385 (5)
O7	1.3135 (2)	0.6129 (2)	-0.29233 (18)	0.0368 (5)
O8	1.1731 (2)	0.7994 (2)	-0.35307 (16)	0.0347 (5)
N2	1.2191 (2)	0.7014 (3)	-0.27572 (19)	0.0287 (5)
C8	0.9814 (3)	0.6492 (3)	0.1953 (2)	0.0256 (5)
C9	1.0424 (3)	0.6604 (3)	0.0705 (2)	0.0233 (5)
C10	1.1613 (3)	0.5752 (3)	0.0471 (2)	0.0253 (5)
H10	1.2030	0.5075	0.1090	0.030*
C11	1.2196 (3)	0.5889 (3)	-0.0670 (2)	0.0264 (5)
H11	1.3010	0.5311	-0.0841	0.032*
C12	1.1562 (3)	0.6885 (3)	-0.1550 (2)	0.0252 (5)
C13	1.0376 (3)	0.7750 (3)	-0.1345 (2)	0.0259 (5)
H13	0.9963	0.8426	-0.1967	0.031*
C14	0.9806 (3)	0.7597 (3)	-0.0201 (2)	0.0255 (5)
H14	0.8989	0.8174	-0.0034	0.031*
N3	0.6982 (2)	0.7378 (2)	0.61824 (18)	0.0254 (5)
H3N	0.665 (3)	0.734 (4)	0.6902 (12)	0.030*
N4	0.7971 (2)	0.7223 (2)	0.41892 (18)	0.0261 (5)
H4N	0.830 (3)	0.728 (4)	0.3454 (12)	0.031*
C15	0.8607 (3)	0.7624 (3)	0.6003 (2)	0.0274 (5)
H15A	0.9072	0.6984	0.6663	0.033*
H15B	0.8832	0.8655	0.5954	0.033*
C16	0.9224 (3)	0.7273 (3)	0.4870 (2)	0.0310 (6)
H16A	0.9946	0.8035	0.4418	0.037*
H16B	0.9733	0.6317	0.5047	0.037*
C17	0.6672 (3)	0.5903 (3)	0.6066 (2)	0.0269 (5)
H17A	0.5616	0.5648	0.6317	0.032*
H17B	0.7276	0.5163	0.6564	0.032*
C18	0.7046 (3)	0.5908 (3)	0.4791 (2)	0.0299 (6)
H18A	0.7592	0.5009	0.4744	0.036*
H18B	0.6127	0.5935	0.4418	0.036*
C19	0.6263 (3)	0.8510 (3)	0.5300 (2)	0.0304 (6)
H19A	0.6328	0.9477	0.5453	0.036*
H19B	0.5206	0.8261	0.5334	0.036*
C20	0.7063 (3)	0.8562 (3)	0.4098 (2)	0.0318 (6)
H20A	0.6330	0.8614	0.3541	0.038*
H20B	0.7704	0.9444	0.3814	0.038*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0368 (10)	0.0397 (11)	0.0185 (9)	0.0127 (8)	-0.0025 (7)	-0.0077 (7)
O2	0.0355 (10)	0.0357 (10)	0.0175 (8)	0.0069 (8)	-0.0028 (7)	-0.0053 (7)
O3	0.0476 (12)	0.0394 (11)	0.0182 (9)	0.0023 (9)	-0.0009 (8)	-0.0022 (8)
O4	0.0375 (11)	0.0371 (10)	0.0264 (9)	0.0021 (8)	0.0072 (8)	-0.0096 (8)
N1	0.0331 (12)	0.0299 (11)	0.0189 (10)	-0.0031 (9)	0.0022 (8)	-0.0060 (8)

C1	0.0243 (11)	0.0254 (12)	0.0183 (11)	-0.0019 (9)	-0.0008 (9)	-0.0047 (9)
C2	0.0255 (12)	0.0223 (11)	0.0190 (11)	-0.0027 (9)	-0.0012 (9)	-0.0045 (9)
C3	0.0255 (12)	0.0255 (12)	0.0198 (11)	-0.0014 (9)	-0.0041 (9)	-0.0041 (9)
C4	0.0249 (12)	0.0250 (12)	0.0229 (12)	-0.0006 (9)	0.0010 (9)	-0.0066 (9)
C5	0.0300 (12)	0.0256 (12)	0.0162 (11)	-0.0037 (10)	0.0026 (9)	-0.0050 (9)
C6	0.0317 (13)	0.0275 (12)	0.0187 (11)	0.0016 (10)	-0.0043 (10)	-0.0013 (9)
C7	0.0257 (12)	0.0250 (12)	0.0222 (12)	0.0019 (9)	0.0003 (9)	-0.0040 (9)
O5	0.0401 (11)	0.0445 (11)	0.0163 (9)	0.0144 (9)	-0.0005 (7)	-0.0062 (8)
O6	0.0454 (12)	0.0458 (12)	0.0193 (9)	0.0158 (9)	-0.0035 (8)	-0.0019 (8)
O7	0.0349 (10)	0.0484 (12)	0.0294 (10)	0.0019 (9)	0.0053 (8)	-0.0180 (9)
O8	0.0352 (10)	0.0478 (12)	0.0194 (9)	-0.0030 (9)	-0.0015 (7)	-0.0068 (8)
N2	0.0256 (11)	0.0382 (12)	0.0230 (11)	-0.0036 (9)	0.0010 (8)	-0.0105 (9)
C8	0.0264 (12)	0.0291 (12)	0.0192 (12)	0.0002 (10)	-0.0015 (9)	-0.0037 (9)
C9	0.0247 (12)	0.0265 (12)	0.0184 (11)	-0.0009 (9)	-0.0018 (9)	-0.0057 (9)
C10	0.0258 (12)	0.0245 (12)	0.0235 (12)	-0.0003 (9)	-0.0039 (10)	-0.0026 (9)
C11	0.0247 (12)	0.0264 (12)	0.0275 (13)	0.0005 (9)	0.0013 (10)	-0.0080 (10)
C12	0.0255 (12)	0.0298 (12)	0.0204 (12)	-0.0048 (10)	0.0015 (9)	-0.0081 (10)
C13	0.0272 (12)	0.0312 (13)	0.0180 (11)	-0.0008 (10)	-0.0035 (9)	-0.0043 (9)
C14	0.0243 (12)	0.0294 (12)	0.0215 (12)	0.0036 (9)	-0.0013 (9)	-0.0056 (10)
N3	0.0270 (11)	0.0308 (11)	0.0176 (10)	0.0066 (9)	-0.0021 (8)	-0.0061 (8)
N4	0.0290 (11)	0.0304 (11)	0.0169 (10)	0.0033 (9)	0.0004 (8)	-0.0045 (8)
C15	0.0289 (13)	0.0309 (13)	0.0228 (12)	0.0008 (10)	-0.0052 (10)	-0.0072 (10)
C16	0.0258 (12)	0.0425 (15)	0.0232 (12)	0.0004 (11)	0.0003 (10)	-0.0075 (11)
C17	0.0280 (12)	0.0312 (13)	0.0188 (12)	-0.0005 (10)	-0.0002 (9)	-0.0033 (9)
C18	0.0365 (14)	0.0306 (13)	0.0217 (12)	-0.0009 (11)	-0.0018 (10)	-0.0062 (10)
C19	0.0341 (14)	0.0339 (14)	0.0214 (12)	0.0121 (11)	-0.0040 (10)	-0.0056 (10)
C20	0.0408 (15)	0.0317 (13)	0.0195 (12)	0.0090 (11)	-0.0038 (10)	-0.0018 (10)

*Geometric parameters (Å, °)*

O1—C1	1.281 (3)	C12—C13	1.384 (4)
O2—C1	1.228 (3)	C13—C14	1.390 (3)
O3—N1	1.229 (3)	C13—H13	0.9500
O4—N1	1.228 (3)	C14—H14	0.9500
N1—C5	1.475 (3)	N3—C17	1.483 (3)
C1—C2	1.520 (3)	N3—C15	1.485 (3)
C2—C3	1.386 (3)	N3—C19	1.487 (3)
C2—C7	1.398 (3)	N3—H3N	0.879 (10)
C3—C4	1.390 (3)	N4—C20	1.486 (3)
C3—H3A	0.9500	N4—C16	1.487 (3)
C4—C5	1.384 (4)	N4—C18	1.487 (3)
C4—H4A	0.9500	N4—H4N	0.890 (10)
C5—C6	1.383 (4)	C15—C16	1.539 (3)
C6—C7	1.388 (3)	C15—H15A	0.9900
C6—H6	0.9500	C15—H15B	0.9900
C7—H7	0.9500	C16—H16A	0.9900
O5—C8	1.273 (3)	C16—H16B	0.9900
O6—C8	1.231 (3)	C17—C18	1.538 (3)

O7—N2	1.228 (3)	C17—H17A	0.9900
O8—N2	1.226 (3)	C17—H17B	0.9900
N2—C12	1.479 (3)	C18—H18A	0.9900
C8—C9	1.523 (3)	C18—H18B	0.9900
C9—C10	1.387 (3)	C19—C20	1.538 (3)
C9—C14	1.395 (3)	C19—H19A	0.9900
C10—C11	1.393 (4)	C19—H19B	0.9900
C10—H10	0.9500	C20—H20A	0.9900
C11—C12	1.382 (4)	C20—H20B	0.9900
C11—H11	0.9500		
O4—N1—O3	124.0 (2)	C17—N3—C15	108.68 (19)
O4—N1—C5	117.9 (2)	C17—N3—C19	109.7 (2)
O3—N1—C5	118.0 (2)	C15—N3—C19	109.7 (2)
O2—C1—O1	125.6 (2)	C17—N3—H3N	105 (2)
O2—C1—C2	119.0 (2)	C15—N3—H3N	110 (2)
O1—C1—C2	115.4 (2)	C19—N3—H3N	114 (2)
C3—C2—C7	120.3 (2)	C20—N4—C16	109.6 (2)
C3—C2—C1	119.7 (2)	C20—N4—C18	109.5 (2)
C7—C2—C1	120.0 (2)	C16—N4—C18	108.9 (2)
C2—C3—C4	120.2 (2)	C20—N4—H4N	103 (2)
C2—C3—H3A	119.9	C16—N4—H4N	111 (2)
C4—C3—H3A	119.9	C18—N4—H4N	115 (2)
C5—C4—C3	118.2 (2)	N3—C15—C16	108.8 (2)
C5—C4—H4A	120.9	N3—C15—H15A	109.9
C3—C4—H4A	120.9	C16—C15—H15A	109.9
C6—C5—C4	123.2 (2)	N3—C15—H15B	109.9
C6—C5—N1	118.9 (2)	C16—C15—H15B	109.9
C4—C5—N1	117.9 (2)	H15A—C15—H15B	108.3
C5—C6—C7	117.9 (2)	N4—C16—C15	108.4 (2)
C5—C6—H6	121.1	N4—C16—H16A	110.0
C7—C6—H6	121.1	C15—C16—H16A	110.0
C6—C7—C2	120.3 (2)	N4—C16—H16B	110.0
C6—C7—H7	119.8	C15—C16—H16B	110.0
C2—C7—H7	119.8	H16A—C16—H16B	108.4
O8—N2—O7	124.1 (2)	N3—C17—C18	109.0 (2)
O8—N2—C12	118.0 (2)	N3—C17—H17A	109.9
O7—N2—C12	118.0 (2)	C18—C17—H17A	109.9
O6—C8—O5	125.5 (2)	N3—C17—H17B	109.9
O6—C8—C9	119.2 (2)	C18—C17—H17B	109.9
O5—C8—C9	115.3 (2)	H17A—C17—H17B	108.3
C10—C9—C14	120.2 (2)	N4—C18—C17	108.3 (2)
C10—C9—C8	120.2 (2)	N4—C18—H18A	110.0
C14—C9—C8	119.6 (2)	C17—C18—H18A	110.0
C9—C10—C11	120.0 (2)	N4—C18—H18B	110.0
C9—C10—H10	120.0	C17—C18—H18B	110.0
C11—C10—H10	120.0	H18A—C18—H18B	108.4
C12—C11—C10	118.5 (2)	N3—C19—C20	108.2 (2)

C12—C11—H11	120.8	N3—C19—H19A	110.1
C10—C11—H11	120.8	C20—C19—H19A	110.1
C11—C12—C13	122.9 (2)	N3—C19—H19B	110.1
C11—C12—N2	117.8 (2)	C20—C19—H19B	110.1
C13—C12—N2	119.3 (2)	H19A—C19—H19B	108.4
C12—C13—C14	117.9 (2)	N4—C20—C19	109.0 (2)
C12—C13—H13	121.1	N4—C20—H20A	109.9
C14—C13—H13	121.1	C19—C20—H20A	109.9
C13—C14—C9	120.5 (2)	N4—C20—H20B	109.9
C13—C14—H14	119.7	C19—C20—H20B	109.9
C9—C14—H14	119.7	H20A—C20—H20B	108.3
O2—C1—C2—C3	6.3 (4)	C10—C11—C12—N2	-179.8 (2)
O1—C1—C2—C3	-173.0 (2)	O8—N2—C12—C11	-172.9 (2)
O2—C1—C2—C7	-173.4 (2)	O7—N2—C12—C11	7.4 (3)
O1—C1—C2—C7	7.3 (3)	O8—N2—C12—C13	7.4 (3)
C7—C2—C3—C4	-1.2 (4)	O7—N2—C12—C13	-172.3 (2)
C1—C2—C3—C4	179.1 (2)	C11—C12—C13—C14	-0.1 (4)
C2—C3—C4—C5	0.5 (4)	N2—C12—C13—C14	179.7 (2)
C3—C4—C5—C6	0.3 (4)	C12—C13—C14—C9	0.2 (4)
C3—C4—C5—N1	178.4 (2)	C10—C9—C14—C13	-0.2 (4)
O4—N1—C5—C6	170.3 (2)	C8—C9—C14—C13	178.0 (2)
O3—N1—C5—C6	-9.1 (4)	C17—N3—C15—C16	50.7 (3)
O4—N1—C5—C4	-7.9 (3)	C19—N3—C15—C16	-69.2 (3)
O3—N1—C5—C4	172.7 (2)	C20—N4—C16—C15	50.2 (3)
C4—C5—C6—C7	-0.3 (4)	C18—N4—C16—C15	-69.6 (3)
N1—C5—C6—C7	-178.4 (2)	N3—C15—C16—N4	15.8 (3)
C5—C6—C7—C2	-0.4 (4)	C15—N3—C17—C18	-69.4 (2)
C3—C2—C7—C6	1.1 (4)	C19—N3—C17—C18	50.5 (3)
C1—C2—C7—C6	-179.2 (2)	C20—N4—C18—C17	-68.8 (3)
O6—C8—C9—C10	-2.6 (4)	C16—N4—C18—C17	51.1 (3)
O5—C8—C9—C10	177.5 (2)	N3—C17—C18—N4	15.8 (3)
O6—C8—C9—C14	179.2 (2)	C17—N3—C19—C20	-68.8 (3)
O5—C8—C9—C14	-0.7 (4)	C15—N3—C19—C20	50.6 (3)
C14—C9—C10—C11	0.1 (4)	C16—N4—C20—C19	-68.9 (3)
C8—C9—C10—C11	-178.1 (2)	C18—N4—C20—C19	50.5 (3)
C9—C10—C11—C12	0.0 (4)	N3—C19—C20—N4	15.6 (3)
C10—C11—C12—C13	0.0 (4)		

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N3—H3N $\cdots$ O1	0.88 (2)	1.66 (2)	2.539 (3)	173 (2)
N4—H4N $\cdots$ O5	0.89 (2)	1.65 (2)	2.542 (3)	175 (3)
C15—H15A $\cdots$ O6 <sup>i</sup>	0.99	2.42	3.193 (3)	134
C16—H16A $\cdots$ O4 <sup>ii</sup>	0.99	2.42	3.375 (3)	161
C17—H17A $\cdots$ O7 <sup>iii</sup>	0.99	2.42	3.338 (3)	153
C17—H17B $\cdots$ O6 <sup>i</sup>	0.99	2.42	3.313 (3)	149

---

C20—H20A···O2 <sup>iv</sup>	0.99	2.41	3.043 (3)	121
C20—H20B···O8 <sup>v</sup>	0.99	2.42	3.339 (3)	154

---

Symmetry codes: (i)  $-x+2, -y+1, -z+1$ ; (ii)  $x+1, y, z-1$ ; (iii)  $x-1, y, z+1$ ; (iv)  $-x+1, -y+2, -z+1$ ; (v)  $-x+2, -y+2, -z$ .