

Keywords: crystal structure; *cis*-platinum(II) complexes; hydrogen bonding; anticancer activity

CCDC reference: 1004305

Supporting information: this article has supporting information at journals.iucr.org/e

Crystal structure of chlorido(piperidine- κN)-(quinoline-2-carboxylato- $\kappa^2 N,O$)platinum(II)

Chi Nguyen Thi Thanh,^a Ngan Nguyen Bich^a and Luc Van Meervelt^{b*}



^aChemistry Department, Hanoi National University of Education, 136 – Xuan Thuy – Cau Giay, Hanoi, Vietnam, and

^bChemistry Department, KU Leuven, Celestijnenlaan 200F, B-3001 Leuven (Heverlee), Belgium. *Correspondence e-mail: luc.vanmeervelt@chem.kuleuven.be

The title compound, [Pt(C₁₀H₆NO₂)Cl(C₅H₁₁N)], crystallizes with one molecule in the asymmetric unit. The Pt^{II} cation has a slightly distorted square-planar coordination environment defined by a chloride anion, the quinoline N atom and a carboxylate O atom of the bidentate quinaldate ligand and a piperidine N atom. An intramolecular C–H···Cl hydrogen bond occurs. In the crystal, molecules are stacked into columns along the *c* axis by the formation of N–H···Cl and C–H···O hydrogen bonds.

1. Chemical context

The title compound belongs to a series of platinum(II) complexes bearing piperidine (pip) as a ligand, which exhibit notable antitumour activity (Da *et al.*, 2001; Rounaq Ali Khan *et al.*, 2000; Solin *et al.*, 1982). In comparison with the earlier reported complex [PtCl₂(pip)(quinoline)] (Nguyen Thi Thanh *et al.*, 2014), the quinoline ligand is replaced by an *N,O*-bidentate quinaldate ligand. It is interesting to note that in the [PtCl₂(pip)(quinoline)] complex, the quinoline and piperidine ligands are arranged in *cis* positions (Nguyen Thi Thanh *et al.*, 2014). In the title compound, the quinoline ring of the quinaldate ligand occupies a *trans* position with respect to the piperidine ring. We suggest that in the reaction solution there exists a chemical equilibrium between the neutral and bipolar forms of quinaldic acid. Thus, the quinaldic acid in its ionic form coordinates with Pt^{II} *via* the O atom of the carboxylate group first and in a *cis* position with respect to piperidine based on the *trans* effect. In a second step, the quinaldic acid coordinates with Pt^{II} also *via* its N atom, resulting in the cyclic complex.



The anticancer activity of the title compound was tested according to the method described in Skehan *et al.* (1990) on four human cancer cells of HepG2, RD, MCF7 and FI. The IC₅₀ values calculated based on OD values taken on an Elisa instrument at 515–540 nm are 4.46, 2.59, >10 and 5.60 μg ml⁻¹, respectively.




Figure 1
View of the molecular structure of the title compound, showing the atom-labelling scheme, with ellipsoids drawn at the 50% probability level. The intramolecular C—H···Cl hydrogen bond is shown as a green dashed line.

2. Structural commentary

The title complex crystallizes with one molecule per asymmetric unit (Fig. 1). The Pt^{II} cation is surrounded by two N atoms, one O atom and one Cl atom, resulting in a slightly distorted square-planar coordination environment [angles around platinum: O1—Pt1—N1 81.38 (9), O1—Pt1—N2 88.26 (9), Cl1—Pt1—N2 84.26 (7) and Cl1—Pt1—N1 106.11 (7)°]. The Cl[−] and the Pt^{II} atoms are displaced from the least-squares plane of the quinoline ring and all other coordinating atoms by 0.2936 (7) and 0.0052 (1) Å, respectively. The piperidine ring adopts a chair conformation and is almost perpendicular to the coordination plane of the Pt^{II} cation [dihedral angle between the best plane through the piperidine ring and the four atoms coordinating to the Pt^{II} cation = 79.66 (13)°]. Bond lengths are normal and agree well with




Figure 2
View of the crystal packing for the title compound, with (N/C)—H···Cl and C—H···O hydrogen bonds drawn as green and red dashed lines, respectively. [Symmetry codes: (i) $x, -y + 1, z - \frac{1}{2}$; (ii) $-x + 1, y, -z + \frac{3}{2}$]

Table 1
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N2—H2A···Cl1 ⁱ	0.93	2.74	3.624 (2)	160
C3—H3···O2 ⁱⁱ	0.96	2.53	3.360 (4)	145
C8—H8···Cl1	0.95	2.40	3.268 (3)	152

Symmetry codes: (i) $-x + 1, y, -z + \frac{3}{2}$; (ii) $x, -y + 1, z - \frac{1}{2}$.

related platinum compounds (Cambridge Structural Database, version 5.34; Allen, 2002). There is an intramolecular hydrogen bond between atom Cl1 and atom H8 (Fig. 1 and Table 1).

3. Supramolecular features

The crystal packing is characterized by N—H···Cl and C—H···O hydrogen bonds (Table 1). Molecules are arranged into columns along the c axis (Fig. 2) with the piperidine rings all directed towards the center of the column, favouring hydrophobic interactions.

4. Synthesis and crystallization

The starting complex K[PtCl₃(piperidine)] (0.425 g, 1 mmol), prepared according to the synthetic protocol of Da *et al.* (2001), was dissolved in water (10 ml) and filtered to afford a clear solution. To this solution, quinaldic acid (1.2 mmol) in an aqueous ethanol solution (5 ml, 1:1 v/v) was added gradually

Table 2
Experimental details.

Crystal data	[Pt(C ₁₀ H ₆ NO ₂)Cl(C ₅ H ₁₁ N)]
Chemical formula	487.85
M_r	Monoclinic, $C2/c$
Crystal system, space group	200
Temperature (K)	22.7542 (8), 9.7540 (3), 14.0139 (5)
a, b, c (Å)	95.542 (3)
β (°)	3095.78 (19)
V (Å ³)	8
Z	Mo $K\alpha$
Radiation type	9.24
μ (mm ^{−1})	0.3 × 0.3 × 0.2
Crystal size (mm)	
Data collection	
Diffractometer	Agilent SuperNova (single source at offset, Eos detector)
Absorption correction	Multi-scan (<i>CrysAlis PRO</i> ; Agilent, 2012)
T_{\min}, T_{\max}	0.473, 1.000
No. of measured, independent and observed [$I > 2\sigma(I)$] reflections	31419, 3166, 2951
R_{int}	0.026
$(\sin \theta/\lambda)_{\text{max}}$ (Å ^{−1})	0.625
Refinement	
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.015, 0.035, 1.12
No. of reflections	3166
No. of parameters	190
H-atom treatment	H-atom parameters constrained
$\Delta\rho_{\text{max}}, \Delta\rho_{\text{min}}$ (e Å ^{−3})	0.80, −0.53

Computer programs: *CrysAlis PRO* (Agilent, 2012), *SHELXS97* and *SHELXL97* (Sheldrick, 2008) and *OLEX2* (Dolomanov *et al.*, 2009).

while stirring at room temperature for 1 h. The reaction mixture was stirred further for 4 h. The precipitated yellow substance was filtered off and washed consecutively with a 0.1 M HCl solution (2 × 2 ml), warm water (3 × 2 ml) and cold ethanol (2 ml). The product was then dried in a vacuum at 323 K for 4 h. The yield was 80%. Single crystals suitable for X-ray diffraction analysis were obtained by slow evaporation from an ethanol–water (1:1 v/v) solution at room temperature. Positive ESI–MS: m/z 1973 [$4M + Na$]⁺, 1483 [$3M + Na$]⁺, 998 [$2M + Na$]⁺, 510 [$M + Na$]⁺, 977 [$2M + H$]⁺, 489 [$M + H$]⁺; IR (KBr) cm^{-1} : 3192 (ν_{NH}); 3080, 2930, 2866 (ν_{CH}); 1678 ($\nu_{C=O}$); 1592, 1459 ($\nu_{C=C}$ arom); 1334 (ν_{C-O}); ¹H NMR (δ p.p.m; CDCl₃, 500Hz): 9.50 (1H, *d*, ³*J* = 9.0 Hz, Ar-*H*), 8.51 (1H, *d*, ³*J* = 8.0 Hz, Ar-*H*), 8.06 (1H, *d*, ³*J* = 8.0 Hz, Ar-*H*), 7.91–7.88 (2H, *ov*, Ar-*H*), 7.71 (1H, *t*, ³*J* = 8.0 Hz, Ar-*H*), 3.52 (2H_α^e, *d*, ²*J*_{ae} = 12.5 Hz, C₅H₁₀NH), 3.27 (2H_α^a, *q*, ²*J*_{ae}, ³*J*_{aa}, ³*J*_{aa(NH)} = 12.5 Hz, C₅H₁₀NH), 1.76–1.61 (4H_β, 2H_γ, *ov*, C₅H₁₀NH), 4.00 (1H, *br*, C₅H₁₀NH).

5. Refinement

All H atoms were refined using a riding model, with C–H = 0.95 Å for aromatic, C–H = 0.99 Å for CH₂ and N–H = 0.93 Å for amino H atoms, with $U_{iso} = 1.2U_{eq}(C,N)$.

Acknowledgements

The authors thank the Vietnamese Ministry of Education (project No. B2013-17-39) for financial support and the Hercules Foundation for supporting the purchase of the diffractometer through project AKUL/09/0035.

References

- Agilent (2012). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.
- Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
- Da, T. T., Vu, D. B. & Dinh, N. H. (2001). *J. Pharm. Sci. (Vietnam)*, **6**, 6–8.
- Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
- Nguyen Thi Thanh, C., Nguyen Bich, N. & Van Meervelt, L. (2014). *Acta Cryst.* **C70**, 297–301.
- Rounaq Ali Khan, S., Guzman-Jimenez, I., Whitmire, K. H. & Khokhar, A. R. (2000). *Polyhedron*, **19**, 975–981.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Skehan, P., Storeng, R., Scudiero, D., Monks, A., McMahon, J., Vistica, D., Warren, J. T., Bokesch, H., Kenney, S. & Boyd, M. R. (1990). *J. Natl Cancer Inst.* **82**, 1107–1112.
- Solin, T., Matsumoto, K. & Fuwa, K. (1982). *Inorg. Chim. Acta*, **65**, L172–L172.

supporting information

Acta Cryst. (2014). E70, 36-38 [https://doi.org/10.1107/S160053681401191X]

Crystal structure of chlorido(piperidine- κ N)(quinoline-2-carboxylato- κ^2 N,O)platinum(II)

Chi Nguyen Thi Thanh, Ngan Nguyen Bich and Luc Van Meervelt

Computing details

Data collection: *CrysAlis PRO* (Agilent, 2012); cell refinement: *CrysAlis PRO* (Agilent, 2012); data reduction: *CrysAlis PRO* (Agilent, 2012); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: OLEX2 (Dolomanov *et al.*, 2009); software used to prepare material for publication: OLEX2 (Dolomanov *et al.*, 2009).

Chlorido(piperidine- κ N)(quinoline-2-carboxylato- κ^2 N,O)platinum(II)

Crystal data

[Pt(C₁₀H₆NO₂)Cl(C₅H₁₁N)]

$M_r = 487.85$

Monoclinic, *C2/c*

$a = 22.7542$ (8) Å

$b = 9.7540$ (3) Å

$c = 14.0139$ (5) Å

$\beta = 95.542$ (3)°

$V = 3095.78$ (19) Å³

$Z = 8$

$F(000) = 1856$

$D_x = 2.093$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.7107$ Å

Cell parameters from 16449 reflections

$\theta = 3.4$ – 29.8 °

$\mu = 9.24$ mm⁻¹

$T = 200$ K

Block, yellow

$0.3 \times 0.3 \times 0.2$ mm

Data collection

Agilent SuperNova (single source at offset, Eos detector)

diffractometer

Radiation source: SuperNova (Mo) X-ray

Source

Mirror monochromator

Detector resolution: 15.9631 pixels mm⁻¹

ω scans

Absorption correction: multi-scan

(*CrysAlis PRO*; Agilent, 2012)

$T_{\min} = 0.473$, $T_{\max} = 1.000$

31419 measured reflections

3166 independent reflections

2951 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.026$

$\theta_{\max} = 26.4$ °, $\theta_{\min} = 2.8$ °

$h = -28 \rightarrow 28$

$k = -12 \rightarrow 12$

$l = -17 \rightarrow 17$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.015$

$wR(F^2) = 0.035$

$S = 1.12$

3166 reflections

190 parameters

0 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0118P)^2 + 9.138P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.002$

$$\Delta\rho_{\max} = 0.80 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.53 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.29348 (12)	0.6098 (3)	0.5028 (2)	0.0268 (6)
C2	0.25609 (13)	0.5746 (3)	0.4211 (2)	0.0327 (7)
H2	0.2227	0.5176	0.4264	0.039*
C3	0.26831 (14)	0.6232 (3)	0.3338 (2)	0.0354 (7)
H3	0.2432	0.6014	0.2777	0.042*
C4	0.31811 (14)	0.7052 (3)	0.3278 (2)	0.0324 (7)
C5	0.33360 (18)	0.7554 (4)	0.2386 (2)	0.0432 (8)
H5	0.3085	0.7372	0.1818	0.052*
C6	0.38367 (19)	0.8290 (4)	0.2332 (2)	0.0516 (10)
H6	0.3934	0.8625	0.1730	0.062*
C7	0.42105 (18)	0.8556 (4)	0.3167 (2)	0.0503 (9)
H7	0.4565	0.9057	0.3121	0.060*
C8	0.40769 (16)	0.8111 (3)	0.4047 (2)	0.0400 (8)
H8	0.4336	0.8309	0.4603	0.048*
C9	0.35561 (14)	0.7362 (3)	0.4130 (2)	0.0285 (6)
C10	0.28109 (13)	0.5514 (3)	0.5984 (2)	0.0318 (7)
C11	0.42217 (15)	0.5840 (3)	0.8206 (2)	0.0402 (8)
H11A	0.4518	0.5401	0.7832	0.048*
H11B	0.3857	0.5281	0.8125	0.048*
C12	0.44551 (16)	0.5867 (4)	0.9262 (3)	0.0479 (9)
H12A	0.4509	0.4914	0.9499	0.058*
H12B	0.4845	0.6323	0.9333	0.058*
C13	0.40413 (18)	0.6613 (5)	0.9861 (2)	0.0550 (10)
H13A	0.4225	0.6690	1.0528	0.066*
H13B	0.3671	0.6083	0.9869	0.066*
C14	0.39015 (16)	0.8039 (4)	0.9459 (2)	0.0443 (9)
H14A	0.4263	0.8607	0.9527	0.053*
H14B	0.3604	0.8481	0.9828	0.053*
C15	0.36650 (13)	0.7956 (3)	0.8404 (2)	0.0337 (7)
H15A	0.3285	0.7455	0.8344	0.040*
H15B	0.3591	0.8893	0.8150	0.040*
Cl1	0.45271 (3)	0.88368 (8)	0.62745 (5)	0.03279 (16)

N1	0.34113 (10)	0.6894 (2)	0.50082 (16)	0.0255 (5)
N2	0.40915 (10)	0.7240 (2)	0.78242 (16)	0.0264 (5)
H2A	0.4443	0.7733	0.7893	0.032*
O1	0.31888 (9)	0.5856 (2)	0.66939 (14)	0.0369 (5)
O2	0.23904 (10)	0.4760 (3)	0.60452 (16)	0.0433 (6)
Pt1	0.380802 (4)	0.720087 (11)	0.639573 (7)	0.02386 (4)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0264 (14)	0.0266 (14)	0.0271 (14)	0.0071 (12)	0.0005 (11)	-0.0021 (12)
C2	0.0288 (15)	0.0374 (17)	0.0308 (15)	0.0016 (13)	-0.0034 (12)	-0.0040 (13)
C3	0.0374 (16)	0.0387 (18)	0.0281 (15)	0.0067 (14)	-0.0071 (12)	-0.0063 (13)
C4	0.0425 (17)	0.0282 (16)	0.0256 (15)	0.0079 (13)	-0.0006 (12)	-0.0035 (12)
C5	0.067 (2)	0.0370 (18)	0.0247 (16)	-0.0022 (17)	-0.0017 (15)	-0.0011 (13)
C6	0.086 (3)	0.042 (2)	0.0269 (16)	-0.014 (2)	0.0094 (17)	0.0007 (15)
C7	0.071 (3)	0.046 (2)	0.0345 (18)	-0.0231 (19)	0.0125 (17)	-0.0054 (16)
C8	0.052 (2)	0.0383 (18)	0.0298 (16)	-0.0106 (15)	0.0033 (14)	-0.0039 (13)
C9	0.0397 (16)	0.0213 (14)	0.0242 (14)	0.0045 (12)	0.0010 (12)	-0.0031 (11)
C10	0.0282 (15)	0.0393 (17)	0.0268 (14)	0.0012 (13)	-0.0028 (12)	-0.0017 (13)
C11	0.0421 (18)	0.0303 (17)	0.0443 (19)	0.0005 (14)	-0.0153 (15)	0.0051 (14)
C12	0.051 (2)	0.040 (2)	0.048 (2)	-0.0110 (16)	-0.0235 (17)	0.0172 (16)
C13	0.060 (2)	0.072 (3)	0.0299 (17)	-0.024 (2)	-0.0086 (16)	0.0155 (18)
C14	0.0404 (18)	0.065 (3)	0.0262 (16)	-0.0016 (17)	-0.0010 (13)	-0.0030 (15)
C15	0.0296 (15)	0.0433 (19)	0.0275 (15)	0.0027 (13)	-0.0014 (12)	0.0001 (13)
C11	0.0348 (4)	0.0335 (4)	0.0292 (3)	-0.0071 (3)	-0.0013 (3)	-0.0006 (3)
N1	0.0274 (12)	0.0244 (12)	0.0241 (11)	0.0033 (9)	-0.0002 (9)	-0.0026 (9)
N2	0.0240 (11)	0.0299 (13)	0.0242 (12)	-0.0014 (10)	-0.0028 (9)	0.0022 (10)
O1	0.0349 (11)	0.0493 (14)	0.0251 (10)	-0.0115 (10)	-0.0044 (9)	0.0061 (10)
O2	0.0350 (12)	0.0585 (16)	0.0350 (12)	-0.0179 (11)	-0.0031 (9)	0.0032 (11)
Pt1	0.02348 (6)	0.02556 (6)	0.02184 (6)	0.00129 (4)	-0.00148 (4)	-0.00049 (4)

Geometric parameters (Å, °)

C1—C2	1.402 (4)	C11—H11B	0.9900
C1—C10	1.507 (4)	C11—C12	1.524 (4)
C1—N1	1.336 (4)	C11—N2	1.486 (4)
C2—H2	0.9500	C12—H12A	0.9900
C2—C3	1.365 (4)	C12—H12B	0.9900
C3—H3	0.9500	C12—C13	1.507 (6)
C3—C4	1.396 (5)	C13—H13A	0.9900
C4—C5	1.419 (4)	C13—H13B	0.9900
C4—C9	1.432 (4)	C13—C14	1.522 (5)
C5—H5	0.9500	C14—H14A	0.9900
C5—C6	1.355 (5)	C14—H14B	0.9900
C6—H6	0.9500	C14—C15	1.526 (4)
C6—C7	1.403 (5)	C15—H15A	0.9900
C7—H7	0.9500	C15—H15B	0.9900

C7—C8	1.369 (5)	C15—N2	1.497 (4)
C8—H8	0.9500	Cl1—Pt1	2.3035 (7)
C8—C9	1.407 (5)	N1—Pt1	2.085 (2)
C9—N1	1.382 (4)	N2—H2A	0.9300
C10—O1	1.295 (3)	N2—Pt1	2.043 (2)
C10—O2	1.217 (4)	O1—Pt1	1.999 (2)
C11—H11A	0.9900		
C2—C1—C10	118.8 (3)	H12A—C12—H12B	107.9
N1—C1—C2	123.8 (3)	C13—C12—C11	111.8 (3)
N1—C1—C10	117.4 (2)	C13—C12—H12A	109.3
C1—C2—H2	120.4	C13—C12—H12B	109.3
C3—C2—C1	119.1 (3)	C12—C13—H13A	109.5
C3—C2—H2	120.4	C12—C13—H13B	109.5
C2—C3—H3	120.3	C12—C13—C14	110.8 (3)
C2—C3—C4	119.3 (3)	H13A—C13—H13B	108.1
C4—C3—H3	120.3	C14—C13—H13A	109.5
C3—C4—C5	121.6 (3)	C14—C13—H13B	109.5
C3—C4—C9	119.5 (3)	C13—C14—H14A	109.5
C5—C4—C9	118.9 (3)	C13—C14—H14B	109.5
C4—C5—H5	119.5	C13—C14—C15	110.6 (3)
C6—C5—C4	120.9 (3)	H14A—C14—H14B	108.1
C6—C5—H5	119.5	C15—C14—H14A	109.5
C5—C6—H6	120.1	C15—C14—H14B	109.5
C5—C6—C7	119.8 (3)	C14—C15—H15A	109.4
C7—C6—H6	120.1	C14—C15—H15B	109.4
C6—C7—H7	119.2	H15A—C15—H15B	108.0
C8—C7—C6	121.6 (3)	N2—C15—C14	111.4 (3)
C8—C7—H7	119.2	N2—C15—H15A	109.4
C7—C8—H8	120.0	N2—C15—H15B	109.4
C7—C8—C9	120.1 (3)	C1—N1—C9	118.3 (2)
C9—C8—H8	120.0	C1—N1—Pt1	110.10 (18)
C8—C9—C4	118.7 (3)	C9—N1—Pt1	131.61 (19)
N1—C9—C4	119.9 (3)	C11—N2—C15	110.5 (2)
N1—C9—C8	121.4 (3)	C11—N2—H2A	107.4
O1—C10—C1	114.8 (3)	C11—N2—Pt1	111.61 (18)
O2—C10—C1	120.5 (3)	C15—N2—H2A	107.4
O2—C10—O1	124.7 (3)	C15—N2—Pt1	112.37 (17)
H11A—C11—H11B	107.9	Pt1—N2—H2A	107.4
C12—C11—H11A	109.2	C10—O1—Pt1	115.95 (19)
C12—C11—H11B	109.2	N1—Pt1—Cl1	106.11 (7)
N2—C11—H11A	109.2	N2—Pt1—Cl1	84.26 (7)
N2—C11—H11B	109.2	N2—Pt1—N1	169.63 (9)
N2—C11—C12	111.9 (3)	O1—Pt1—Cl1	171.99 (6)
C11—C12—H12A	109.3	O1—Pt1—N1	81.38 (9)
C11—C12—H12B	109.3	O1—Pt1—N2	88.26 (9)
C1—C2—C3—C4	0.8 (5)	C9—N1—Pt1—Cl1	8.2 (3)

C1—C10—O1—Pt1	5.1 (3)	C9—N1—Pt1—N2	-171.6 (4)
C1—N1—Pt1—Cl1	-171.51 (17)	C9—N1—Pt1—O1	-174.7 (3)
C1—N1—Pt1—N2	8.6 (6)	C10—C1—C2—C3	-177.6 (3)
C1—N1—Pt1—O1	5.58 (18)	C10—C1—N1—C9	175.6 (2)
C2—C1—C10—O1	177.8 (3)	C10—C1—N1—Pt1	-4.7 (3)
C2—C1—C10—O2	-0.7 (4)	C10—O1—Pt1—N1	-6.0 (2)
C2—C1—N1—C9	-2.1 (4)	C10—O1—Pt1—N2	174.6 (2)
C2—C1—N1—Pt1	177.6 (2)	C11—C12—C13—C14	-53.6 (4)
C2—C3—C4—C5	178.4 (3)	C11—N2—Pt1—Cl1	-129.0 (2)
C2—C3—C4—C9	0.4 (4)	C11—N2—Pt1—N1	50.9 (6)
C3—C4—C5—C6	-176.6 (3)	C11—N2—Pt1—O1	53.9 (2)
C3—C4—C9—C8	175.7 (3)	C12—C11—N2—C15	-56.3 (3)
C3—C4—C9—N1	-2.5 (4)	C12—C11—N2—Pt1	177.9 (2)
C4—C5—C6—C7	0.3 (6)	C12—C13—C14—C15	54.5 (4)
C4—C9—N1—C1	3.3 (4)	C13—C14—C15—N2	-56.9 (4)
C4—C9—N1—Pt1	-176.4 (2)	C14—C15—N2—C11	57.6 (3)
C5—C4—C9—C8	-2.3 (4)	C14—C15—N2—Pt1	-177.0 (2)
C5—C4—C9—N1	179.4 (3)	C15—N2—Pt1—Cl1	106.25 (19)
C5—C6—C7—C8	-1.3 (6)	C15—N2—Pt1—N1	-73.9 (6)
C6—C7—C8—C9	0.3 (6)	C15—N2—Pt1—O1	-70.9 (2)
C7—C8—C9—C4	1.5 (5)	N1—C1—C2—C3	0.0 (5)
C7—C8—C9—N1	179.7 (3)	N1—C1—C10—O1	0.0 (4)
C8—C9—N1—C1	-174.9 (3)	N1—C1—C10—O2	-178.5 (3)
C8—C9—N1—Pt1	5.4 (4)	N2—C11—C12—C13	54.9 (4)
C9—C4—C5—C6	1.4 (5)	O2—C10—O1—Pt1	-176.5 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N2—H2A \cdots Cl1 ⁱ	0.93	2.74	3.624 (2)	160
C3—H3 \cdots O2 ⁱⁱ	0.96	2.53	3.360 (4)	145
C8—H8 \cdots Cl1	0.95	2.40	3.268 (3)	152

Symmetry codes: (i) $-x+1, y, -z+3/2$; (ii) $x, -y+1, z-1/2$.