

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3,10,14,21-Tetrakis(4-methoxyphenyl)-pentacyclo[11.8.0.0^{2,11}.0^{4,9}.0^{15,20}]-henicosa-1(21),2,4(9),5,7,10,13,15(20),-16,18-decaen-12-one chloroform monosolvate

S. Gopinath,^a P. Narayanan,^a K. Sethusankar,^{a*} Meganathan Nandakumar^b and Arasambattu K. Mohanakrishnan^b

^aDepartment of Physics, RKM Vivekananda College (Autonomous), Chennai 600 004, India, and ^bDepartment of Organic Chemistry, University of Madras, Guindy Campus, Chennai 600 025, India
Correspondence e-mail: ksethusankar@yahoo.co.in

Received 23 June 2014; accepted 15 July 2014


Edited by E. R. T. Tiekink, University of Malaya, Malaysia

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; some non-H atoms missing; R factor = 0.040; wR factor = 0.125; data-to-parameter ratio = 14.0.

The asymmetric unit of the title compound, $\text{C}_{49}\text{H}_{36}\text{O}_6 \cdot \text{CHCl}_3$, contains half an organic molecule, the complete molecule being generated by the operation of a crystallographic twofold rotation axis, and half a highly disordered chloroform molecule. The contribution to the diffraction pattern of the latter was removed using the program SQUEEZE in PLATON [Spek (2009). *Acta Cryst.* **D65**, 148–155]; the unit-cell characteristics take into account the presence of CHCl_3 . The dihedral angles between the planes of the naphthalene ring system and the methoxybenzene rings are 71.05 (7) (*syn* to the central $\text{C}=\text{O}$ group) and 57.27 (6)° (*anti* to the central $\text{C}=\text{O}$ group). In the crystal, molecules are linked by $\text{C}-\text{H} \cdots \text{O}$ interactions, generating $C(12)$ chains running parallel to the b axis.

Related literature

For the uses and biological importance of naphthalene, see: Morikawa & Takahashi (2004); Rokade & Sayyed (2009).



Experimental

Crystal data

 $\text{C}_{49}\text{H}_{36}\text{O}_6 \cdot \text{CHCl}_3$ $M_r = 824.15$ Monoclinic, $P2_1/n$ $a = 9.7720$ (4) Å $b = 12.0351$ (4) Å $c = 17.3162$ (6) Å $\beta = 93.947$ (1)° $V = 2031.68$ (13) Å³ $Z = 2$ Mo $K\alpha$ radiation $\mu = 0.28$ mm⁻¹ $T = 296$ K $0.35 \times 0.30 \times 0.30$ mm

Data collection

Bruker Kappa APEXII CCD diffractometer

Absorption correction: multi-scan (SADABS; Bruker, 2008)

 $T_{\min} = 0.892$, $T_{\max} = 0.934$

20614 measured reflections

3474 independent reflections

2664 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.025$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.040$ $wR(F^2) = 0.125$ $S = 1.09$

3474 reflections

248 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.13$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.15$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{C}24-\text{H}24\text{c} \cdots \text{O}3^i$	0.96	2.39	3.199 (3)	141

Symmetry code: (i) $x, y + 1, z$.

Data collection: APEX2 (Bruker, 2008); cell refinement: APEX2; data reduction: SAINT (Bruker, 2008); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 2012) and Mercury (Macrae *et al.*, 2008); software used to prepare material for publication: SHELXL97 and PLATON (Spek, 2009).

SG and KS thank Dr Babu Varghese, Senior Scientific Officer, SAIF, IIT, Madras (India), for the X-ray intensity data collection.

Supporting information for this paper is available from the IUCr electronic archives (Reference: TK5322).

References

Bruker (2008). *APEX2*, *SAINTE* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.

- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Morikawa, H. & Takahashi, M. (2004). US Patent No. 6800482.
- Rokade, Y. B. & Sayyed, R. Z. (2009). *Rasayan J. Chem.* **2**, 972–980.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supporting information

Acta Cryst. (2014). E70, o873–o874 [doi:10.1107/S1600536814016389]

3,10,14,21-Tetrakis(4-methoxyphenyl)pentacyclo- [11.8.0.0^{2,11}.0^{4,9}.0^{15,20}]henicosa-1(21),2,4(9),5,7,10,13,15(20),16,18-decaen-12- one chloroform monosolvate

S. Gopinath, P. Narayanan, K. Sethusankar, Meganathan Nandakumar and Arasambattu K. Mohanakrishnan

S1. Structural commentary

Naphthalene derivatives have been extensively employed in many fields and some possess important biological and commercial applications such as disinfectants, insecticides, plant hormones and rooting agents (Morikawa & Takahashi, 2004). Naphthalene has been identified as new range of potent anti-microbiols effective against wide range of human pathogens (Rokade & Sayyed, 2009).

The title compound, C₄₉H₃₆O₆·CHCl₃, comprises half the organic molecule in the asymmetric unit, the complete molecule is being generated by two fold rotation, as well as half a disordered chloroform molecule. The X-ray analysis confirms the molecular structure and atom connectivity of the compound as illustrated in Fig. 1.

The dihedral angles between naphthalene ring and methoxy-substituted benzene rings are 71.05 (7) and 57.27 (6)°, respectively. The dihedral angle between naphthalene and cyclopenta-2,4-dienone residue is 13.26 (6)°.


In the crystal packing, molecules are linked *via* bifurcated C24—H24cⁱⁱ···O3ⁱⁱⁱ intermolecular hydrogen bonding, which generates C(12) infinite chains running parallel to the *b* axis. A view of the supramolecular chain is shown in Fig. 2.

S2. Synthesis and crystallization

To a solution of benzo[*c*]furan (0.20 g, 0.60 mmol) in dry dichloromethane (15 ml), indenone (0.24 g, 0.60 mmol) was added and refluxed till the disappearance of colour due to the benzo[*c*]furan (10 h). To this, PTSA (0.46 g, 2.42 mmol) was added (9 h). The reaction mixture was poured into saturated solution of NaHCO₃ (40 ml), extracted with CHCl₃ (3 x 15 ml) and dried (Na₂SO₄). Removal of solvent followed by column chromatographic purification (silica gel; 10% ethyl acetate in hexane) afforded the fluorenone derivative as a yellow solid. Single crystals suitable for X-ray diffraction were prepared by slow evaporation of a solution of the title compound in chloroform at room temperature.

S3. Refinement

The hydrogen atoms bound to the C atoms are treated as riding atoms, with d(C–H) = 0.93–0.96 Å, and with $U_{iso}(H) = 1.2–1.5U_{eq}(C)$. The void in the crystal contains a highly disordered molecule of chloroform which has been removed from the intensity data using the SQUEEZE routine in *PLATON* [Spek (2009). *Acta Cryst.* D65, 148–155]. The squeeze results show removal of 58 electrons equivalent of scattering materials each from two regions in the unit cell. This corresponds exactly to the number of electrons present per molecule of chloroform.

**Figure 1**

The molecular structure of the title compound (chloroform molecule omitted) showing the atom numbering scheme and displacement ellipsoids drawn at 30% probability level. The symmetry code of the unlabelled atoms is $-x + 3/2, y, -z + 1/2$.




Figure 2

A partial packing diagram for the title compound viewed down the c axis, showing C24—H24c \cdots O3ⁱⁱ hydrogen bonds resulting in the formation of C(12) chains running parallel to the b axis. [hydrogen atoms not involved in the hydrogen bonding and chloroform molecules have been omitted for the clarity]. Symmetry code: (ii) $x, 1 + y, z$.

3,10,14,21-Tetrakis(4-methoxyphenyl)pentacyclo[11.8.0.0^{2,11}.0^{4,9}.0^{15,20}]henicosa-1(21),2,4(9),5,7,10,13,15 (20),16,18-decaen-12-one chloroform monosolvate

Crystal data

C₄₉H₃₆O₅·CHCl₃

$M_r = 824.15$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yac

$a = 9.7720$ (4) Å

$b = 12.0351$ (4) Å

$c = 17.3162$ (6) Å

$\beta = 93.947$ (1)°

$V = 2031.68$ (13) Å³

$Z = 2$

$F(000) = 740$

$D_x = 1.347$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3474 reflections

$\theta = 1.7$ – 24.8°

$\mu = 0.28$ mm⁻¹

$T = 296$ K

Block, yellow

$0.35 \times 0.30 \times 0.30$ mm

Data collection

Bruker Kappa APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω & φ scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2008)

$T_{\min} = 0.892$, $T_{\max} = 0.934$

20614 measured reflections

3474 independent reflections

2664 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.025$

$\theta_{\max} = 24.8^\circ$, $\theta_{\min} = 1.7^\circ$

$h = -11 \rightarrow 11$

$k = -14 \rightarrow 14$

$l = -20 \rightarrow 20$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.040$

$wR(F^2) = 0.125$

$S = 1.09$

3474 reflections

248 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0692P)^2 + 0.1579P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.13$ e Å⁻³

$\Delta\rho_{\min} = -0.15$ e Å⁻³

Extinction correction: *SHELXL97* (Sheldrick, 2008), $F_c^* = kFc[1 + 0.001xFe^2\lambda^3/\sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0041 (10)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	U_{iso}^*/U_{eq}
O1	0.70322 (14)	0.19072 (11)	-0.01039 (8)	0.0743 (4)
O2	1.07295 (15)	1.23830 (11)	0.27627 (8)	0.0773 (4)
O3	0.7500	0.48212 (15)	0.2500	0.0629 (5)
C1	0.78181 (15)	0.71567 (14)	0.05173 (8)	0.0461 (4)
C2	0.77909 (16)	0.69508 (16)	-0.02894 (8)	0.0524 (4)
H2	0.7573	0.6243	-0.0476	0.063*
C3	0.80786 (17)	0.77710 (16)	-0.07986 (9)	0.0575 (5)
H3	0.8026	0.7626	-0.1327	0.069*
C4	0.84506 (19)	0.88229 (16)	-0.05267 (9)	0.0600 (5)
H4	0.8680	0.9371	-0.0873	0.072*
C5	0.84809 (18)	0.90560 (16)	0.02447 (9)	0.0564 (4)
H5	0.8746	0.9761	0.0416	0.068*
C6	0.81187 (15)	0.82512 (14)	0.07917 (8)	0.0468 (4)
C7	0.80505 (16)	0.85209 (13)	0.16025 (8)	0.0465 (4)
C8	0.76176 (15)	0.76959 (13)	0.20778 (8)	0.0449 (4)
C9	0.75067 (15)	0.65740 (13)	0.18137 (8)	0.0464 (4)
C10	0.75817 (15)	0.62789 (13)	0.10526 (8)	0.0452 (4)
C11	0.74288 (16)	0.51114 (14)	0.07819 (8)	0.0471 (4)
C12	0.61738 (17)	0.45959 (15)	0.07359 (10)	0.0575 (5)
H12	0.5419	0.4976	0.0903	0.069*
C13	0.59946 (18)	0.35266 (15)	0.04487 (11)	0.0627 (5)
H13	0.5132	0.3197	0.0424	0.075*
C14	0.71000 (18)	0.29579 (14)	0.02007 (9)	0.0534 (4)
C15	0.83762 (18)	0.34534 (16)	0.02556 (10)	0.0599 (5)
H15	0.9134	0.3068	0.0098	0.072*
C16	0.85313 (17)	0.45107 (15)	0.05413 (10)	0.0548 (4)
H16	0.9398	0.4833	0.0575	0.066*
C17	0.86294 (16)	0.95837 (13)	0.19126 (8)	0.0463 (4)
C18	0.97613 (17)	0.95292 (14)	0.24523 (9)	0.0545 (4)
H18	1.0070	0.8839	0.2631	0.065*
C19	1.04222 (19)	1.04640 (15)	0.27230 (10)	0.0608 (5)
H19	1.1177	1.0405	0.3078	0.073*
C20	0.99703 (18)	1.15006 (14)	0.24698 (10)	0.0554 (4)

C21	0.88557 (18)	1.15873 (15)	0.19489 (9)	0.0549 (4)
H21	0.8545	1.2282	0.1780	0.066*
C22	0.81949 (17)	1.06346 (14)	0.16761 (9)	0.0521 (4)
H22	0.7437	1.0701	0.1324	0.063*
C23	0.5727 (3)	0.14149 (19)	-0.02289 (16)	0.0952 (7)
H23A	0.5322	0.1330	0.0257	0.143*
H23B	0.5817	0.0699	-0.0465	0.143*
H23C	0.5152	0.1880	-0.0564	0.143*
C24	1.0313 (3)	1.34544 (17)	0.25541 (15)	0.0896 (7)
H24A	1.0311	1.3536	0.2003	0.134*
H24B	1.0935	1.3983	0.2801	0.134*
H24C	0.9405	1.3583	0.2715	0.134*
C25	0.7500	0.5833 (2)	0.2500	0.0468 (5)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0740 (9)	0.0585 (8)	0.0895 (10)	-0.0005 (7)	-0.0016 (7)	-0.0103 (7)
O2	0.0810 (10)	0.0622 (9)	0.0863 (10)	-0.0095 (7)	-0.0112 (7)	-0.0093 (7)
O3	0.0864 (13)	0.0581 (11)	0.0450 (9)	0.000	0.0103 (8)	0.000
C1	0.0364 (8)	0.0661 (11)	0.0362 (8)	-0.0031 (7)	0.0044 (6)	0.0009 (7)
C2	0.0466 (9)	0.0760 (11)	0.0348 (8)	-0.0049 (8)	0.0037 (6)	-0.0056 (8)
C3	0.0545 (10)	0.0845 (13)	0.0337 (8)	-0.0034 (9)	0.0052 (7)	0.0020 (8)
C4	0.0624 (11)	0.0786 (13)	0.0398 (9)	-0.0077 (9)	0.0087 (7)	0.0087 (9)
C5	0.0576 (10)	0.0685 (12)	0.0437 (9)	-0.0113 (8)	0.0072 (7)	0.0044 (8)
C6	0.0400 (8)	0.0644 (10)	0.0362 (8)	-0.0036 (7)	0.0045 (6)	0.0022 (7)
C7	0.0440 (9)	0.0581 (10)	0.0375 (8)	-0.0006 (7)	0.0041 (6)	0.0026 (7)
C8	0.0458 (9)	0.0565 (10)	0.0327 (8)	-0.0012 (7)	0.0045 (6)	0.0001 (7)
C9	0.0446 (9)	0.0598 (10)	0.0354 (8)	-0.0022 (7)	0.0066 (6)	0.0002 (7)
C10	0.0370 (8)	0.0613 (10)	0.0379 (8)	-0.0033 (7)	0.0073 (6)	-0.0039 (7)
C11	0.0462 (9)	0.0639 (10)	0.0314 (7)	-0.0036 (7)	0.0042 (6)	0.0003 (7)
C12	0.0416 (9)	0.0690 (12)	0.0627 (11)	-0.0015 (8)	0.0105 (7)	-0.0104 (9)
C13	0.0471 (10)	0.0662 (12)	0.0749 (12)	-0.0092 (8)	0.0048 (8)	-0.0065 (9)
C14	0.0564 (10)	0.0553 (10)	0.0480 (9)	0.0004 (8)	-0.0001 (7)	-0.0010 (8)
C15	0.0509 (10)	0.0713 (12)	0.0579 (10)	0.0069 (9)	0.0070 (8)	-0.0058 (9)
C16	0.0424 (9)	0.0687 (12)	0.0541 (10)	-0.0057 (8)	0.0080 (7)	-0.0070 (8)
C17	0.0456 (9)	0.0572 (10)	0.0365 (8)	-0.0026 (7)	0.0056 (6)	0.0022 (7)
C18	0.0574 (10)	0.0554 (10)	0.0498 (9)	0.0041 (8)	-0.0032 (7)	0.0042 (8)
C19	0.0582 (10)	0.0671 (12)	0.0549 (10)	-0.0007 (9)	-0.0119 (8)	0.0001 (8)
C20	0.0553 (10)	0.0579 (11)	0.0532 (10)	-0.0032 (8)	0.0053 (8)	-0.0036 (8)
C21	0.0607 (11)	0.0566 (10)	0.0483 (9)	0.0021 (8)	0.0094 (8)	0.0072 (8)
C22	0.0498 (9)	0.0648 (11)	0.0414 (8)	0.0004 (8)	0.0009 (7)	0.0076 (8)
C23	0.0904 (17)	0.0707 (14)	0.123 (2)	-0.0214 (12)	-0.0012 (14)	-0.0209 (13)
C24	0.1026 (18)	0.0595 (13)	0.1068 (18)	-0.0005 (12)	0.0071 (13)	-0.0168 (12)
C25	0.0465 (12)	0.0525 (15)	0.0421 (12)	0.000	0.0088 (9)	0.000

Geometric parameters (Å, °)

O1—C14	1.370 (2)	C12—C13	1.386 (2)
O1—C23	1.410 (3)	C12—H12	0.9300
O2—C20	1.372 (2)	C13—C14	1.373 (2)
O2—C24	1.392 (2)	C13—H13	0.9300
O3—C25	1.217 (3)	C14—C15	1.380 (2)
C1—C2	1.417 (2)	C15—C16	1.370 (2)
C1—C6	1.424 (2)	C15—H15	0.9300
C1—C10	1.435 (2)	C16—H16	0.9300
C2—C3	1.366 (2)	C17—C22	1.387 (2)
C2—H2	0.9300	C17—C18	1.400 (2)
C3—C4	1.390 (3)	C18—C19	1.364 (2)
C3—H3	0.9300	C18—H18	0.9300
C4—C5	1.363 (2)	C19—C20	1.385 (2)
C4—H4	0.9300	C19—H19	0.9300
C5—C6	1.417 (2)	C20—C21	1.369 (2)
C5—H5	0.9300	C21—C22	1.383 (2)
C6—C7	1.447 (2)	C21—H21	0.9300
C7—C8	1.375 (2)	C22—H22	0.9300
C7—C17	1.484 (2)	C23—H23A	0.9600
C8—C9	1.427 (2)	C23—H23B	0.9600
C8—C8 ⁱ	1.496 (3)	C23—H23C	0.9600
C9—C10	1.372 (2)	C24—H24A	0.9600
C9—C25	1.486 (2)	C24—H24B	0.9600
C10—C11	1.486 (2)	C24—H24C	0.9600
C11—C12	1.372 (2)	C25—C9 ⁱ	1.486 (2)
C11—C16	1.385 (2)		
C14—O1—C23	117.79 (16)	O1—C14—C15	116.43 (16)
C20—O2—C24	118.67 (16)	C13—C14—C15	119.34 (16)
C2—C1—C6	118.69 (14)	C16—C15—C14	120.23 (16)
C2—C1—C10	121.01 (15)	C16—C15—H15	119.9
C6—C1—C10	120.29 (13)	C14—C15—H15	119.9
C3—C2—C1	121.29 (16)	C15—C16—C11	121.63 (16)
C3—C2—H2	119.4	C15—C16—H16	119.2
C1—C2—H2	119.4	C11—C16—H16	119.2
C2—C3—C4	120.01 (15)	C22—C17—C18	116.85 (15)
C2—C3—H3	120.0	C22—C17—C7	125.30 (14)
C4—C3—H3	120.0	C18—C17—C7	117.72 (14)
C5—C4—C3	120.45 (16)	C19—C18—C17	121.63 (16)
C5—C4—H4	119.8	C19—C18—H18	119.2
C3—C4—H4	119.8	C17—C18—H18	119.2
C4—C5—C6	121.66 (17)	C18—C19—C20	120.08 (15)
C4—C5—H5	119.2	C18—C19—H19	120.0
C6—C5—H5	119.2	C20—C19—H19	120.0
C5—C6—C1	117.69 (14)	C21—C20—O2	124.73 (16)
C5—C6—C7	121.72 (15)	C21—C20—C19	119.94 (16)

C1—C6—C7	120.59 (14)	O2—C20—C19	115.32 (15)
C8—C7—C6	117.10 (14)	C20—C21—C22	119.56 (16)
C8—C7—C17	122.09 (13)	C20—C21—H21	120.2
C6—C7—C17	120.16 (14)	C22—C21—H21	120.2
C7—C8—C9	120.74 (13)	C21—C22—C17	121.93 (15)
C7—C8—C8 ⁱ	131.26 (10)	C21—C22—H22	119.0
C9—C8—C8 ⁱ	107.44 (8)	C17—C22—H22	119.0
C10—C9—C8	123.01 (14)	O1—C23—H23A	109.5
C10—C9—C25	128.04 (15)	O1—C23—H23B	109.5
C8—C9—C25	108.43 (13)	H23A—C23—H23B	109.5
C9—C10—C1	116.85 (14)	O1—C23—H23C	109.5
C9—C10—C11	122.49 (14)	H23A—C23—H23C	109.5
C1—C10—C11	120.66 (13)	H23B—C23—H23C	109.5
C12—C11—C16	117.21 (16)	O2—C24—H24A	109.5
C12—C11—C10	121.07 (15)	O2—C24—H24B	109.5
C16—C11—C10	121.68 (14)	H24A—C24—H24B	109.5
C11—C12—C13	122.07 (16)	O2—C24—H24C	109.5
C11—C12—H12	119.0	H24A—C24—H24C	109.5
C13—C12—H12	119.0	H24B—C24—H24C	109.5
C14—C13—C12	119.50 (16)	O3—C25—C9	126.89 (10)
C14—C13—H13	120.2	O3—C25—C9 ⁱ	126.89 (10)
C12—C13—H13	120.2	C9—C25—C9 ⁱ	106.22 (19)
O1—C14—C13	124.23 (16)		
C6—C1—C2—C3	1.6 (2)	C1—C10—C11—C16	70.8 (2)
C10—C1—C2—C3	-176.88 (14)	C16—C11—C12—C13	-1.2 (3)
C1—C2—C3—C4	2.1 (3)	C10—C11—C12—C13	176.72 (16)
C2—C3—C4—C5	-2.5 (3)	C11—C12—C13—C14	0.0 (3)
C3—C4—C5—C6	-1.0 (3)	C23—O1—C14—C13	5.4 (3)
C4—C5—C6—C1	4.6 (2)	C23—O1—C14—C15	-174.72 (18)
C4—C5—C6—C7	-175.27 (16)	C12—C13—C14—O1	-178.90 (16)
C2—C1—C6—C5	-4.9 (2)	C12—C13—C14—C15	1.2 (3)
C10—C1—C6—C5	173.64 (14)	O1—C14—C15—C16	178.89 (16)
C2—C1—C6—C7	175.05 (14)	C13—C14—C15—C16	-1.2 (3)
C10—C1—C6—C7	-6.4 (2)	C14—C15—C16—C11	0.0 (3)
C5—C6—C7—C8	176.60 (15)	C12—C11—C16—C15	1.2 (2)
C1—C6—C7—C8	-3.3 (2)	C10—C11—C16—C15	-176.70 (15)
C5—C6—C7—C17	-12.5 (2)	C8—C7—C17—C22	-128.22 (17)
C1—C6—C7—C17	167.62 (13)	C6—C7—C17—C22	61.3 (2)
C6—C7—C8—C9	12.1 (2)	C8—C7—C17—C18	56.0 (2)
C17—C7—C8—C9	-158.68 (14)	C6—C7—C17—C18	-114.44 (17)
C6—C7—C8—C8 ⁱ	-177.58 (19)	C22—C17—C18—C19	-1.2 (3)
C17—C7—C8—C8 ⁱ	11.7 (3)	C7—C17—C18—C19	174.98 (16)
C7—C8—C9—C10	-11.7 (2)	C17—C18—C19—C20	0.6 (3)
C8 ⁱ —C8—C9—C10	175.83 (15)	C24—O2—C20—C21	3.5 (3)
C7—C8—C9—C25	160.57 (13)	C24—O2—C20—C19	-177.98 (18)
C8 ⁱ —C8—C9—C25	-11.85 (19)	C18—C19—C20—C21	0.2 (3)
C8—C9—C10—C1	1.6 (2)	C18—C19—C20—O2	-178.36 (16)

C25—C9—C10—C1	-169.11 (12)	O2—C20—C21—C22	178.03 (16)
C8—C9—C10—C11	-178.39 (14)	C19—C20—C21—C22	-0.4 (3)
C25—C9—C10—C11	10.9 (2)	C20—C21—C22—C17	-0.2 (3)
C2—C1—C10—C9	-174.37 (14)	C18—C17—C22—C21	1.0 (2)
C6—C1—C10—C9	7.2 (2)	C7—C17—C22—C21	-174.84 (15)
C2—C1—C10—C11	5.6 (2)	C10—C9—C25—O3	-3.49 (18)
C6—C1—C10—C11	-172.82 (13)	C8—C9—C25—O3	-175.31 (7)
C9—C10—C11—C12	73.0 (2)	C10—C9—C25—C9 ⁱ	176.51 (18)
C1—C10—C11—C12	-107.00 (18)	C8—C9—C25—C9 ⁱ	4.69 (7)
C9—C10—C11—C16	-109.20 (17)		

Symmetry code: (i) $-x+3/2, y, -z+1/2$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C24—H24c \cdots O3 ⁱⁱ	0.96	2.39	3.199 (3)	141

Symmetry code: (ii) $x, y+1, z$.