

Crystal structure of (Z)-2,3-dichloro-1,4-bis(4-methoxyphenyl)but-2-ene-1,4-dione

Ram K. Tittal,^{a*‡} Satish Kumar^b and R. N. Ram^a

^aDepartment of Chemistry, Indian Institute of Technology Delhi, Hauz Khas, New Delhi 110 016, India, and ^bDepartment of Chemistry, St Stephen's College, University Enclave, Delhi 110 007, India. *Correspondence e-mail: rktittaliitd@nitkkr.ac.in

Received 4 July 2014; accepted 19 August 2014

Edited by L. Fabian, University of East Anglia, England

The title compound, C₁₈H₁₄Cl₂O₄, adopts a *Z* conformation around the central C=C bond. The two aromatic rings of the molecule are nearly perpendicular to each other, with a dihedral angle between of 86.22 (14)°. The methoxy substituents lie close to the plane of the attached benzene rings. The C(ar)–C(ar)–O–C(Me) torsion angles are –2.4 (7) and 7.5 (6)°. Weak C–H···O interactions link the molecules forming a three-dimensional network. The crystal packing also displays short [3.160 (3) Å] Cl···O halogen-bonding contacts between molecules related by the screw axis. The structure exhibits disorder of one carbonyl O atom with a refined occupancy ratio of 0.21 (6):0.79 (6).

Keywords: crystal structure; 2,3-dichlorobut-2-ene-1,4-dione; dichloromethyl radical; CuCl/bpy; trichloromethyl groups; stereoselectivity.

CCDC reference: 1019996

1. Related literature

For a review of radical reactions, see: Clark (2002); Ram & Tittal (2014a,b); Pintauer & Matyjaszewski (2008). For details of the synthesis, see: Kurosawa & Yamaguchi (1981); Ram *et al.* (2007). For halogen-bond interactions, see: Agarwal *et al.* (2014). For a similar structure and short aromatic contacts, see: Tittal *et al.* (2014).

2. Experimental

2.1. Crystal data

C ₁₈ H ₁₄ Cl ₂ O ₄	<i>V</i> = 848.3 (5) Å ³
<i>M_r</i> = 365.19	<i>Z</i> = 2
Monoclinic, <i>P</i> 2 ₁	Mo <i>K</i> α radiation
<i>a</i> = 8.877 (3) Å	<i>μ</i> = 0.40 mm ^{−1}
<i>b</i> = 9.914 (3) Å	<i>T</i> = 273 K
<i>c</i> = 10.294 (3) Å	0.31 × 0.23 × 0.14 mm
<i>β</i> = 110.565 (5)°	

2.2. Data collection

Bruker SMART APEX CCD diffractometer	9879 measured reflections
Absorption correction: multi-scan (SADABS; Bruker, 2000)	4012 independent reflections
<i>T</i> _{min} = 0.860, <i>T</i> _{max} = 1.000	3755 reflections with <i>I</i> > 2σ(<i>I</i>)
	<i>R</i> _{int} = 0.025

2.3. Refinement

<i>R</i> [<i>F</i> ² > 2σ(<i>F</i> ²)] = 0.057	<i>Δρ</i> _{min} = −0.17 e Å ^{−3}
<i>wR</i> (<i>F</i> ²) = 0.134	Absolute structure: Flack <i>x</i>
<i>S</i> = 1.17	determined using 1548 quotients
4012 reflections	[(<i>I</i> ⁺ − <i>I</i> [−])]/[(<i>I</i> ⁺ + <i>I</i> [−])] (Parsons <i>et al.</i> , 2013)
223 parameters	Absolute structure parameter:
2 restraints	0.01 (2)
H-atom parameters constrained	
<i>Δρ</i> _{max} = 0.41 e Å ^{−3}	

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> –H··· <i>A</i>	<i>D</i> –H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> –H··· <i>A</i>
C7–H7···O4A ⁱ	0.93	2.49	3.23 (3)	137
C7–H7···O4B ⁱ	0.93	2.62	3.245 (9)	125
C12–H12···O1 ⁱⁱ	0.93	2.66	3.488 (5)	149
C14–H14···O1 ⁱⁱⁱ	0.93	2.72	3.363 (5)	127

Symmetry codes: (i) $-x + 1, y - \frac{1}{2}, -z + 1$; (ii) $-x, y - \frac{1}{2}, -z$; (iii) $x + 1, y - 1, z$.

Data collection: SMART (Bruker, 2000); cell refinement: SAINT (Bruker, 2000); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: OLEX2 (Dolomanov *et al.*, 2009) and Mercury (Macrae *et al.*, 2008); software used to prepare material for publication: OLEX2, PLATON (Spek, 2009) and publCIF (Westrip, 2010).

Acknowledgements

The authors are thankful to Shailesh Upreti for providing assistance in solving the X-ray structure.

Supporting information for this paper is available from the IUCr electronic archives (Reference: FY2117).

[‡] Present address: Department of Chemistry, National Institute of Technology, Kurukshetra, Haryana 136 119, India.

References

- Agarwal, P., Mishra, P., Gupta, N., Neelam, Sahoo, P. & Kumar, S. (2014). *Acta Cryst.* **E70**, o418.
- Bruker (2000). *SMART, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Clark, A. J. (2002). *Chem. Soc. Rev.* **31**, 1–11.
- Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
- Kurosawa, K. & Yamaguchi, K. (1981). *Bull. Chem. Soc. Jpn.* **54**, 1757–1760.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Parsons, S., Flack, H. D. & Wagner, T. (2013). *Acta Cryst.* **B69**, 249–259.
- Pintauer, T. & Matyjaszewski, K. (2008). *Chem. Soc. Rev.* **37**, 1087–1097.
- Ram, R. N. & Tittal, R. K. (2014a). *Tetrahedron Lett.* **55**, 4448–4451.
- Ram, R. N. & Tittal, R. K. (2014b). *Tetrahedron Lett.* **55**, 4342–4345.
- Ram, R. N., Tittal, R. K. & Upreti, S. (2007). *Tetrahedron Lett.* **48**, 7994–7997.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Tittal, R. K., Kumar, S. & Ram, R. N. (2014). *Acta Cryst.* **E70**, o861–o862.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2014). E70, o1049–o1050 [doi:10.1107/S1600536814018790]

Crystal structure of (Z)-2,3-dichloro-1,4-bis(4-methoxyphenyl)but-2-ene-1,4-dione

Ram K. Tittal, Satish Kumar and R. N. Ram

S1. Chemical context

Trichloromethyl groups are known to generate free radicals through homolysis of a C—Cl bond with relative ease in presence of radical initiators or UV-light or redox active metal salts (such as CuCl) or its complexes. A number of research papers are available that show radical intermediates produced from trichloromethyl group containing compounds. For example, CuCl or its complexes with bipyridine or bi- or tridentate amine ligands have been used with trichloromethyl groups on a variety of organic substrates under non reducing conditions to generate free radicals. Such radicals have been reported to undergo intramolecular and intermolecular cyclization or addition reactions along with mono- and/or-di reduction products (Clark, 2002; Ram *et al.*, 2007). The radicals produced in these reactions can also acts as a radical initiator in ATRP (Pintauer & Matyjaszewski, 2008). However, the presence of relatively better leaving group at the β -position of the radical centre leads to predominantly rearrangement and/or fragmentation reactions through the intermediate formation of contact ion pairs (Ram & Tittal, 2014*a,b*). It is important to mention here that 2,2,2-trichloroethylalkyl ethers and trichloromethyl carbinols with no leaving group at β -position to the trichloromethyl around carbon carbon double bond undergoes 1,2-H shift under similar conditions through the intermediate copper-carbenoid species (Ram & Tittal, 2014*b*). Keeping in view the above discussion, we have decided to study the behavior of the radicals produced from substituted trichloromethyl compounds with no suitably located carbon-carbon double bond and leaving group or any hydrogen atom at the β -position of the radical centre in order to restrict the intermolecular or intramolecular addition; ATRP; rearrangement and/or fragmentation or 1,2-H shift. The major product obtained under such conditions is reported here.

S2. Structural commentary

In the asymmetric unit (Fig. 1) the molecule adopts *Z* conformation about the C=C bond: C8=C9. The two aromatic rings are nearly perpendicular with an interplanar angle between the two phenyl rings of 86.22 (14)°. The two methoxy substituents lie close to the plane of the attached benzene rings. The C(ar)—C(ar)—O—C(Me) torsion angles are -2.4 (7)° (C21—C4—O1—C22) and 7.5 (6)° (C14—C9—O2—C20). The structure exhibits disorder of carbonyl group with a refined occupancy ratio of 0.21 (6): 0.79 (6).

S3. Supramolecular features

The crystal packing is stabilized by short halogen bond Cl \cdots O interactions (Fig. 2) [3.160 (3)Å]. The crystal structure of also displays intermolecular C(ar)—H \cdots O (Table 1, Fig. 3) short contacts.

S4. Synthesis and crystallization

A two-neck round bottom flask fitted with a rubber septum was charged with CuCl (0.8 g, 0.008 mol), 2,2'-bipyridine (bpy, 1.25 g, 0.008 mol) under continuous flow of nitrogen followed by addition of 1 mL dry dichloroethane (DCE) or benzene to ensure CuCl-bpy complex formation. To this reaction flask a solution of the 2,2,2-trichloro-1-(4-methoxyphenyl)-ethanone (0.004 mol) in dry DCE or benzene (5 mL) was injected through the septum with the help of a syringe and the reaction mixture was heated at reflux with stirring under a slow and continuous flow of nitrogen. The completion of the reaction was indicated by TLC (1-2 h). The reaction mixture was cooled and filtered through a celite pad. The filtrate was evaporated under reduced pressure and purified on a silica gel column chromatography using *n*-hexane and ethylacetate mixture as the solvent to get title compound in 60 or 70% isolated yields in DCE or benzene, respectively. Suitable single crystals were obtained from a chloroform:hexane (3:7) mixture by slow evaporation at room temperature. Melting point 132°C, literature melting point 132-134°C (Kurosawa & Yamaguchi, 1981).

S5. Refinement

H atoms were placed in calculated positions and refined using riding model, with C—H 0.93 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for aromatic C—H. Methyl groups were refined as idealised rotating groups, with C—H = 0.96 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$. Disorder was modeled for O4 atom of the carbonyl group over two sets of sites with minor:major occupancy ratio of 0.21 (6): 0.79 (6). Similarity restraints were used for the C=O bond distances using SADI. Anisotropic displacement parameters of the minor component of oxygen atom O4 were constrained to those of the major component using EADP.




Figure 1

Molecular structure of the title compound, with atom labels and 50% probability displacement ellipsoids for non-H atoms.




Figure 2

A view of the crystal packing of the title compound. A weak C—H...O hydrogen bond is shown as a dashed line.




Figure 3

Part of the structure of the title compound showing O...Cl and C—H...O interactions.

(Z)-2,3-Dichloro-1,4-bis(4-methoxyphenyl)but-2-ene-1,4-dione

Crystal data

$C_{18}H_{14}Cl_2O_4$

$M_r = 365.19$

Monoclinic, $P2_1$

$a = 8.877(3) \text{ \AA}$

$b = 9.914(3) \text{ \AA}$

$c = 10.294(3) \text{ \AA}$

$\beta = 110.565(5)^\circ$

$V = 848.3(5) \text{ \AA}^3$

$Z = 2$

$F(000) = 376$

$D_x = 1.430 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 5754 reflections

$\theta = 3.2\text{--}26.1^\circ$

$\mu = 0.40 \text{ mm}^{-1}$

$T = 273 \text{ K}$

Block, colorless

$0.31 \times 0.23 \times 0.14 \text{ mm}$

Data collection

Bruker SMART APEX CCD
diffractometer

Radiation source: fine-focus sealed tube

φ and ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 2000)

$T_{\min} = 0.860$, $T_{\max} = 1.000$

9879 measured reflections

4012 independent reflections

3755 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.025$

$\theta_{\max} = 28.3^\circ$, $\theta_{\min} = 2.1^\circ$

$h = -11 \rightarrow 11$

$k = -12 \rightarrow 13$

$l = -13 \rightarrow 13$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.057$

$wR(F^2) = 0.134$

$S = 1.17$

4012 reflections

223 parameters

2 restraints

Primary atom site location: structure-invariant
direct methods

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0741P)^2 + 0.0555P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.41 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.17 \text{ e } \text{\AA}^{-3}$

Absolute structure: Flack x determined using

1548 quotients $[(I^+)-(I^-)]/[(I^+)+(I^-)]$ (Parsons *et al.*, 2013)

Absolute structure parameter: 0.01 (2)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C1	0.6776 (4)	0.5542 (4)	0.2160 (4)	0.0447 (8)	
C2	0.5707 (5)	0.6341 (4)	0.1270 (4)	0.0443 (8)	
C3	0.6859 (4)	0.3333 (4)	0.3536 (4)	0.0424 (7)	
C4	0.1894 (4)	1.0044 (4)	0.1824 (4)	0.0456 (8)	
C6	0.3256 (4)	0.7652 (4)	0.1360 (4)	0.0420 (7)	
C7	0.7162 (5)	0.1437 (5)	0.5065 (4)	0.0505 (9)	
H7	0.7053	0.1059	0.5854	0.061*	
C8	0.7473 (4)	0.2551 (4)	0.2713 (3)	0.0444 (8)	
H8	0.7567	0.2925	0.1916	0.053*	
C9	0.7798 (4)	0.0673 (4)	0.4242 (4)	0.0449 (8)	
C11	0.0916 (5)	0.9011 (5)	0.1102 (5)	0.0544 (10)	
H11	-0.0196	0.9119	0.0769	0.065*	
C12	0.1581 (4)	0.7821 (4)	0.0872 (4)	0.0507 (9)	
H12	0.0916	0.7125	0.0391	0.061*	

C13	0.6700 (5)	0.2740 (4)	0.4710 (4)	0.0494 (9)	
H13	0.6271	0.3242	0.5261	0.059*	
C14	0.7946 (5)	0.1233 (4)	0.3056 (4)	0.0463 (8)	
H14	0.8360	0.0725	0.2499	0.056*	
C15	0.6357 (5)	0.4749 (4)	0.3229 (4)	0.0535 (9)	
O4A	0.516 (5)	0.509 (5)	0.347 (7)	0.088 (5)	0.20 (6)
O4B	0.569 (3)	0.5385 (17)	0.390 (2)	0.088 (5)	0.80 (6)
C16	0.3938 (5)	0.6362 (4)	0.1112 (5)	0.0496 (9)	
C18	0.4208 (4)	0.8699 (4)	0.2085 (4)	0.0463 (8)	
H18	0.5320	0.8598	0.2417	0.056*	
C20	0.9065 (6)	-0.1371 (5)	0.3997 (6)	0.0724 (14)	
H20A	0.8372	-0.1526	0.3055	0.109*	
H20B	0.9380	-0.2221	0.4460	0.109*	
H20C	1.0004	-0.0888	0.4003	0.109*	
C21	0.3548 (4)	0.9895 (4)	0.2329 (4)	0.0456 (8)	
H21	0.4208	1.0587	0.2826	0.055*	
C22	0.2043 (6)	1.2268 (6)	0.2784 (6)	0.0742 (14)	
H22A	0.2795	1.2571	0.2367	0.111*	
H22B	0.2617	1.1972	0.3715	0.111*	
H22C	0.1337	1.2996	0.2800	0.111*	
O1	0.1126 (4)	1.1181 (3)	0.2000 (4)	0.0633 (8)	
O2	0.8234 (4)	-0.0603 (3)	0.4692 (3)	0.0620 (8)	
O3	0.3164 (4)	0.5356 (4)	0.0744 (6)	0.0929 (14)	
Cl1	0.62060 (13)	0.74187 (12)	0.01732 (11)	0.0625 (3)	
Cl2	0.87618 (12)	0.54992 (12)	0.22528 (13)	0.0651 (3)	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0420 (17)	0.0381 (17)	0.0560 (19)	0.0057 (16)	0.0198 (15)	-0.0067 (16)
C2	0.0487 (19)	0.0387 (18)	0.0471 (19)	-0.0001 (15)	0.0186 (16)	-0.0032 (14)
C3	0.0382 (16)	0.0429 (19)	0.0446 (17)	0.0046 (15)	0.0126 (14)	-0.0020 (14)
C4	0.0460 (18)	0.0446 (19)	0.048 (2)	0.0078 (15)	0.0193 (16)	0.0042 (15)
C6	0.0404 (17)	0.0397 (19)	0.0452 (17)	0.0031 (15)	0.0141 (13)	0.0032 (14)
C7	0.052 (2)	0.053 (2)	0.049 (2)	-0.0062 (18)	0.0214 (17)	0.0025 (17)
C8	0.0446 (18)	0.048 (2)	0.0401 (16)	0.0033 (17)	0.0146 (14)	-0.0021 (15)
C9	0.0390 (16)	0.0403 (19)	0.0507 (19)	-0.0057 (15)	0.0098 (15)	-0.0021 (15)
C11	0.0343 (18)	0.061 (3)	0.064 (2)	0.0039 (17)	0.0122 (17)	-0.0033 (19)
C12	0.0400 (18)	0.050 (2)	0.058 (2)	-0.0061 (16)	0.0123 (16)	-0.0080 (17)
C13	0.050 (2)	0.052 (2)	0.052 (2)	0.0006 (17)	0.0252 (16)	-0.0062 (17)
C14	0.0454 (19)	0.0433 (19)	0.0496 (19)	0.0016 (16)	0.0161 (16)	-0.0090 (16)
C15	0.060 (2)	0.049 (2)	0.056 (2)	0.0144 (19)	0.0261 (19)	0.0014 (18)
O4A	0.131 (10)	0.066 (5)	0.099 (8)	0.046 (6)	0.079 (8)	0.018 (5)
O4B	0.131 (10)	0.066 (5)	0.099 (8)	0.046 (6)	0.079 (8)	0.018 (5)
C16	0.0439 (19)	0.0402 (19)	0.061 (2)	0.0017 (16)	0.0140 (17)	-0.0052 (17)
C18	0.0321 (16)	0.047 (2)	0.055 (2)	0.0029 (15)	0.0091 (15)	0.0002 (16)
C20	0.073 (3)	0.046 (2)	0.086 (3)	0.019 (2)	0.013 (2)	-0.001 (2)
C21	0.0411 (18)	0.0391 (18)	0.053 (2)	-0.0010 (15)	0.0124 (16)	-0.0070 (15)

C22	0.074 (3)	0.060 (3)	0.089 (3)	0.014 (3)	0.029 (3)	-0.018 (3)
O1	0.0521 (16)	0.0538 (19)	0.083 (2)	0.0161 (14)	0.0219 (15)	-0.0103 (16)
O2	0.072 (2)	0.0416 (16)	0.0703 (19)	0.0056 (14)	0.0220 (15)	0.0056 (13)
O3	0.0590 (18)	0.050 (2)	0.166 (4)	-0.0067 (17)	0.035 (2)	-0.032 (2)
C11	0.0694 (6)	0.0620 (6)	0.0619 (6)	0.0065 (6)	0.0302 (5)	0.0126 (5)
C12	0.0461 (5)	0.0549 (6)	0.0968 (8)	0.0105 (5)	0.0284 (5)	0.0012 (6)

Geometric parameters (Å, °)

C1—C2	1.326 (5)	C9—O2	1.356 (5)
C1—C15	1.502 (6)	C11—H11	0.9300
C1—C12	1.732 (3)	C11—C12	1.377 (6)
C2—C16	1.521 (5)	C12—H12	0.9300
C2—C11	1.722 (4)	C13—H13	0.9300
C3—C8	1.393 (5)	C14—H14	0.9300
C3—C13	1.395 (6)	C15—O4A	1.219 (13)
C3—C15	1.473 (6)	C15—O4B	1.229 (8)
C4—C11	1.379 (6)	C16—O3	1.195 (5)
C4—C21	1.383 (5)	C18—H18	0.9300
C4—O1	1.362 (5)	C18—C21	1.384 (6)
C6—C12	1.402 (5)	C20—H20A	0.9600
C6—C16	1.476 (5)	C20—H20B	0.9600
C6—C18	1.380 (5)	C20—H20C	0.9600
C7—H7	0.9300	C20—O2	1.417 (6)
C7—C9	1.395 (6)	C21—H21	0.9300
C7—C13	1.367 (6)	C22—H22A	0.9600
C8—H8	0.9300	C22—H22B	0.9600
C8—C14	1.380 (6)	C22—H22C	0.9600
C9—C14	1.389 (6)	C22—O1	1.419 (6)
C2—C1—C15	121.2 (3)	C7—C13—H13	119.4
C2—C1—C12	121.4 (3)	C8—C14—C9	119.3 (3)
C15—C1—C12	117.1 (3)	C8—C14—H14	120.4
C1—C2—C16	122.9 (4)	C9—C14—H14	120.4
C1—C2—C11	122.5 (3)	C3—C15—C1	121.4 (3)
C16—C2—C11	114.6 (3)	O4A—C15—C1	117 (2)
C8—C3—C13	118.3 (3)	O4A—C15—C3	116 (2)
C8—C3—C15	123.6 (4)	O4B—C15—C1	115.8 (6)
C13—C3—C15	118.2 (3)	O4B—C15—C3	122.5 (6)
C11—C4—C21	120.5 (4)	C6—C16—C2	117.7 (3)
O1—C4—C11	115.9 (3)	O3—C16—C2	118.7 (4)
O1—C4—C21	123.6 (4)	O3—C16—C6	123.6 (4)
C12—C6—C16	119.3 (3)	C6—C18—H18	119.2
C18—C6—C12	118.4 (3)	C6—C18—C21	121.6 (3)
C18—C6—C16	122.3 (3)	C21—C18—H18	119.2
C9—C7—H7	120.0	H20A—C20—H20B	109.5
C13—C7—H7	120.0	H20A—C20—H20C	109.5
C13—C7—C9	119.9 (4)	H20B—C20—H20C	109.5

C3—C8—H8	119.3	O2—C20—H20A	109.5
C14—C8—C3	121.3 (4)	O2—C20—H20B	109.5
C14—C8—H8	119.3	O2—C20—H20C	109.5
C14—C9—C7	120.0 (4)	C4—C21—C18	119.0 (3)
O2—C9—C7	115.3 (3)	C4—C21—H21	120.5
O2—C9—C14	124.7 (4)	C18—C21—H21	120.5
C4—C11—H11	119.9	H22A—C22—H22B	109.5
C12—C11—C4	120.1 (3)	H22A—C22—H22C	109.5
C12—C11—H11	119.9	H22B—C22—H22C	109.5
C6—C12—H12	119.8	O1—C22—H22A	109.5
C11—C12—C6	120.4 (4)	O1—C22—H22B	109.5
C11—C12—H12	119.8	O1—C22—H22C	109.5
C3—C13—H13	119.4	C4—O1—C22	119.3 (3)
C7—C13—C3	121.2 (4)	C9—O2—C20	117.7 (4)
C1—C2—C16—C6	121.1 (4)	C13—C7—C9—C14	-0.6 (6)
C1—C2—C16—O3	-61.8 (6)	C13—C7—C9—O2	179.4 (4)
C2—C1—C15—C3	137.3 (4)	C14—C9—O2—C20	7.5 (6)
C2—C1—C15—O4A	-17 (4)	C15—C1—C2—C16	-9.4 (6)
C2—C1—C15—O4B	-48.1 (18)	C15—C1—C2—C11	171.9 (3)
C3—C8—C14—C9	0.3 (6)	C15—C3—C8—C14	179.1 (4)
C4—C11—C12—C6	-0.5 (7)	C15—C3—C13—C7	-179.1 (4)
C6—C18—C21—C4	-0.5 (6)	C16—C6—C12—C11	179.4 (4)
C7—C9—C14—C8	0.6 (6)	C16—C6—C18—C21	-178.9 (4)
C7—C9—O2—C20	-172.5 (4)	C18—C6—C12—C11	0.7 (6)
C8—C3—C13—C7	1.3 (6)	C18—C6—C16—C2	-18.6 (6)
C8—C3—C15—C1	-12.4 (6)	C18—C6—C16—O3	164.4 (5)
C8—C3—C15—O4A	142 (4)	C21—C4—C11—C12	-0.3 (7)
C8—C3—C15—O4B	173.5 (19)	C21—C4—O1—C22	-2.4 (7)
C9—C7—C13—C3	-0.4 (6)	O1—C4—C11—C12	-180.0 (4)
C11—C4—C21—C18	0.8 (6)	O1—C4—C21—C18	-179.6 (4)
C11—C4—O1—C22	177.3 (4)	O2—C9—C14—C8	-179.4 (3)
C12—C6—C16—C2	162.7 (4)	C11—C2—C16—C6	-60.1 (4)
C12—C6—C16—O3	-14.2 (7)	C11—C2—C16—O3	117.0 (5)
C12—C6—C18—C21	-0.2 (6)	C12—C1—C2—C16	177.2 (3)
C13—C3—C8—C14	-1.3 (5)	C12—C1—C2—C11	-1.5 (5)
C13—C3—C15—C1	168.0 (4)	C12—C1—C15—C3	-49.0 (5)
C13—C3—C15—O4A	-38 (4)	C12—C1—C15—O4A	157 (4)
C13—C3—C15—O4B	-6.1 (19)	C12—C1—C15—O4B	125.5 (17)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C7—H7...O4A ⁱ	0.93	2.49	3.23 (3)	137
C7—H7...O4B ⁱ	0.93	2.62	3.245 (9)	125

C12—H12···O1 ⁱⁱ	0.93	2.66	3.488 (5)	149
C14—H14···O1 ⁱⁱⁱ	0.93	2.72	3.363 (5)	127

Symmetry codes: (i) $-x+1, y-1/2, -z+1$; (ii) $-x, y-1/2, -z$; (iii) $x+1, y-1, z$.