

# Crystal structure of methyl 2-[2,4-bis(4-fluorophenyl)-3-azabicyclo[3.3.1]nonan-9-ylidene]-hydrazinecarboxylate

A. Kamaraj,<sup>a</sup> R. Rajkumar,<sup>a</sup> K. Krishnasamy<sup>a\*</sup> and S. Murugavel<sup>b\*</sup>

Received 14 August 2014

Accepted 21 August 2014

Edited by H. Stoeckli-Evans, University of Neuchâtel, Switzerland

**Keywords:** crystal structure; 3-azabicyclo[3.3.1]nonane; hydrazinecarboxylate; twin-chair conformation

**CCDC reference:** 1020373

**Supporting information:** this article has supporting information at journals.iucr.org/e



<sup>a</sup>Department of Chemistry, Annamalai University, Annamalaiagar 608 002, Chidambaram, Tamilnadu, India, and


<sup>b</sup>Department of Physics, Thanthai Periyar Government Institute of Technology, Vellore 632 002, India. \*Correspondence e-mail: krishnasamybala56@gmail.com, smurugavel27@gmail.com

In the title compound,  $C_{22}H_{23}F_2N_3O_2$ , the bicyclic ring system exists in a twin-chair conformation with an equatorial disposition of the 4-fluorophenyl groups on the heterocycle. These aromatic rings are inclined to one another by  $19.4(1)^\circ$ . In the crystal, molecules are linked by pairs of  $N-H\cdots O$  and  $C-H\cdots O$  hydrogen bonds into inversion dimers, incorporating  $R_1^2(7)$  and  $R_2^2(8)$  ring motifs; the same O atom accepts both hydrogen bonds. These dimers are further linked by a pair of  $C-H\cdots F$  hydrogen bonds, enclosing  $R_2^2(28)$  ring motifs, forming supramolecular chains along [010]. The NH group of the pyridine ring is not involved in hydrogen bonding, probably due to the steric hindrance of the fluorophenyl groups.

## 1. Chemical context

Molecules containing the 3-azabicyclo[3.3.1]nonane nucleus are of great interest due to their presence in a wide range of naturally occurring diterpenoid/norditerpenoid alkaloids and their broad-spectrum biological activities, such as anti-microbial, analgesic, antagonistic, anti-inflammatory and local anesthetic hypotensive activity (Parthiban *et al.*, 2009; Hardick *et al.*, 1996; Jeyaraman & Avila, 1981). Hence, the synthesis of new molecules with the 3-azabicyclo[3.3.1]nonane nucleus and their stereochemical investigation are of interest in the field of medicinal chemistry. Also, the stereochemistry of such molecules is a major criterium for their biological response. As a consequence, the present study was undertaken to examine the configuration and conformation of the synthesized title compound.





**Figure 1**  
The molecular structure of the title molecule (I), showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level.

## 2. Structural commentary

The molecular structure of the title compound, (I), is illustrated in Fig. 1. The bicyclo ring system adopts a twin-chair conformation, with puckering parameters  $Q = 0.593$  (2) Å,  $\theta = 170.8$  (2)° and  $\varphi = 353.9$  (1)° for the N1/C1–C5 piperidine ring, and  $Q = 0.546$  (2) Å,  $\theta = 10.9$  (2)° and  $\varphi = 65.0$  (1)° for the C2–C4/C6–C8 cyclohexane ring. The fluorophenyl groups on the heterocycle occupy equatorial positions and are inclined to one another by 19.4 (1)°. The geometric parameters of the title molecule agree well with those reported for similar structures, for example, 2,4-bis(4-fluorophenyl)-3-azabicyclo[3.3.1]nonan-9-one, (II) (Parthiban *et al.*, 2008), and 2,4-bis(4-fluorophenyl)-1,5-dimethyl-3-azabicyclo[3.3.1]nonan-9-one, (III) (Rizwana Begum *et al.*, 2013).

## 3. Supramolecular features

In the crystal, pairs of bifurcated acceptor N3–H3···O3<sup>i</sup> and C2–H2···O3<sup>i</sup> (Table 1) hydrogen bonds link molecules into inversion dimers, incorporating  $R_1^2(7)$  and  $R_2^2(8)$  ring motifs (Fig. 2). These dimers are further linked through a pair of C22–H22C···F1<sup>ii</sup> (Table 1) hydrogen bonds, enclosing  $R_2^2(28)$  ring motifs, forming supramolecular chains along the *b*-axis direction (Fig. 2).

The NH group of the pyridine ring is not involved in hydrogen bonding, probably due to the steric hindrance of the fluorophenyl groups. Such a situation was reported for a similar bicyclic system substituted by difluorophenyl rings, *viz.* compound (III) (Rizwana Begum *et al.*, 2013).


**Table 1**  
Hydrogen-bond geometry (Å, °).

<i>D</i> –H··· <i>A</i>	<i>D</i> –H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> –H··· <i>A</i>
N3–H3···O3 <sup>i</sup>	0.86	2.12	2.930 (2)	157
C2–H2···O3 <sup>i</sup>	0.98	2.49	3.442 (2)	163
C22–H22C···F1 <sup>ii</sup>	0.96	2.49	3.243 (3)	136

Symmetry codes: (i)  $-x + \frac{1}{2}, -y + \frac{5}{2}, -z$ ; (ii)  $-x + \frac{1}{2}, -y + \frac{3}{2}, -z$ .

## 4. Database survey

38 ‘hits’ for crystal structures containing the 3-azabicyclo[3.3.1]nonane subunit were obtained for a search of the Cambridge Structural Database (CSD, Version 5.35, last update of February 2014; Allen, 2002). However, extending the search to allow additional substitution of 4-fluorophenyl groups on the bicyclic ring gave two hits, namely compounds (II) and (III) mentioned above (Section 2). Compound (III) crystallized with two independent molecules (*A* and *B*) in the asymmetric unit. In all three compounds, the bicyclo rings have twin-chair conformations with equatorially disposed 4-fluorophenyl groups on the heterocycle. The fluorophenyl rings are oriented at an angle of 28.7 (1)° in (II), and 55.3 (1)° (molecule *A*) and 56.4 (1)° (molecule *B*) for (III), compared to 19.4 (1)° in the title compound, (I).



**Figure 2**  
Partial view of the crystal packing of the title compound, showing the hydrogen bonds as dashed lines (see Table 1 for details). H atoms not involved in hydrogen bonding have been omitted for clarity.

## 5. Synthesis and crystallization

A mixture of 2,4-diphenyl-3-azabicyclo[3.3.1]nonan-9-one (0.1 mmol), methyl hydrazinecarboxylate (1.5 mmol) in an ethanol–chloroform (1:1 *v/v*) medium, with the addition of few drops of acetic acid, was stirred for 10–12 h. After completion of the reaction a solid mass was formed. The precipitate was filtered off and washed with an ethanol–water mixture. The crude product was then recrystallized from ethanol–chloroform to obtain colourless diffraction-quality crystals of title compound.

## 6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. All H atoms were positioned geometrically and constrained to ride on their parent atom, with N–H = 0.86 Å and C–H = 0.93–0.97 Å, and with  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$  for methyl H atoms and  $1.2U_{\text{eq}}(\text{N,C})$  for all other H atoms.

## Acknowledgements

KK thanks the University grant commission for the financial support (major research project, F. No. 42-342/2013) of this research work. We are also grateful to the UGC Networking Resource Centre, University of Hyderabad, for providing characterization facilities and Dr R. Nagarajan, School of Chemistry, University of Hyderabad, for providing laboratory facilities.

## References

- Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.  
 Bruker (2002). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.  
 Hardick, D. J., Blagbrough, I. S., Cooper, G., Potter, B. V. L., Critchley, T. & Wonnacott, S. (1996). *J. Med. Chem.* **39**, 4860–4866.  
 Jeyaraman, R. & Avila, S. (1981). *Chem. Rev.* **81**, 149–174.

Table 2

Experimental details.

Crystal data	
Chemical formula	C <sub>22</sub> H <sub>23</sub> F <sub>2</sub> N <sub>3</sub> O <sub>2</sub>
$M_r$	399.43
Crystal system, space group	Monoclinic, C2/c
Temperature (K)	293
$a, b, c$ (Å)	19.751 (6), 7.087 (2), 28.492 (9)
$\beta$ (°)	102.997 (4)
$V$ (Å <sup>3</sup> )	3886 (2)
$Z$	8
Radiation type	Mo $K\alpha$
$\mu$ (mm <sup>-1</sup> )	0.10
Crystal size (mm)	0.38 × 0.36 × 0.34
Data collection	
Diffractometer	Bruker SMART CCD area detector
Absorption correction	Multi-scan ( <i>SADABS</i> ; Sheldrick, 1996)
$T_{\text{min}}, T_{\text{max}}$	0.962, 0.966
No. of measured, independent and observed [ $I > 2\sigma(I)$ ] reflections	18613, 3793, 2872
$R_{\text{int}}$	0.030
$(\sin \theta/\lambda)_{\text{max}}$ (Å <sup>-1</sup> )	0.618
Refinement	
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.050, 0.147, 1.04
No. of reflections	3793
No. of parameters	263
H-atom treatment	H-atom parameters constrained
$\Delta\rho_{\text{max}}, \Delta\rho_{\text{min}}$ (e Å <sup>-3</sup> )	0.52, -0.44

Computer programs: *SMART* and *SAINT* (Bruker, 2002), *SHELXS97* and *SHELXL97* (Sheldrick, 2008), *ORTEP-3 for Windows* (Farrugia (2012) and *PLATON* (Spek, 2009).

- Parthiban, P., Aridoss, G., Rathika, P., Ramkumar, V. & Kabilan, S. (2009). *Bioorg. Med. Chem. Lett.* **19**, 6981–6985.  
 Parthiban, P., Ramkumar, V., Santan, H. D., Kim, J. T. & Jeong, Y. T. (2008). *Acta Cryst.* **E64**, o1710.  
 Rizwana Begum, S., Hema, R., Venkateswaramoorthi, R., Krishnasamy, K. & Anitha, A. G. (2013). *Acta Cryst.* **E69**, o1525.  
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

## supporting information

*Acta Cryst.* (2014). E70, 145-147 [doi:10.1107/S1600536814018935]

## Crystal structure of methyl 2-[2,4-bis(4-fluorophenyl)-3-azabicyclo-[3.3.1]nonan-9-ylidene]hydrazinecarboxylate

A. Kamaraj, R. Rajkumar, K. Krishnasamy and S. Murugavel

### Computing details

Data collection: *SMART* (Bruker, 2002); cell refinement: *SAINTE* (Bruker, 2002); data reduction: *SAINTE* (Bruker, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia (2012)); software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008) and *PLATON* (Spek, 2009).

### 2-[2,4-Bis(4-fluorophenyl)-3-azabicyclo[3.3.1]nonan-9-ylidene]hydrazinecarboxylate

#### Crystal data

$C_{22}H_{23}F_2N_3O_2$

$M_r = 399.43$

Monoclinic, *C2/c*

Hall symbol: -C 2yc

$a = 19.751$  (6) Å

$b = 7.087$  (2) Å

$c = 28.492$  (9) Å

$\beta = 102.997$  (4)°

$V = 3886$  (2) Å<sup>3</sup>

$Z = 8$

$F(000) = 1680$

$D_x = 1.365$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 3821 reflections

$\theta = 1.5$ – $26.1$ °

$\mu = 0.10$  mm<sup>-1</sup>

$T = 293$  K

Needle, colourless

$0.38 \times 0.36 \times 0.34$  mm

#### Data collection

Bruker SMART CCD area-detector  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan  
(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.962$ ,  $T_{\max} = 0.966$

18613 measured reflections

3793 independent reflections

2872 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.030$

$\theta_{\max} = 26.1$ °,  $\theta_{\min} = 1.5$ °

$h = -24 \rightarrow 24$

$k = -8 \rightarrow 8$

$l = -35 \rightarrow 35$

#### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.050$

$wR(F^2) = 0.147$

$S = 1.04$

3793 reflections

263 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0752P)^2 + 2.7221P]$$

where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$

$$\Delta\rho_{\max} = 0.52 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.44 \text{ e } \text{\AA}^{-3}$$

### Special details

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor wR and goodness of fit S are based on  $F^2$ , conventional R-factors R are based on F, with F set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.34340 (10)	0.7153 (3)	0.09187 (6)	0.0395 (4)
H1	0.2935	0.7216	0.0775	0.047*
C2	0.37276 (10)	0.9185 (3)	0.09443 (6)	0.0397 (4)
H2	0.3614	0.9740	0.0621	0.048*
C3	0.33596 (9)	1.0266 (3)	0.12629 (6)	0.0375 (4)
C4	0.35286 (10)	0.9504 (3)	0.17646 (6)	0.0418 (4)
H4	0.3280	1.0258	0.1960	0.050*
C5	0.32457 (10)	0.7461 (3)	0.17394 (6)	0.0411 (4)
H5	0.2740	0.7514	0.1625	0.049*
C6	0.45136 (10)	0.9334 (3)	0.11470 (7)	0.0488 (5)
H6A	0.4747	0.8521	0.0960	0.059*
H6B	0.4658	1.0619	0.1105	0.059*
C7	0.47506 (11)	0.8806 (4)	0.16730 (8)	0.0568 (6)
H7A	0.4737	0.7445	0.1704	0.068*
H7B	0.5228	0.9211	0.1790	0.068*
C8	0.43043 (11)	0.9689 (3)	0.19781 (7)	0.0506 (5)
H8A	0.4421	1.1016	0.2021	0.061*
H8B	0.4411	0.9100	0.2294	0.061*
C9	0.37645 (10)	0.5907 (3)	0.06043 (7)	0.0415 (4)
C10	0.35221 (11)	0.5986 (3)	0.01104 (7)	0.0500 (5)
H10	0.3158	0.6797	-0.0020	0.060*
C11	0.38109 (13)	0.4882 (4)	-0.01910 (8)	0.0609 (6)
H11	0.3648	0.4941	-0.0523	0.073*
C12	0.43378 (14)	0.3710 (3)	0.00080 (9)	0.0622 (7)
C13	0.45997 (13)	0.3609 (3)	0.04883 (9)	0.0631 (6)
H13	0.4969	0.2808	0.0614	0.076*
C14	0.43080 (12)	0.4716 (3)	0.07882 (8)	0.0530 (5)
H14	0.4482	0.4656	0.1119	0.064*
C15	0.34057 (10)	0.6549 (3)	0.22311 (6)	0.0435 (5)
C16	0.38849 (13)	0.5154 (3)	0.23568 (8)	0.0603 (6)
H16	0.4112	0.4693	0.2128	0.072*
C17	0.40412 (15)	0.4407 (4)	0.28160 (9)	0.0718 (7)

H17	0.4368	0.3449	0.2897	0.086*
C18	0.37098 (14)	0.5097 (4)	0.31434 (8)	0.0628 (6)
C19	0.32234 (13)	0.6474 (4)	0.30379 (7)	0.0618 (6)
H19	0.3001	0.6920	0.3271	0.074*
C20	0.30660 (12)	0.7197 (3)	0.25771 (7)	0.0530 (5)
H20	0.2728	0.8129	0.2497	0.064*
C21	0.22233 (9)	1.3588 (3)	0.05966 (6)	0.0383 (4)
C22	0.14426 (13)	1.5496 (4)	0.08890 (9)	0.0650 (6)
H22A	0.1636	1.6721	0.0859	0.098*
H22B	0.1214	1.5500	0.1153	0.098*
H22C	0.1113	1.5189	0.0597	0.098*
N1	0.35236 (8)	0.6352 (2)	0.13996 (5)	0.0416 (4)
H1A	0.3728	0.5289	0.1477	0.050*
N2	0.29174 (8)	1.1591 (2)	0.11708 (5)	0.0384 (4)
N3	0.27210 (8)	1.2269 (2)	0.07053 (5)	0.0396 (4)
H3	0.2917	1.1846	0.0486	0.048*
O2	0.19861 (7)	1.4123 (2)	0.09762 (5)	0.0516 (4)
O3	0.20192 (7)	1.4228 (2)	0.01966 (5)	0.0487 (4)
F1	0.46171 (10)	0.2594 (2)	-0.02847 (6)	0.0940 (6)
F2	0.38685 (10)	0.4378 (3)	0.35949 (5)	0.0986 (6)

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0405 (10)	0.0478 (11)	0.0313 (9)	0.0021 (8)	0.0106 (7)	0.0020 (8)
C2	0.0476 (10)	0.0434 (10)	0.0304 (9)	0.0083 (8)	0.0135 (7)	0.0069 (7)
C3	0.0404 (10)	0.0407 (10)	0.0313 (9)	0.0023 (8)	0.0076 (7)	0.0024 (7)
C4	0.0489 (11)	0.0472 (11)	0.0297 (9)	0.0082 (9)	0.0099 (8)	0.0010 (8)
C5	0.0401 (10)	0.0529 (11)	0.0321 (9)	0.0018 (9)	0.0118 (7)	0.0029 (8)
C6	0.0480 (11)	0.0475 (11)	0.0548 (12)	0.0002 (9)	0.0196 (9)	0.0048 (9)
C7	0.0443 (11)	0.0638 (14)	0.0594 (13)	-0.0012 (10)	0.0058 (10)	0.0053 (11)
C8	0.0551 (12)	0.0540 (12)	0.0389 (10)	-0.0035 (10)	0.0025 (9)	0.0043 (9)
C9	0.0470 (11)	0.0421 (10)	0.0393 (10)	-0.0047 (9)	0.0176 (8)	-0.0004 (8)
C10	0.0563 (12)	0.0567 (13)	0.0398 (10)	-0.0074 (10)	0.0166 (9)	-0.0040 (9)
C11	0.0782 (16)	0.0654 (15)	0.0465 (12)	-0.0200 (13)	0.0297 (11)	-0.0146 (11)
C12	0.0825 (17)	0.0447 (12)	0.0751 (16)	-0.0148 (12)	0.0510 (14)	-0.0160 (11)
C13	0.0748 (16)	0.0466 (12)	0.0798 (17)	0.0107 (11)	0.0427 (13)	0.0097 (11)
C14	0.0636 (13)	0.0503 (12)	0.0507 (12)	0.0054 (10)	0.0247 (10)	0.0076 (9)
C15	0.0483 (11)	0.0491 (11)	0.0357 (10)	-0.0036 (9)	0.0148 (8)	0.0044 (8)
C16	0.0775 (16)	0.0636 (14)	0.0453 (12)	0.0138 (12)	0.0253 (11)	0.0119 (10)
C17	0.0903 (19)	0.0704 (17)	0.0563 (14)	0.0167 (14)	0.0196 (13)	0.0236 (12)
C18	0.0850 (17)	0.0676 (15)	0.0365 (11)	-0.0139 (13)	0.0149 (11)	0.0140 (10)
C19	0.0779 (16)	0.0761 (16)	0.0387 (11)	-0.0159 (13)	0.0286 (11)	-0.0025 (11)
C20	0.0594 (13)	0.0634 (14)	0.0411 (11)	-0.0017 (11)	0.0217 (9)	0.0018 (10)
C21	0.0400 (10)	0.0384 (10)	0.0379 (10)	0.0013 (8)	0.0114 (8)	0.0006 (8)
C22	0.0655 (15)	0.0639 (15)	0.0700 (15)	0.0241 (12)	0.0245 (12)	-0.0003 (12)
N1	0.0524 (9)	0.0429 (9)	0.0319 (8)	0.0040 (7)	0.0146 (7)	0.0046 (6)
N2	0.0431 (9)	0.0424 (9)	0.0303 (8)	0.0052 (7)	0.0093 (6)	0.0031 (6)

N3	0.0439 (8)	0.0449 (9)	0.0317 (8)	0.0089 (7)	0.0122 (6)	0.0041 (6)
O2	0.0583 (9)	0.0564 (9)	0.0440 (8)	0.0180 (7)	0.0196 (6)	0.0038 (6)
O3	0.0555 (8)	0.0514 (8)	0.0389 (7)	0.0143 (7)	0.0099 (6)	0.0087 (6)
F1	0.1259 (14)	0.0680 (10)	0.1142 (13)	-0.0134 (9)	0.0820 (11)	-0.0356 (9)
F2	0.1409 (15)	0.1101 (13)	0.0446 (8)	-0.0095 (11)	0.0204 (9)	0.0314 (8)

*Geometric parameters (Å, °)*

C1—N1	1.457 (2)	C11—H11	0.9300
C1—C9	1.507 (3)	C12—C13	1.352 (3)
C1—C2	1.548 (3)	C12—F1	1.354 (2)
C1—H1	0.9800	C13—C14	1.378 (3)
C2—C3	1.495 (2)	C13—H13	0.9300
C2—C6	1.533 (3)	C14—H14	0.9300
C2—H2	0.9800	C15—C16	1.360 (3)
C3—N2	1.269 (2)	C15—C20	1.389 (3)
C3—C4	1.494 (2)	C16—C17	1.381 (3)
C4—C8	1.522 (3)	C16—H16	0.9300
C4—C5	1.548 (3)	C17—C18	1.346 (4)
C4—H4	0.9800	C17—H17	0.9300
C5—N1	1.447 (2)	C18—F2	1.353 (2)
C5—C15	1.510 (3)	C18—C19	1.355 (4)
C5—H5	0.9800	C19—C20	1.378 (3)
C6—C7	1.513 (3)	C19—H19	0.9300
C6—H6A	0.9700	C20—H20	0.9300
C6—H6B	0.9700	C21—O3	1.208 (2)
C7—C8	1.506 (3)	C21—O2	1.327 (2)
C7—H7A	0.9700	C21—N3	1.342 (2)
C7—H7B	0.9700	C22—O2	1.428 (3)
C8—H8A	0.9700	C22—H22A	0.9600
C8—H8B	0.9700	C22—H22B	0.9600
C9—C14	1.373 (3)	C22—H22C	0.9600
C9—C10	1.382 (3)	N1—H1A	0.8600
C10—C11	1.376 (3)	N2—N3	1.3814 (19)
C10—H10	0.9300	N3—H3	0.8600
C11—C12	1.352 (4)		
N1—C1—C9	110.73 (15)	C11—C10—H10	119.5
N1—C1—C2	110.67 (15)	C9—C10—H10	119.5
C9—C1—C2	111.42 (15)	C12—C11—C10	118.4 (2)
N1—C1—H1	108.0	C12—C11—H11	120.8
C9—C1—H1	108.0	C10—C11—H11	120.8
C2—C1—H1	108.0	C13—C12—C11	122.7 (2)
C3—C2—C6	108.99 (16)	C13—C12—F1	118.4 (2)
C3—C2—C1	106.09 (14)	C11—C12—F1	118.8 (2)
C6—C2—C1	114.73 (16)	C12—C13—C14	118.6 (2)
C3—C2—H2	109.0	C12—C13—H13	120.7
C6—C2—H2	109.0	C14—C13—H13	120.7

C1—C2—H2	109.0	C9—C14—C13	120.9 (2)
N2—C3—C4	117.43 (16)	C9—C14—H14	119.6
N2—C3—C2	131.26 (16)	C13—C14—H14	119.6
C4—C3—C2	111.18 (15)	C16—C15—C20	118.16 (18)
C3—C4—C8	109.85 (16)	C16—C15—C5	122.92 (17)
C3—C4—C5	107.04 (15)	C20—C15—C5	118.90 (19)
C8—C4—C5	114.81 (16)	C15—C16—C17	121.5 (2)
C3—C4—H4	108.3	C15—C16—H16	119.2
C8—C4—H4	108.3	C17—C16—H16	119.2
C5—C4—H4	108.3	C18—C17—C16	118.5 (2)
N1—C5—C15	110.89 (16)	C18—C17—H17	120.7
N1—C5—C4	110.59 (15)	C16—C17—H17	120.7
C15—C5—C4	111.04 (15)	C17—C18—F2	118.5 (3)
N1—C5—H5	108.1	C17—C18—C19	122.6 (2)
C15—C5—H5	108.1	F2—C18—C19	118.9 (2)
C4—C5—H5	108.1	C18—C19—C20	118.4 (2)
C7—C6—C2	114.68 (16)	C18—C19—H19	120.8
C7—C6—H6A	108.6	C20—C19—H19	120.8
C2—C6—H6A	108.6	C19—C20—C15	120.8 (2)
C7—C6—H6B	108.6	C19—C20—H20	119.6
C2—C6—H6B	108.6	C15—C20—H20	119.6
H6A—C6—H6B	107.6	O3—C21—O2	123.87 (17)
C8—C7—C6	112.23 (18)	O3—C21—N3	123.44 (17)
C8—C7—H7A	109.2	O2—C21—N3	112.68 (15)
C6—C7—H7A	109.2	O2—C22—H22A	109.5
C8—C7—H7B	109.2	O2—C22—H22B	109.5
C6—C7—H7B	109.2	H22A—C22—H22B	109.5
H7A—C7—H7B	107.9	O2—C22—H22C	109.5
C7—C8—C4	113.64 (17)	H22A—C22—H22C	109.5
C7—C8—H8A	108.8	H22B—C22—H22C	109.5
C4—C8—H8A	108.8	C5—N1—C1	115.71 (15)
C7—C8—H8B	108.8	C5—N1—H1A	122.1
C4—C8—H8B	108.8	C1—N1—H1A	122.1
H8A—C8—H8B	107.7	C3—N2—N3	119.17 (15)
C14—C9—C10	118.38 (19)	C21—N3—N2	119.81 (14)
C14—C9—C1	122.62 (17)	C21—N3—H3	120.1
C10—C9—C1	118.98 (18)	N2—N3—H3	120.1
C11—C10—C9	121.0 (2)	C21—O2—C22	116.23 (16)
N1—C1—C2—C3	-56.46 (19)	C11—C12—C13—C14	1.4 (4)
C9—C1—C2—C3	179.87 (15)	F1—C12—C13—C14	-178.8 (2)
N1—C1—C2—C6	63.9 (2)	C10—C9—C14—C13	-0.8 (3)
C9—C1—C2—C6	-59.8 (2)	C1—C9—C14—C13	-179.51 (19)
C6—C2—C3—N2	124.6 (2)	C12—C13—C14—C9	-0.3 (3)
C1—C2—C3—N2	-111.4 (2)	N1—C5—C15—C16	14.8 (3)
C6—C2—C3—C4	-59.9 (2)	C4—C5—C15—C16	-108.6 (2)
C1—C2—C3—C4	64.16 (19)	N1—C5—C15—C20	-167.30 (17)
N2—C3—C4—C8	-122.34 (19)	C4—C5—C15—C20	69.3 (2)



C2—C3—C4—C8	61.4 (2)	C20—C15—C16—C17	-0.9 (4)
N2—C3—C4—C5	112.41 (19)	C5—C15—C16—C17	177.1 (2)
C2—C3—C4—C5	-63.81 (19)	C15—C16—C17—C18	-0.3 (4)
C3—C4—C5—N1	55.25 (19)	C16—C17—C18—F2	-179.3 (2)
C8—C4—C5—N1	-67.0 (2)	C16—C17—C18—C19	0.9 (4)
C3—C4—C5—C15	178.80 (15)	C17—C18—C19—C20	-0.3 (4)
C8—C4—C5—C15	56.6 (2)	F2—C18—C19—C20	180.0 (2)
C3—C2—C6—C7	52.2 (2)	C18—C19—C20—C15	-1.0 (4)
C1—C2—C6—C7	-66.5 (2)	C16—C15—C20—C19	1.5 (3)
C2—C6—C7—C8	-45.8 (3)	C5—C15—C20—C19	-176.5 (2)
C6—C7—C8—C4	46.3 (3)	C15—C5—N1—C1	-176.69 (15)
C3—C4—C8—C7	-54.2 (2)	C4—C5—N1—C1	-53.1 (2)
C5—C4—C8—C7	66.5 (2)	C9—C1—N1—C5	178.09 (15)
N1—C1—C9—C14	-26.0 (3)	C2—C1—N1—C5	54.0 (2)
C2—C1—C9—C14	97.7 (2)	C4—C3—N2—N3	-176.76 (16)
N1—C1—C9—C10	155.29 (17)	C2—C3—N2—N3	-1.5 (3)
C2—C1—C9—C10	-81.1 (2)	O3—C21—N3—N2	-179.05 (17)
C14—C9—C10—C11	0.8 (3)	O2—C21—N3—N2	1.6 (2)
C1—C9—C10—C11	179.62 (18)	C3—N2—N3—C21	175.91 (17)
C9—C10—C11—C12	0.2 (3)	O3—C21—O2—C22	2.0 (3)
C10—C11—C12—C13	-1.4 (3)	N3—C21—O2—C22	-178.61 (18)
C10—C11—C12—F1	178.83 (19)		

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N3—H3 $\cdots$ O3 <sup>i</sup>	0.86	2.12	2.930 (2)	157
C2—H2 $\cdots$ O3 <sup>i</sup>	0.98	2.49	3.442 (2)	163
C22—H22C $\cdots$ F1 <sup>ii</sup>	0.96	2.49	3.243 (3)	136

Symmetry codes: (i)  $-x+1/2, -y+5/2, -z$ ; (ii)  $-x+1/2, -y+3/2, -z$ .