



Crystal structure of 1-(4-fluorophenyl)-4-(4-methoxyphenyl)-1*H*-1,2,3-triazole

Balbir Kumar,^a Madhvi Bhardwaj,^b Satya Paul,^b Rajni Kant^a and Vivek K. Gupta^{a*}

^aPost-Graduate Department of Physics & Electronics, University of Jammu, Jammu Tawi 180 006, India, and ^bDepartment of Chemistry, University of Jammu, Jammu Tawi 180 006, India. *Correspondence e-mail: vivek_gupta2k2@hotmail.com

Received 11 June 2015; accepted 24 June 2015

Edited by A. J. Lough, University of Toronto, Canada


In the title compound, C₁₅H₁₂FN₃O, the triazole ring forms dihedral angles of 30.57 (8) and 21.81 (9)° with the fluoro-substituted and methoxy-substituted benzene rings, respectively. The dihedral angle between the benzene rings is 51.53 (7)°. In the crystal, π - π interactions between the triazole rings [centroid-centroid separations = 3.774 (2) and 3.841 (2) Å] form chains along [010].

Keywords: crystal structure; 1,2,3-triazole; π - π interactions.

CCDC reference: 1408544

1. Related literature

For related literature on 1,2,3-triazoles, see: Aher *et al.* (2009); Jordao *et al.* (2009); Vijaya Raghava Reddy *et al.* (2010); Soltis *et al.* (1996). For applications of 1,2,3-triazoles, see: Pérez-Balderas *et al.* (2003); Wu *et al.* (2004); Kumar & Pandey (2008); Haridas *et al.* (2008); Turner *et al.*, (2007); Angell & Burgess (2007); For the synthesis of 1,2,3-triazoles, see: Huisgen *et al.* (1965); Wang *et al.* (2010). For related structures, see: Abdel-Wahab *et al.* (2012); Zhang *et al.* (2004).



2. Experimental

2.1. Crystal data

C₁₅H₁₂FN₃O

$M_r = 269.28$

Triclinic, $P\bar{1}$
 $a = 5.6572$ (5) Å
 $b = 7.3692$ (8) Å
 $c = 15.5711$ (15) Å
 $\alpha = 79.202$ (9)°
 $\beta = 81.159$ (8)°
 $\gamma = 89.442$ (8)°

$V = 629.95$ (11) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹
 $T = 293$ K
 $0.30 \times 0.20 \times 0.20$ mm

2.2. Data collection

Oxford Diffraction Xcalibur
 Sapphire3 diffractometer
 Absorption correction: multi-scan
 (*CrysAlis PRO*; Oxford
 Diffraction, 2010)
 $T_{\min} = 0.806$, $T_{\max} = 1.000$

4369 measured reflections
 2461 independent reflections
 1575 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.034$

2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.057$
 $wR(F^2) = 0.179$
 $S = 1.04$
 2461 reflections

182 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.26$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.22$ e Å⁻³

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *PLATON* (Spek, 2009).

Acknowledgements

RK acknowledges the Department of Science & Technology for the single-crystal X-ray diffractometer sanctioned as a National Facility under Project No. SR/S2/CMP-47/2003.

Supporting information for this paper is available from the IUCr electronic archives (Reference: LH5772).

References

- Abdel-Wahab, B. F., Mohamed, H. A., Ng, S. W. & Tiekink, E. R. T. (2012). *Acta Cryst.* **E68**, o1956–o1957.
- Aher, N. G., Pore, V. S., Mishra, N. N., Kumar, A., Shukla, P. K., Sharma, A. & Bhat, M. K. (2009). *Bioorg. Med. Chem. Lett.* **19**, 759–763.
- Angell, Y. & Burgess, K. (2007). *Chem. Soc. Rev.* **36**, 1674–1689.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Haridas, V., Lal, K., Sharma, Y. K. & Upreti, S. (2008). *Org. Lett.* **10**, 1645–1647.
- Huisgen, R., Knorr, R., Möbius, L. & Szeimies, G. (1965). *Chem. Ber.* **98**, 4014–4021.
- Jordão, A. K., Ferreira, V. F., Lima, E. S., de Souza, M. C. B. V., Carlos, E. C. L., Castro, H. C., Geraldo, R. B., Rodrigues, C. R., Almeida, M. C. B. & Cunha, A. C. (2009). *Bioorg. Med. Chem.* **17**, 3713–3719.
- Kumar, A. & Pandey, P. S. (2008). *Org. Lett.* **10**, 165–168.
- Oxford Diffraction (2010). *CrysAlis PRO*. Oxford Diffraction Ltd, Yarnton, England.
- Pérez-Balderas, F., Ortega-Muñoz, M., Morales-Sanfrutos, J., Hernández-Mateo, F., Calvo-Flores, F. G., Calvo-Asín, J. A., Isac-García, J. & Santoyo-González, F. (2003). *Org. Lett.* **5**, 1951–1954.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Soltis, M. J., Yeh, H. J., Cole, K. A., Whittaker, N., Wersto, R. P. & Kohn, E. C. (1996). *Drug Metab. Dispos.* **24**, 799–806.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

- Turner, R. A., Oliver, A. G. & Lokey, R. S. (2007). *Org. Lett.* **9**, 5011–5014.
- Vijaya Raghava Reddy, L., Venkat Reddy, P., Mishra, N. N., Shukla, P. K., Yadav, G., Srivastava, R. & Shaw, A. K. (2010). *Carbohydr. Res.* **345**, 1515–1521.
- Wang, D., Li, N., Zhao, M., Shi, W., Ma, C. & Chen, B. (2010). *Green Chem.* **12**, 2120.
- Wu, P., Feldman, A. K., Nugent, A. K., Hawker, C. J., Scheel, A., Voit, B., Pyun, J., Fréchet, J. M. J., Sharpless, K. B. & Fokin, V. V. (2004). *Angew. Chem. Int. Ed.* **43**, 3928–3932.
- Zhang, L.-X., Zhang, A.-J., Lei, X.-X., Zou, K.-H. & Ng, S. W. (2004). *Acta Cryst.* **E60**, o613–o615.

supporting information

Acta Cryst. (2015). E71, o534–o535 [https://doi.org/10.1107/S2056989015012153]

Crystal structure of 1-(4-fluorophenyl)-4-(4-methoxyphenyl)-1*H*-1,2,3-triazole

Balbir Kumar, Madhvi Bhardwaj, Satya Paul, Rajni Kant and Vivek K. Gupta

S1. Comment

1,2,3-Triazoles are an important class of organic compounds which have become prominent in recent years as superbly versatile five membered nitrogen heterocycles. The 1,2,3-triazole family exhibit a broad spectrum of bioactivities such as antifungal (Aher *et al.*, 2009) antiviral (Jordao *et al.*, 2009), antibacterial (Vijaya Raghava Reddy *et al.*, 2010) and anticancer (Soltis *et al.*, 1996) activities. Furthermore 1,4-disubstituted 1,2,3-triazoles have also been used as a ligation tool for the synthesis of neoglyco-conjugates (Perez-Balderas *et al.*, 2003), multivalent dendrimeric peptides (Wu *et al.*, 2004), ionic receptors (Kumar *et al.*, 2008), triazolophanes (Haridas *et al.*, 2008), cyclic peptides (Turner *et al.*, 2007) and peptidomimetics (Angell *et al.*, 2007). 1,2,3-Triazoles are traditionally obtained using the thermal 1,3-dipolar cyclo-addition of organic azides with alkynes (Huisgen *et al.*, 1965) that has been known for nearly five decades. Recently, copper based catalysis was found to dramatically accelerate the reaction under mild conditions while achieving a high regioselectivity towards the 1,4-regioisomer of the triazole product (Wang *et al.*, 2010). This powerful, highly reliable, and selective reaction is the paradigm of a click reaction, which placed it in a class of its own and has enabled many novel applications.

The molecular structure of the title compound is shown in Fig. 1. The triazole ring forms dihedral angles of 30.57 (8)° and 21.81 (9)° with the fluoro-substituted and methoxy-substituted benzen rings, respectively. The dihedral angle between the benzene rings is 51.53 (7)°. All bond lengths and angles are normal and correspond to those observed in the related structures (Zhang *et al.*, 2004; Abdel-Wahab *et al.*, 2012). The C15—F1 bond length [1.357 (4) Å] agrees well with the accepted value of 1.340 Å for the F-C_{aromatic} length and is in good agreement with a structure of this type (Abdel-Wahab *et al.*, 2012). In the crystal, π - π interactions observed between the triazole rings [centroid-centroid separations = 3.774 (2) and 3.841 (2) Å] form chains along [010] (Fig. 2).

S2. Experimental

Synthesis of 1-(4-fluorophenyl)-4-(4-methoxyphenyl)-1*H*-1,2,3-triazole: To 4-fluoroaniline (0.22 g, 2 mmol) in a round bottomed flask maintained at 273-278 K, mixture of conc. HCl: H₂O (1.5 ml, 1:1) was added and stirred for 5 min. Then solution of NaNO₂ (0.17 g, 2.5 mmol in 1 ml water) was added dropwise over a period of 5 min. After stirring for another 5 min, sodium azide (0.19 g, 3 mmol) was added and the reaction mixture was further stirred for another 10 min. Finally, 4-methoxyphenylacetylene (0.19 g, 1.5 mmol) and catalyst [Cu(0)-Fe₃O₄@SiO₂/NH₂Cel] (0.05 g) were added to the reaction mixture followed by stirring at room temperature for 6 h. The reaction was then stopped and the catalyst was separated using an external magnet. The reaction mixture was extracted with EtOAc, washed with water and dried over Na₂SO₄. Finally, the product was obtained after removal of the solvent under reduced pressure followed by crystallization with EtOAc: pet ether. The product, 1-(4-fluorophenyl)-4-(4-methoxyphenyl)-1*H*-1,2,3-triazole was obtained as shiny white crystals.

S3. Refinement

All H atoms were geometrically fixed and allowed to ride on their parent C atoms, with C—H distances of 0.93–0.96 Å; and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$, except for the methyl group where $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$.




Figure 1

The molecular structure of the title compound with displacement ellipsoids drawn at the 40% probability level. H atoms are shown as small spheres of arbitrary radii.




Figure 2

The packing arrangement of molecules viewed along the *a* axis.

1-(4-Fluorophenyl)-4-(4-methoxyphenyl)-1*H*-1,2,3-triazole

Crystal data

$C_{15}H_{12}FN_3O$

$M_r = 269.28$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 5.6572$ (5) Å

$b = 7.3692$ (8) Å

$c = 15.5711$ (15) Å

$\alpha = 79.202$ (9)°

$\beta = 81.159$ (8)°

$\gamma = 89.442$ (8)°

$V = 629.95$ (11) Å³

$Z = 2$

$F(000) = 280$

$D_x = 1.420$ Mg m⁻³

$D_m = 1.42$ Mg m⁻³

D_m measured by not measured

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 1205 reflections

$\theta = 4.0$ – 28.0 °

$\mu = 0.10$ mm⁻¹

$T = 293$ K

Block, white

$0.30 \times 0.20 \times 0.20$ mm

Data collection

Oxford Diffraction Xcalibur Sapphire3
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
Detector resolution: 16.1049 pixels mm⁻¹
 ω scans
Absorption correction: multi-scan
(*CrysAlis PRO*; Oxford Diffraction, 2010)
 $T_{\min} = 0.806$, $T_{\max} = 1.000$

4369 measured reflections
2461 independent reflections
1575 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.034$
 $\theta_{\max} = 26.0^\circ$, $\theta_{\min} = 3.7^\circ$
 $h = -4 \rightarrow 6$
 $k = -7 \rightarrow 9$
 $l = -18 \rightarrow 19$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.057$
 $wR(F^2) = 0.179$
 $S = 1.04$
2461 reflections
182 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0744P)^2 + 0.0652P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.26 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.22 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
F1	0.2052 (4)	-0.1487 (2)	0.41281 (12)	0.0782 (6)
N1	0.1232 (4)	0.1852 (3)	0.07275 (13)	0.0404 (5)
N2	-0.0970 (4)	0.1982 (3)	0.04842 (15)	0.0505 (6)
N3	-0.0714 (4)	0.2743 (3)	-0.03467 (15)	0.0489 (6)
C4	0.1657 (4)	0.3132 (3)	-0.06593 (16)	0.0367 (6)
C5	0.2893 (4)	0.2552 (3)	0.00235 (15)	0.0395 (6)
H5	0.4540	0.2623	0.0009	0.047*
C6	0.2436 (4)	0.3971 (3)	-0.15737 (16)	0.0371 (6)
C7	0.1015 (5)	0.3849 (3)	-0.22101 (17)	0.0426 (6)
H7	-0.0454	0.3228	-0.2040	0.051*
C8	0.1707 (5)	0.4613 (4)	-0.30769 (18)	0.0483 (7)
H8	0.0717	0.4501	-0.3490	0.058*
C9	0.3880 (5)	0.5558 (3)	-0.33479 (17)	0.0457 (7)
O9	0.4360 (4)	0.6328 (3)	-0.42195 (14)	0.0718 (7)
C10	0.5341 (5)	0.5698 (3)	-0.27319 (17)	0.0452 (6)
H10	0.6804	0.6327	-0.2906	0.054*

C11	0.4626 (5)	0.4900 (3)	-0.18530 (16)	0.0414 (6)
H11	0.5629	0.4987	-0.1441	0.050*
C12	0.1484 (4)	0.1035 (3)	0.16056 (16)	0.0371 (6)
C13	-0.0340 (5)	0.1177 (3)	0.22842 (16)	0.0430 (6)
H13	-0.1694	0.1844	0.2166	0.052*
C14	-0.0162 (5)	0.0338 (4)	0.31322 (18)	0.0518 (7)
H14	-0.1391	0.0417	0.3594	0.062*
C15	0.1878 (5)	-0.0630 (3)	0.32897 (18)	0.0500 (7)
C16	0.3719 (5)	-0.0751 (3)	0.26282 (19)	0.0503 (7)
H16	0.5088	-0.1392	0.2752	0.060*
C17	0.3531 (5)	0.0081 (3)	0.17797 (17)	0.0425 (6)
H17	0.4773	0.0007	0.1322	0.051*
C18	0.6617 (8)	0.6997 (5)	-0.4596 (2)	0.0918 (12)
H18A	0.7031	0.7988	-0.4321	0.138*
H18B	0.6648	0.7446	-0.5218	0.138*
H18C	0.7749	0.6025	-0.4512	0.138*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
F1	0.0830 (15)	0.0859 (12)	0.0607 (12)	0.0042 (11)	-0.0198 (10)	0.0062 (9)
N1	0.0268 (11)	0.0459 (12)	0.0481 (13)	0.0005 (9)	-0.0039 (9)	-0.0096 (9)
N2	0.0252 (11)	0.0691 (14)	0.0558 (15)	-0.0006 (10)	-0.0048 (10)	-0.0098 (11)
N3	0.0260 (12)	0.0675 (15)	0.0508 (14)	-0.0006 (11)	-0.0033 (10)	-0.0069 (11)
C4	0.0268 (13)	0.0389 (12)	0.0455 (15)	0.0003 (10)	-0.0056 (10)	-0.0105 (10)
C5	0.0224 (12)	0.0470 (14)	0.0483 (15)	-0.0018 (11)	-0.0025 (10)	-0.0090 (11)
C6	0.0279 (13)	0.0391 (12)	0.0458 (15)	0.0054 (10)	-0.0075 (11)	-0.0108 (10)
C7	0.0303 (13)	0.0450 (13)	0.0536 (16)	-0.0001 (11)	-0.0086 (11)	-0.0107 (11)
C8	0.0393 (15)	0.0583 (16)	0.0513 (17)	0.0049 (13)	-0.0158 (13)	-0.0136 (12)
C9	0.0450 (16)	0.0499 (14)	0.0404 (15)	0.0101 (13)	-0.0063 (12)	-0.0049 (11)
O9	0.0608 (14)	0.0911 (15)	0.0542 (13)	-0.0043 (12)	-0.0021 (11)	0.0045 (11)
C10	0.0321 (14)	0.0458 (14)	0.0553 (17)	-0.0029 (12)	0.0001 (12)	-0.0083 (12)
C11	0.0331 (14)	0.0447 (13)	0.0484 (15)	0.0003 (11)	-0.0088 (11)	-0.0120 (11)
C12	0.0293 (13)	0.0358 (12)	0.0458 (15)	-0.0034 (10)	-0.0048 (11)	-0.0070 (10)
C13	0.0316 (13)	0.0452 (14)	0.0497 (16)	0.0032 (11)	-0.0020 (11)	-0.0063 (11)
C14	0.0441 (16)	0.0556 (16)	0.0522 (18)	0.0000 (14)	0.0027 (13)	-0.0092 (12)
C15	0.0532 (18)	0.0471 (15)	0.0493 (17)	-0.0049 (13)	-0.0165 (14)	-0.0011 (12)
C16	0.0401 (15)	0.0457 (14)	0.0657 (19)	0.0059 (12)	-0.0131 (14)	-0.0079 (13)
C17	0.0317 (13)	0.0425 (13)	0.0535 (16)	0.0001 (11)	-0.0038 (11)	-0.0121 (11)
C18	0.091 (3)	0.109 (3)	0.068 (2)	-0.032 (2)	0.000 (2)	-0.0060 (19)

Geometric parameters (Å, °)

F1—C15	1.357 (3)	O9—C18	1.376 (4)
N1—C5	1.354 (3)	C10—C11	1.384 (3)
N1—N2	1.354 (3)	C10—H10	0.9300
N1—C12	1.415 (3)	C11—H11	0.9300
N2—N3	1.297 (3)	C12—C13	1.377 (3)

N3—C4	1.368 (3)	C12—C17	1.385 (3)
C4—C5	1.362 (3)	C13—C14	1.367 (3)
C4—C6	1.444 (3)	C13—H13	0.9300
C5—H5	0.9300	C14—C15	1.381 (4)
C6—C7	1.384 (3)	C14—H14	0.9300
C6—C11	1.390 (3)	C15—C16	1.363 (4)
C7—C8	1.360 (3)	C16—C17	1.368 (3)
C7—H7	0.9300	C16—H16	0.9300
C8—C9	1.387 (4)	C17—H17	0.9300
C8—H8	0.9300	C18—H18A	0.9600
C9—O9	1.356 (3)	C18—H18B	0.9600
C9—C10	1.377 (4)	C18—H18C	0.9600
C5—N1—N2	109.5 (2)	C10—C11—C6	121.3 (2)
C5—N1—C12	130.8 (2)	C10—C11—H11	119.4
N2—N1—C12	119.7 (2)	C6—C11—H11	119.4
N3—N2—N1	107.7 (2)	C13—C12—C17	120.3 (2)
N2—N3—C4	109.6 (2)	C13—C12—N1	119.3 (2)
C5—C4—N3	107.4 (2)	C17—C12—N1	120.4 (2)
C5—C4—C6	131.8 (2)	C14—C13—C12	120.1 (2)
N3—C4—C6	120.8 (2)	C14—C13—H13	120.0
N1—C5—C4	105.8 (2)	C12—C13—H13	120.0
N1—C5—H5	127.1	C13—C14—C15	118.7 (2)
C4—C5—H5	127.1	C13—C14—H14	120.7
C7—C6—C11	117.5 (2)	C15—C14—H14	120.7
C7—C6—C4	120.5 (2)	F1—C15—C16	119.1 (2)
C11—C6—C4	122.0 (2)	F1—C15—C14	119.0 (3)
C8—C7—C6	121.9 (3)	C16—C15—C14	121.9 (3)
C8—C7—H7	119.1	C15—C16—C17	119.2 (2)
C6—C7—H7	119.1	C15—C16—H16	120.4
C7—C8—C9	120.3 (2)	C17—C16—H16	120.4
C7—C8—H8	119.9	C16—C17—C12	119.7 (2)
C9—C8—H8	119.9	C16—C17—H17	120.1
O9—C9—C10	125.0 (3)	C12—C17—H17	120.1
O9—C9—C8	115.6 (3)	O9—C18—H18A	109.5
C10—C9—C8	119.3 (2)	O9—C18—H18B	109.5
C9—O9—C18	120.7 (3)	H18A—C18—H18B	109.5
C9—C10—C11	119.8 (3)	O9—C18—H18C	109.5
C9—C10—H10	120.1	H18A—C18—H18C	109.5
C11—C10—H10	120.1	H18B—C18—H18C	109.5
C5—N1—N2—N3	-0.1 (3)	O9—C9—C10—C11	178.0 (2)
C12—N1—N2—N3	-178.86 (19)	C8—C9—C10—C11	-0.1 (4)
N1—N2—N3—C4	-0.3 (3)	C9—C10—C11—C6	-0.7 (4)
N2—N3—C4—C5	0.6 (3)	C7—C6—C11—C10	1.0 (3)
N2—N3—C4—C6	179.3 (2)	C4—C6—C11—C10	-179.9 (2)
N2—N1—C5—C4	0.5 (2)	C5—N1—C12—C13	150.9 (2)
C12—N1—C5—C4	179.1 (2)	N2—N1—C12—C13	-30.7 (3)

N3—C4—C5—N1	-0.7 (2)	C5—N1—C12—C17	-30.0 (3)
C6—C4—C5—N1	-179.1 (2)	N2—N1—C12—C17	148.5 (2)
C5—C4—C6—C7	156.9 (2)	C17—C12—C13—C14	-1.6 (4)
N3—C4—C6—C7	-21.4 (3)	N1—C12—C13—C14	177.6 (2)
C5—C4—C6—C11	-22.2 (4)	C12—C13—C14—C15	0.6 (4)
N3—C4—C6—C11	159.5 (2)	C13—C14—C15—F1	-179.1 (2)
C11—C6—C7—C8	-0.4 (3)	C13—C14—C15—C16	0.8 (4)
C4—C6—C7—C8	-179.5 (2)	F1—C15—C16—C17	178.8 (2)
C6—C7—C8—C9	-0.4 (4)	C14—C15—C16—C17	-1.2 (4)
C7—C8—C9—O9	-177.6 (2)	C15—C16—C17—C12	0.1 (4)
C7—C8—C9—C10	0.7 (4)	C13—C12—C17—C16	1.2 (4)
C10—C9—O9—C18	14.0 (4)	N1—C12—C17—C16	-177.9 (2)
C8—C9—O9—C18	-167.9 (3)		
