



Crystal structure of dichloridobis[2-(phenyldiazenyl)pyridine- κ N¹]zinc

Luksamee Vittaya,^{a*} Nararak Leesakul,^b Saowanit Saithong^b and Kittipong Chainok^c

^aFaculty of Science and Fisheries Technology, Rajamangala University of Technology Srivijaya, Sikao, Trang 92150, Thailand, ^bDepartment of Chemistry and Center for Innovation in Chemistry, Faculty of Science, Prince of Songkla University, Hat Yai, Songkhla 90112, Thailand, and ^cDepartment of Physics, Faculty of Science and Technology, Thammasat University, Khlong Luang, Pathum Thani 12121, Thailand.
*Correspondence e-mail: nokluksamee@hotmail.com

Received 1 October 2015; accepted 10 October 2015

Edited by M. Zeller, Youngstown State University, USA


The structure of the title complex, $[\text{ZnCl}_2(\text{C}_{11}\text{H}_9\text{N}_3)_2]$, comprises two 2-(phenyldiazenyl)pyridine ligands coordinating to a central Zn^{II} dichloride unit *via* the pyridyl N-atom donors, resulting in a slightly distorted tetrahedral geometry. The complex exhibits twofold rotation symmetry, with the rotation axis bisecting the zinc cation. The structure is stabilized by weak intermolecular $\text{C}\cdots\text{H}\cdots\text{Cl}$ interactions [$\text{C}\cdots\text{Cl} = 3.411(2)$ and $3.675(2)$ Å], connecting neighbouring molecules into layers perpendicular to the *c* axis.

Keywords: crystal structure; zinc complex; $\text{C}\cdots\text{H}\cdots\text{Cl}$ interactions.

CCDC reference: 1430587

1. Related literature

For background to diazenylpyridine compounds, see: Krause & Krause (1980). For applications of diazenylpyridine complexes, see: Wong & Giandomenico (1999); Wu *et al.* (2006); Hotze *et al.* (2004); Velders *et al.* (2000); Barf & Sheldon (1995). For applications of zinc–diazenyl complexes, see: Saha *et al.* (2014); Dutta *et al.* (2014); Datta *et al.* (2014); Zhang *et al.* (2012). For related structures, see: Leesakul *et al.* (2011); Panneerselvam *et al.* (2000); Steffen & Palenik (1976).



2. Experimental

2.1. Crystal data

$[\text{ZnCl}_2(\text{C}_{11}\text{H}_9\text{N}_3)_2]$
 $M_r = 502.69$
Orthorhombic, *Pbcn*
 $a = 13.7960(4)$ Å
 $b = 10.1905(3)$ Å
 $c = 16.1305(5)$ Å

$V = 2267.76(12)$ Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 1.34$ mm⁻¹
 $T = 298$ K
 $0.36 \times 0.32 \times 0.30$ mm

2.2. Data collection

Bruker APEXII CCD
diffractometer
Absorption correction: multi-scan
(*SADABS*; Bruker, 2013)
 $T_{\text{min}} = 0.708$, $T_{\text{max}} = 0.746$

65168 measured reflections
2820 independent reflections
2160 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.041$

2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.083$
 $S = 1.06$
2820 reflections

141 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.35$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.22$ e Å⁻³

Table 1
Hydrogen-bond geometry (Å, °).

<i>D</i> — <i>H</i> ⋯ <i>A</i>	<i>D</i> — <i>H</i>	<i>H</i> ⋯ <i>A</i>	<i>D</i> ⋯ <i>A</i>	<i>D</i> — <i>H</i> ⋯ <i>A</i>
$\text{C3}—\text{H3}\cdots\text{Cl1}^{\dagger}$	0.93	2.75	3.675 (2)	173
$\text{C1}—\text{H1}\cdots\text{Cl1}$	0.93	2.81	3.411 (2)	124

Symmetry code: (i) $x - \frac{1}{2}, y + \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: *SMART* (Bruker, 2013); cell refinement: *SAINT* (Bruker, 2013); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *WinGX* (Farrugia, 2012) and *pubCIF* (Westrip, 2010).

Acknowledgements

We are grateful to the Faculty of Science and Fisheries Technology, Rajamangala University of Technology Srivijaya, for financial support and to the Center for Innovation in Chemistry (PERCH–CIC) Commission on Higher Education,

Ministry of Education, Thailand, for partial support. We are also grateful for support with facilities from the Department of Chemistry, Faculty of Science, Prince of Songkla University. We express our acknowledgements to Dr Brian Hodgson, Faculty of Pharmaceutical Science, Prince of Songkla University, for reading the manuscript and providing comments.

Supporting information for this paper is available from the IUCr electronic archives (Reference: ZL2646).

References

- Barf, G. A. & Sheldon, R. A. (1995). *J. Mol. Catal. A Chem.* **98**, 143–146.
- Bruker (2013). *SMART, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Datta, P., Mallick, D., Mondal, T.-K. & Sinha, C. (2014). *Polyhedron*, **71**, 47–61.
- Dutta, P., Mallick, D., Roy, S., Torres, E.-L. & Sinha, C. (2014). *Inorg. Chim. Acta*, **423**, 397–407.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Hotze, A. C. G., Caspers, S. E., de Vos, D., Kooijman, H., Spek, A. L., Flamigni, A., Bacac, M., Sava, G., Haasnoot, J. G. & Reedijk, J. (2004). *J. Biol. Inorg. Chem.* **9**, 354–364.
- Krause, R. A. & Krause, K. (1980). *Inorg. Chem.* **19**, 2600–2603.
- Leesakul, N., Pakawatchai, C., Saithong, S., Tantirungrotechai, Y. & Kwanplod, K. (2011). *Acta Cryst.* **E67**, m955–m956.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Panneerselvam, K., Hansongnern, K., Rattanawit, N., Liao, F.-L. & Lu, T.-H. (2000). *Anal. Sci.* **16**, 1107–1108.
- Saha (Halder), S., Mitra, P. & Sinha, C. (2014). *Polyhedron*, **67**, 321–328.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
- Steffen, W. L. & Palenik, G. J. (1976). *Acta Cryst.* **B32**, 298–300.
- Velders, A. H., Kooijman, H., Spek, A. L., Haasnoot, J. G., de Vos, D. & Reedijk, J. (2000). *Inorg. Chem.* **39**, 2966–2967.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Wong, E. & Giandomenico, C. M. (1999). *Chem. Rev.* **99**, 2451–2466.
- Wu, B.-Z., Chang, C.-C., Sree, U., Chiu, K.-H. & Lo, J.-G. (2006). *Anal. Chim. Acta*, **576**, 91–99.
- Zhang, G., Wang, S., Ma, J.-S. & Yang, G. (2012). *Inorg. Chim. Acta*, **384**, 97–104.

supporting information

Acta Cryst. (2015). E71, m201–m202 [https://doi.org/10.1107/S2056989015019143]

Crystal structure of dichloridobis[2-(phenyldiazenyl)pyridine- κN^1]zinc

Luksamee Vittaya, Nararak Leesakul, Saowanit Saithong and Kittipong Chainok

S1. Comment

Nitrogen containing heteroaromatic systems like pyridines are some of the most investigated organic compounds primarily because of their importance in the pharmaceutical and chemical industries. Pyridine derivatives have been widely studied in analytical chemistry as an acid-base, redox and biomedical agent and for their photo-physical properties. A common synthetic pyridine derivative is 2-(phenyldiazenyl)pyridine (Krause & Krause, 1980). This compound is a well known ligand with a basic nitrogen atom on the pyridine ring able to coordinate to transition metal complexes. It has various applications, for example as a chemotherapeutic drug (Wong & Giandomenico, 1999), it has anticancer properties (Wu *et al.*, 2006; Hotze *et al.*, 2004; Velder *et al.*, 2000) and is active in catalysing the epoxidation of olefins (Barf & Sheldon 1995). Coordination compounds of Zn^{II} incorporating the diazenyl moiety were shown to have photochromic properties (Saha *et al.*, 2014; Dutta *et al.*, 2014; Datta *et al.*, 2014) as well as non-linear optical properties, which were investigated for the storage of optical information (Zhang *et al.*, 2012).

We here report the preparation and crystal structure of a new Zn^{II} complex with the well known diazenylpyridine ligand 2-(phenyldiazenyl)pyridine (C₁₁H₉N₃), or azpy. The molecular structure of the title complex, [Zn(C₁₁H₉N₃)₂Cl₂], is slightly distorted tetrahedral as illustrated in Fig.1. The Zn atom is four coordinated by two azpy ligands *via* two N(pyridine) atoms [Zn1—N1 and Zn1—N1ⁱ = 2.0618 (15) Å] and two Cl⁻ ions [Zn1—Cl1 and Zn1—Cl1ⁱ = 2.2472 (5) Å ; (i): -x+2,y,-z+3/2]. These values compare well with those of other related dichloro Zn^{II} compounds with pyridine ligands, such as [Zn(N,N-diethyl-4-[(pyridin-2-yl-*k*N)diazenyl]aniline)₂Cl₂] (Leesakul *et al.*, 2011) with Zn—N distances of 2.0513 (9) and 2.0439 (19) Å or [Zn(pyridine)₂Cl₂] (Steffen & Palenik, 1976) with Zn—N distances of 2.046 (5) and 2.052 (6) Å. The reported Zn—Cl bond distances in the complex of Leesakul *et al.* averaged to 2.264 (6) Å, those in [Zn(pyridine)₂Cl₂] (Steffen & Palenik, 1976) averaged to 2.2215 Å, which are thus slightly longer and shorter respectively than those of the title complex [Zn(azpy)₂Cl₂].

The azo (N=N) distance in the Zn^{II} complex is 1.244 (2) Å, which is comparable to that in the free azpy ligand, 1.248 (4) Å (Panneerselvam *et al.*, 2000), which is expected as the azo nitrogen of the azpy ligand is not metal coordinated. All N—Zn—Cl, Cl—Zn—Cl and N—Zn—N bond angles deviate somewhat from the ideal from 109.5°, especially for the angle N1—Zn1—N1ⁱ = 124.45 (9)°, probably due to steric demands of the azpy ligand. The torsion angle of the pyridine-azo-phenyl atoms, C5—N2—N3—C6 is 179.59 (15)°. The dihedral angle of the mean planes between the pyridine and the phenyl ring in the ligand molecule is 11.9 (1)°.

S2. Supramolecular features

Weak intra and intermolecular C—H...Cl interactions (C...Cl = 3.411 (2) and 3.675 (2) Å; see Table 1, Hydrogen-bond geometry) connect neighboring molecules into layers perpendicular to the *c*-axis (Fig. 2–4).

S3. Experimental

An acetonitrile solution (10 mL) of 2-(phenyldiazenyl)pyridine (azpy) (0.183 g, 1.0 mmol) was added dropwise to zinc(II) chloride (0.068 g, 0.50 mmol), then refluxed for 4 h. After being filtered, the filtrate was left standing overnight at 4°C. Orange crystals were obtained (yield 71.31%, 0.179 g). Anal. Calcd for $\text{ZnC}_{22}\text{H}_{18}\text{N}_6\text{Cl}_2$: C, 52.56; H, 3.61; N, 16.72. Found: C, 52.56; H, 3.55; N, 16.96.

S4. Refinement

All H atoms of aromatic carbon were positioned geometrically and refined as riding atoms with $\text{C—H} = 0.93 \text{ \AA}$, and with $U_{\text{eq}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$.




Figure 1

Molecular structure of $[\text{Zn}(\text{C}_{11}\text{H}_9\text{N}_3)_2\text{Cl}_2]$ with thermal ellipsoids plotted at the 30% probability level. Non-labelled atoms are created by the twofold symmetry axis [symmetry operator: (i) $-x + 2, y, -z + 3/2$].




Figure 2

Two-dimensional interaction sheet of $[\text{Zn}(\text{C}_{11}\text{H}_9\text{N}_3)_2\text{Cl}_2]$ plotted down c , formed through weak $\text{C-H}\cdots\text{Cl}$ interactions.




Figure 3

Two-dimensional interaction sheet of $[\text{Zn}(\text{C}_{11}\text{H}_9\text{N}_3)_2\text{Cl}_2]$ plotted down b axis, formed through weak $\text{C}-\text{H}\cdots\text{Cl}$ interactions.




Figure 4

The arrangement of two-dimensional layers plotted down the *a* axis showing a lateral view of alternating C...Cl contact directions of adjacent sheets (A and B layers). H atoms are omitted for clarity.

Dichloridobis[2-(phenyldiazenyl)pyridine- κN^1]zinc

Crystal data

[ZnCl₂(C₁₁H₉N₃)₂]

M_r = 502.69

Orthorhombic, *Pbcn*

a = 13.7960 (4) Å

b = 10.1905 (3) Å

c = 16.1305 (5) Å

V = 2267.76 (12) Å³

Z = 4

F(000) = 1024

Data collection

Bruker APEXII CCD

diffractometer

φ and ω scans

D_x = 1.472 Mg m⁻³

Mo *K* α radiation, λ = 0.71073 Å

Cell parameters from 9900 reflections

θ = 3.2–28.2°

μ = 1.34 mm⁻¹

T = 298 K

Block, orange

0.36 × 0.32 × 0.30 mm

Absorption correction: multi-scan

(*SADABS*; Bruker, 2013)

T_{min} = 0.708, *T_{max}* = 0.746

65168 measured reflections

2820 independent reflections
 2160 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.041$
 $\theta_{\text{max}} = 28.3^\circ$, $\theta_{\text{min}} = 3.5^\circ$

$h = -18 \rightarrow 18$
 $k = -13 \rightarrow 13$
 $l = -21 \rightarrow 21$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.083$
 $S = 1.06$
 2820 reflections
 141 parameters
 0 restraints

Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0384P)^2 + 0.7794P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.35 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.22 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	1.0000	0.21091 (3)	0.7500	0.04573 (11)
C11	0.96785 (4)	0.08709 (5)	0.63805 (3)	0.06114 (15)
N1	0.86977 (11)	0.30520 (15)	0.76959 (10)	0.0468 (3)
N2	0.95560 (12)	0.45206 (15)	0.84789 (10)	0.0514 (4)
N3	0.95301 (13)	0.55766 (16)	0.88646 (11)	0.0584 (4)
C1	0.78854 (15)	0.2566 (2)	0.73676 (13)	0.0571 (5)
H1	0.7927	0.1832	0.7026	0.069*
C2	0.69884 (17)	0.3110 (2)	0.75155 (14)	0.0676 (6)
H2	0.6434	0.2755	0.7275	0.081*
C3	0.69288 (16)	0.4185 (2)	0.80247 (16)	0.0730 (6)
H3	0.6330	0.4563	0.8140	0.088*
C4	0.77541 (15)	0.4698 (2)	0.83614 (14)	0.0627 (5)
H4	0.7726	0.5429	0.8706	0.075*
C5	0.86318 (13)	0.41115 (18)	0.81815 (11)	0.0484 (4)
C6	1.04406 (15)	0.60029 (19)	0.91730 (12)	0.0555 (5)
C7	1.04401 (19)	0.7222 (2)	0.95484 (16)	0.0696 (6)
H7	0.9866	0.7696	0.9593	0.084*
C8	1.1291 (2)	0.7733 (3)	0.98561 (16)	0.0789 (7)
H8	1.1293	0.8553	1.0110	0.095*
C9	1.2132 (2)	0.7037 (3)	0.97888 (15)	0.0763 (7)
H9	1.2707	0.7390	0.9990	0.092*
C10	1.21338 (17)	0.5810 (2)	0.94239 (14)	0.0706 (6)
H10	1.2709	0.5338	0.9386	0.085*
C11	1.12915 (16)	0.5283 (2)	0.91168 (13)	0.0623 (5)
H11	1.1291	0.4455	0.8874	0.075*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.03382 (16)	0.04842 (18)	0.05494 (19)	0.000	0.00041 (11)	0.000
Cl1	0.0475 (3)	0.0725 (3)	0.0635 (3)	0.0004 (2)	-0.0018 (2)	-0.0153 (2)
N1	0.0390 (8)	0.0491 (8)	0.0522 (8)	0.0044 (6)	0.0020 (6)	0.0048 (6)
N2	0.0533 (9)	0.0493 (8)	0.0518 (9)	0.0059 (7)	0.0038 (7)	0.0004 (7)
N3	0.0597 (11)	0.0517 (9)	0.0638 (10)	0.0067 (8)	0.0061 (8)	-0.0026 (8)
C1	0.0431 (10)	0.0613 (11)	0.0669 (13)	0.0021 (9)	-0.0025 (8)	0.0013 (9)
C2	0.0393 (10)	0.0797 (15)	0.0837 (16)	0.0077 (10)	-0.0056 (10)	0.0097 (12)
C3	0.0490 (12)	0.0840 (16)	0.0861 (16)	0.0254 (11)	0.0070 (11)	0.0133 (13)
C4	0.0602 (12)	0.0615 (12)	0.0663 (12)	0.0211 (10)	0.0057 (10)	0.0027 (10)
C5	0.0472 (10)	0.0497 (10)	0.0483 (10)	0.0089 (8)	0.0034 (8)	0.0093 (8)
C6	0.0584 (12)	0.0551 (11)	0.0530 (11)	0.0001 (9)	0.0074 (9)	0.0013 (9)
C7	0.0696 (14)	0.0602 (13)	0.0790 (15)	0.0002 (10)	0.0151 (12)	-0.0122 (11)
C8	0.0851 (18)	0.0683 (15)	0.0832 (16)	-0.0170 (13)	0.0133 (14)	-0.0153 (12)
C9	0.0742 (16)	0.0856 (17)	0.0690 (14)	-0.0211 (13)	0.0029 (12)	0.0037 (12)
C10	0.0624 (13)	0.0789 (16)	0.0704 (14)	0.0030 (11)	0.0025 (11)	0.0065 (12)
C11	0.0657 (13)	0.0609 (12)	0.0602 (12)	0.0051 (10)	0.0035 (10)	-0.0020 (10)

Geometric parameters (\AA , $^\circ$)

Zn1—N1	2.0618 (15)	C3—H3	0.9300
Zn1—N1 ⁱ	2.0619 (15)	C4—C5	1.381 (3)
Zn1—Cl1 ⁱ	2.2471 (5)	C4—H4	0.9300
Zn1—Cl1	2.2472 (5)	C6—C7	1.382 (3)
N1—C1	1.335 (3)	C6—C11	1.387 (3)
N1—C5	1.337 (2)	C7—C8	1.377 (3)
N2—N3	1.244 (2)	C7—H7	0.9300
N2—C5	1.425 (2)	C8—C9	1.364 (4)
N3—C6	1.419 (3)	C8—H8	0.9300
C1—C2	1.377 (3)	C9—C10	1.382 (3)
C1—H1	0.9300	C9—H9	0.9300
C2—C3	1.372 (3)	C10—C11	1.372 (3)
C2—H2	0.9300	C10—H10	0.9300
C3—C4	1.365 (3)	C11—H11	0.9300
N1—Zn1—N1 ⁱ	124.45 (9)	C5—C4—H4	120.6
N1—Zn1—Cl1 ⁱ	108.09 (4)	N1—C5—C4	122.15 (19)
N1 ⁱ —Zn1—Cl1 ⁱ	102.29 (5)	N1—C5—N2	111.89 (15)
N1—Zn1—Cl1	102.29 (5)	C4—C5—N2	125.96 (18)
N1 ⁱ —Zn1—Cl1	108.09 (4)	C7—C6—C11	120.3 (2)
Cl1 ⁱ —Zn1—Cl1	111.68 (3)	C7—C6—N3	115.36 (19)
C1—N1—C5	118.37 (17)	C11—C6—N3	124.36 (19)
C1—N1—Zn1	119.85 (13)	C8—C7—C6	119.8 (2)
C5—N1—Zn1	121.69 (13)	C8—C7—H7	120.1
N3—N2—C5	113.33 (16)	C6—C7—H7	120.1
N2—N3—C6	114.51 (17)	C9—C8—C7	120.0 (2)

N1—C1—C2	122.4 (2)	C9—C8—H8	120.0
N1—C1—H1	118.8	C7—C8—H8	120.0
C2—C1—H1	118.8	C8—C9—C10	120.4 (2)
C3—C2—C1	118.6 (2)	C8—C9—H9	119.8
C3—C2—H2	120.7	C10—C9—H9	119.8
C1—C2—H2	120.7	C11—C10—C9	120.4 (2)
C4—C3—C2	119.6 (2)	C11—C10—H10	119.8
C4—C3—H3	120.2	C9—C10—H10	119.8
C2—C3—H3	120.2	C10—C11—C6	119.1 (2)
C3—C4—C5	118.8 (2)	C10—C11—H11	120.5
C3—C4—H4	120.6	C6—C11—H11	120.5
C5—N2—N3—C6	179.59 (15)	N3—N2—C5—N1	173.32 (16)
C5—N1—C1—C2	-0.4 (3)	N3—N2—C5—C4	-7.2 (3)
Zn1—N1—C1—C2	176.25 (16)	N2—N3—C6—C7	175.28 (19)
N1—C1—C2—C3	-0.5 (3)	N2—N3—C6—C11	-4.7 (3)
C1—C2—C3—C4	0.9 (3)	C11—C6—C7—C8	1.0 (3)
C2—C3—C4—C5	-0.3 (3)	N3—C6—C7—C8	-179.0 (2)
C1—N1—C5—C4	1.0 (3)	C6—C7—C8—C9	0.1 (4)
Zn1—N1—C5—C4	-175.57 (14)	C7—C8—C9—C10	-1.0 (4)
C1—N1—C5—N2	-179.50 (16)	C8—C9—C10—C11	0.7 (4)
Zn1—N1—C5—N2	3.9 (2)	C9—C10—C11—C6	0.4 (3)
C3—C4—C5—N1	-0.7 (3)	C7—C6—C11—C10	-1.3 (3)
C3—C4—C5—N2	179.92 (19)	N3—C6—C11—C10	178.7 (2)

Symmetry code: (i) $-x+2, y, -z+3/2$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C3—H3 \cdots C11 ⁱⁱ	0.93	2.75	3.675 (2)	173
C1—H1 \cdots C11	0.93	2.81	3.411 (2)	124

Symmetry code: (ii) $x-1/2, y+1/2, -z+3/2$.