

## Crystal structure of bis(quinolin-1-ium) tetrachloridoferrate(III) chloride

Azzedine Boudjarda,<sup>a</sup> Karim Bouchouit,<sup>a\*</sup> Samiha Arroudj,<sup>b</sup> Sofiane Bouacida<sup>c,d</sup> and Hocine Merzig<sup>c</sup>

<sup>a</sup>Faculté des Sciences Exactes et Informatique, Département de Chimie, Université de Jijel, 18000 Jijel, Algeria, <sup>b</sup>Laboratoire des Structures, Propriétés et Interactions InterAtomiques, Université de Khenchela, 40000 Khenchela, Algeria, <sup>c</sup>Unité de Recherche de Chimie de l'Environnement et Moléculaire Structurale, CHEMS, Université Constantine 1, 25000, Algeria, and <sup>d</sup>Département Sciences de la Matière, Université Oum El Bouaghi, Algeria. \*Correspondence e-mail: karim.bouchouit@laposte.net

Received 19 November 2015; accepted 21 December 2015

Edited by A. M. Chippindale, University of Reading, England

The asymmetric unit of the title hybrid compound,  $(C_9H_8N)[FeCl_4]Cl$ , comprises a tetrahedral tetrachloridoferrate(III) anion,  $[FeCl_4]^-$ , a  $Cl^-$  anion and two quinolinium cations. There are  $N-H \cdots Cl$  hydrogen-bonding interactions between the protonated N atoms of the quinolinium cations and the chloride anion, which together with  $\pi-\pi$  stacking between adjacent quinolinium rings [centroid-to-centroid distances between  $C_6$  and  $C_5N$  rings in adjacent stacked quinolinium cations of 3.609 (2) and 3.802 (2) Å] serve to hold the structure together.

**Keywords:** crystal structure; hybrid compounds; tetrachloridoferrate(III) anion;  $N-H \cdots Cl$  hydrogen bonding.

**CCDC reference:** 1443665

### 1. Related literature

For non-linear optical properties of hybrid compounds, see: Bouchouit *et al.* (2008, 2010, 2015); Jayalakshmi & Kumar (2006); Sankar *et al.* (2007). For similar structures containing the  $[FeCl_4]^-$  anion, see: Khadri *et al.* (2013); Chen & Huang (2010); Prommon *et al.* (2012); Kruszynski *et al.* (2007).

### 2. Experimental

#### 2.1. Crystal data

$(C_9H_8N)_2[FeCl_4]Cl$   
 $M_r = 493.43$   
 Triclinic,  $P\bar{1}$   
 $a = 8.424$  (2) Å  
 $b = 10.435$  (3) Å  
 $c = 13.022$  (4) Å  
 $\alpha = 109.626$  (18)°  
 $\beta = 100.197$  (19)°

$\gamma = 90.893$  (19)°  
 $V = 1057.7$  (5) Å<sup>3</sup>  
 $Z = 2$   
 Mo  $K\alpha$  radiation  
 $\mu = 1.35$  mm<sup>-1</sup>  
 $T = 295$  K  
 $0.12 \times 0.05 \times 0.04$  mm

#### 2.2. Data collection

Bruker APEXII diffractometer  
 Absorption correction: multi-scan  
 (*SADABS*; Sheldrick, 2002)  
 $T_{min} = 0.899$ ,  $T_{max} = 0.922$

9378 measured reflections  
 3738 independent reflections  
 2927 reflections with  $I > 2\sigma(I)$   
 $R_{int} = 0.043$

#### 2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.033$   
 $wR(F^2) = 0.074$   
 $S = 1.01$   
 3738 reflections

235 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{max} = 0.29$  e Å<sup>-3</sup>  
 $\Delta\rho_{min} = -0.31$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$N1A-H1A \cdots Cl5^i$	0.86	2.16	3.014 (3)	174
$N1B-H1B \cdots Cl5$	0.86	2.21	3.043 (3)	163

Symmetry code: (i)  $-x + 1, -y + 1, -z + 2$ .

Data collection: *APEX2* (Bruker, 2011); cell refinement: *SAINTE* (Bruker, 2011); data reduction: *SAINTE*; program(s) used to solve structure: *SIR2002* (Burla *et al.*, 2005); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012) and *DIAMOND* (Brandenburg & Berndt, 2001); software used to prepare material for publication: *WinGX* (Farrugia, 2012).

### Acknowledgements

MESRS and DG-RSDT (Ministère de l'Enseignement Supérieur et de la Recherche Scientifique et la Direction Générale de la Recherche – Algérie) are thanked for financial support.

Supporting information for this paper is available from the IUCr electronic archives (Reference: CQ2018).

## References

- Bouchouit, K., Bendeif, E. E., EL Ouazzani, H., Dahaoui, S., Lecomte, C., Benali-cherif, N. & Sahraoui, B. (2010). *Chem. Phys.* **375**, 1–7.
- Bouchouit, K., Bougharraf, H., Derkowska-Zielinska, B., Benali-cherif, N. & Sahraoui, B. (2015). *Opt. Mater.* **48**, 215–221.
- Bouchouit, K., Essaidi, Z., Abed, S., Migalska-Zalas, A., Derkowska, B., Benali-cherif, N., Mihaly, M., Meghea, A. & Sahraoui, B. (2008). *Chem. Phys. Lett.* **455**, 270–274.
- Brandenburg, K. & Berndt, M. (2001). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2011). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Burla, M. C., Caliendo, R., Camalli, M., Carrozzini, B., Cascarano, G. L., De Caro, L., Giacovazzo, C., Polidori, G. & Spagna, R. (2005). *J. Appl. Cryst.* **38**, 381–388.
- Chen, L.-Z. & Huang, M.-N. (2010). *Acta Cryst.* **E66**, m377.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Jayalakshmi, D. & Kumar, J. (2006). *Cryst. Res. Technol.* **41**, 37–40.
- Khadri, A., Bouchene, R., Bouacida, S., Merazig, H. & Roisnel, T. (2013). *Acta Cryst.* **E69**, m190.
- Kruszynski, R., Wyrzykowski, D., Styczen, E. & Chmurzynski, L. (2007). *Acta Cryst.* **E63**, m2279–m2280.
- Prommon, P., Promseenong, P. & Chainok, K. (2012). *Acta Cryst.* **E68**, m211–m212.
- Sankar, R., Raghavan, C. M. & Jayavel, R. (2007). *Cryst. Growth Des.* **7**, 501–505.
- Sheldrick, G. M. (2002). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

## supporting information

*Acta Cryst.* (2015). E71, m273–m274 [https://doi.org/10.1107/S2056989015024548]

## Crystal structure of bis(quinolin-1-ium) tetrachloridoferrate(III) chloride

Azzedine Boudjarda, Karim Bouchouit, Samiha Arroudj, Sofiane Bouacida and Hocine Merazig

### S1. Comment

Hybrid compounds are one of the important categories of materials. They have received much attention in research areas including nonlinear optics, second harmonic generation (SHG), third harmonic generation (THG) and optical switching [Bouchouit *et al.* (2008); Bouchouit, *et al.* (2010); Jayalakshmi *et al.* (2006); Sankar *et al.* (2007); Bouchouit *et al.* (2015)]. A considerable number of hybrid organic/inorganic compounds have been extensively studied for their promising properties. Crystals of many of these materials can be grown from aqueous solution (Khadri *et al.* (2013); Chen *et al.* (2010); Prommon *et al.* (2012); Kruszynski *et al.* (2007)]. In the present work, a mixture of water and acetonitrile is used as solvent for the reaction of quinoline with iron (III) chloride and leads to the generation of crystals of bis-(quinolinium)tetrachloroferrate(III) chloride.

The asymmetric unit of the title hybrid compound consists of a tetrachloroferrate anion,  $(\text{FeCl}_4)^-$ , a chloride  $\text{Cl}^-$  anion and two quinolinium cations,  $(\text{C}_9\text{H}_8\text{N})^+$  (Fig. 1). The iron atom lies at the centre of a regular tetrahedron and it is coordinated to four Cl atoms with Fe—Cl bond lengths in the range 2.1862 (10) to 2.2013 (10) Å. The lengths of the C—C and C—N bonds in the two independent quinolinium cations are comparable to the related distances found in the literature. The quinolinium cations stack on top of each other, held together by  $\pi$ – $\pi$  interactions. The centroid to centroid distances between  $\text{C}_6$  and  $\text{C}_5\text{N}$  rings in adjacent stacked quinolinium cations are 3.609 (2) and 3.802 (2) Å.

The projection of the structure onto the *a-c* plane (Fig. 2) shows the N—H $\cdots$ Cl hydrogen bonding interactions between the N—H groups of the quinolinium cations and the  $\text{Cl}^-$  anions which, together with the  $\pi$ – $\pi$  interactions, serve to stabilise the structure.

### S2. Experimental

Quinoline,  $\text{C}_9\text{H}_7\text{N}$ , (0.2 mmol) and iron (III) chloride,  $\text{FeCl}_3$ , (0.1 mmol) were dissolved in a mixture of water (10 ml) and acetonitrile (10 ml) at ambient temperature over a period of approximately 30 minutes. After this period, a brown precipitate appeared which was removed by filtration. The filtrate was then left at room temperature until brown crystals appeared.

### S3. Refinement

All non-H atoms were refined with anisotropic atomic displacement parameters. The remaining H atoms were localized on Fourier maps but introduced in calculated positions and treated as riding on their parent atom (C and N) with C—H = 0.93 Å and N—H = 0.86 Å with  $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}$




Figure 1

An ORTEP-3 (Farrugia, 2012) plot of the title compound, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.




Figure 2

A packing diagram of the title compound, viewed along the *b* axis, showing the N—H...Cl hydrogen bonds as dashed lines.

### Bis(quinolin-1-ium) tetrachloridoferrate(III) chloride

#### Crystal data

$(C_9H_8N)_2[FeCl_4]Cl$

$M_r = 493.43$

Triclinic,  $P\bar{1}$

Hall symbol: -P 1

$a = 8.424$  (2) Å

$b = 10.435$  (3) Å

$c = 13.022$  (4) Å

$\alpha = 109.626$  (18)°

$\beta = 100.197$  (19)°

$\gamma = 90.893$  (19)°

$V = 1057.7$  (5) Å<sup>3</sup>

$Z = 2$

$F(000) = 498$   
 $D_x = 1.549 \text{ Mg m}^{-3}$   
 Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$   
 Cell parameters from 2434 reflections  
 $\theta = 2.5\text{--}24.9^\circ$

$\mu = 1.35 \text{ mm}^{-1}$   
 $T = 295 \text{ K}$   
 Prism, brown  
 $0.12 \times 0.05 \times 0.04 \text{ mm}$

*Data collection*

Bruker APEXII  
 diffractometer  
 Graphite monochromator  
 CCD rotation images, thin slices scans  
 Absorption correction: multi-scan  
 (SADABS; Sheldrick, 2002)  
 $T_{\min} = 0.899$ ,  $T_{\max} = 0.922$   
 9378 measured reflections

3738 independent reflections  
 2927 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.043$   
 $\theta_{\max} = 25.1^\circ$ ,  $\theta_{\min} = 3.1^\circ$   
 $h = -10 \rightarrow 9$   
 $k = -12 \rightarrow 12$   
 $l = -15 \rightarrow 15$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.033$   
 $wR(F^2) = 0.074$   
 $S = 1.01$   
 3738 reflections  
 235 parameters  
 0 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0268P)^2 + 0.0315P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 0.29 \text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.31 \text{ e \AA}^{-3}$

*Special details*

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional R-factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and R-factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Fe1	0.11617 (5)	-0.18619 (4)	0.64487 (3)	0.02192 (12)
Cl4	0.29503 (9)	-0.33664 (7)	0.60671 (6)	0.03500 (19)
Cl3	-0.12466 (8)	-0.28089 (7)	0.55023 (6)	0.02990 (18)
Cl1	0.17477 (8)	-0.00943 (7)	0.59992 (6)	0.02792 (17)
Cl5	0.40919 (9)	0.25683 (8)	1.05985 (6)	0.0361 (2)
Cl2	0.11071 (11)	-0.11756 (8)	0.82228 (6)	0.0407 (2)
N1A	0.8158 (3)	0.5396 (2)	0.83436 (18)	0.0227 (5)
H1A	0.7565	0.5992	0.8689	0.027*
N1B	0.4922 (3)	0.1123 (2)	0.83299 (18)	0.0263 (5)
H1B	0.4486	0.1429	0.8906	0.032*
C4A	0.9484 (3)	0.3350 (3)	0.8123 (2)	0.0209 (6)
C3B	0.6376 (3)	0.0137 (3)	0.6543 (2)	0.0286 (7)

H3B	0.689	-0.0202	0.5941	0.034*
C3A	1.0018 (3)	0.3528 (3)	0.7215 (2)	0.0256 (6)
H3A	1.0639	0.2886	0.6816	0.031*
C8B	0.4286 (3)	0.3164 (3)	0.7925 (2)	0.0269 (6)
H8B	0.3793	0.348	0.8539	0.032*
C9A	0.8504 (3)	0.4318 (3)	0.8695 (2)	0.0214 (6)
C2B	0.6258 (3)	-0.0606 (3)	0.7210 (2)	0.0317 (7)
H2B	0.6671	-0.1457	0.7061	0.038*
C6A	0.9269 (3)	0.2136 (3)	0.9376 (2)	0.0312 (7)
H6A	0.9539	0.1416	0.9625	0.037*
C7A	0.8267 (3)	0.3101 (3)	0.9917 (2)	0.0282 (7)
H7A	0.7859	0.2996	1.0505	0.034*
C7B	0.4349 (3)	0.3917 (3)	0.7256 (2)	0.0310 (7)
H7B	0.3901	0.4754	0.7418	0.037*
C4B	0.5727 (3)	0.1417 (3)	0.6753 (2)	0.0211 (6)
C9B	0.4972 (3)	0.1909 (3)	0.7676 (2)	0.0217 (6)
C5A	0.9848 (3)	0.2239 (3)	0.8496 (2)	0.0277 (6)
H5A	1.0486	0.1579	0.8137	0.033*
C1B	0.5514 (3)	-0.0082 (3)	0.8115 (2)	0.0319 (7)
H1B1	0.5431	-0.0585	0.8576	0.038*
C5B	0.5761 (3)	0.2235 (3)	0.6085 (2)	0.0274 (7)
H5B	0.6258	0.1942	0.5471	0.033*
C8A	0.7892 (3)	0.4185 (3)	0.9588 (2)	0.0244 (6)
H8A	0.724	0.4826	0.9951	0.029*
C1A	0.8687 (3)	0.5569 (3)	0.7502 (2)	0.0274 (7)
H1A1	0.8423	0.6326	0.7302	0.033*
C2A	0.9637 (3)	0.4630 (3)	0.6912 (2)	0.0282 (7)
H2A	1.0007	0.4752	0.6317	0.034*
C6B	0.5079 (3)	0.3445 (3)	0.6328 (2)	0.0309 (7)
H6B	0.5097	0.3965	0.5873	0.037*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Fe1	0.0269 (2)	0.0194 (2)	0.0219 (2)	0.00239 (16)	0.00657 (17)	0.00933 (17)
Cl4	0.0363 (4)	0.0294 (4)	0.0470 (5)	0.0126 (3)	0.0145 (4)	0.0193 (4)
Cl3	0.0271 (4)	0.0295 (4)	0.0319 (4)	-0.0008 (3)	0.0076 (3)	0.0082 (3)
Cl1	0.0317 (4)	0.0243 (4)	0.0322 (4)	0.0010 (3)	0.0065 (3)	0.0154 (3)
Cl5	0.0490 (5)	0.0359 (4)	0.0313 (4)	0.0214 (4)	0.0196 (4)	0.0149 (3)
Cl2	0.0691 (6)	0.0341 (4)	0.0230 (4)	0.0088 (4)	0.0136 (4)	0.0126 (3)
N1A	0.0248 (12)	0.0170 (12)	0.0259 (13)	0.0046 (9)	0.0097 (10)	0.0043 (10)
N1B	0.0307 (13)	0.0301 (14)	0.0209 (12)	0.0049 (11)	0.0092 (11)	0.0101 (11)
C4A	0.0192 (13)	0.0188 (14)	0.0216 (14)	0.0011 (11)	0.0015 (11)	0.0042 (12)
C3B	0.0278 (15)	0.0293 (17)	0.0266 (16)	0.0024 (13)	0.0097 (13)	0.0046 (13)
C3A	0.0248 (15)	0.0246 (16)	0.0260 (15)	0.0023 (12)	0.0104 (12)	0.0043 (13)
C8B	0.0255 (15)	0.0257 (16)	0.0272 (16)	0.0052 (12)	0.0086 (13)	0.0041 (13)
C9A	0.0180 (13)	0.0197 (15)	0.0228 (14)	-0.0033 (11)	0.0016 (12)	0.0041 (12)
C2B	0.0358 (17)	0.0210 (16)	0.0387 (18)	0.0083 (13)	0.0113 (15)	0.0085 (14)

C6A	0.0361 (17)	0.0306 (17)	0.0308 (17)	0.0038 (14)	0.0027 (14)	0.0175 (14)
C7A	0.0302 (16)	0.0335 (17)	0.0215 (15)	-0.0024 (13)	0.0030 (13)	0.0116 (13)
C7B	0.0296 (16)	0.0240 (16)	0.0372 (18)	0.0035 (13)	0.0004 (14)	0.0108 (14)
C4B	0.0190 (14)	0.0234 (15)	0.0186 (14)	-0.0006 (11)	0.0030 (11)	0.0048 (12)
C9B	0.0186 (14)	0.0222 (15)	0.0217 (14)	-0.0022 (11)	0.0011 (12)	0.0059 (12)
C5A	0.0278 (15)	0.0238 (16)	0.0315 (16)	0.0070 (12)	0.0041 (13)	0.0101 (13)
C1B	0.0357 (17)	0.0297 (17)	0.0359 (18)	0.0046 (14)	0.0097 (14)	0.0170 (14)
C5B	0.0268 (15)	0.0330 (17)	0.0223 (15)	-0.0012 (13)	0.0042 (13)	0.0099 (13)
C8A	0.0236 (14)	0.0273 (16)	0.0197 (14)	0.0036 (12)	0.0059 (12)	0.0039 (12)
C1A	0.0300 (16)	0.0217 (15)	0.0345 (17)	-0.0011 (12)	0.0081 (14)	0.0138 (13)
C2A	0.0302 (16)	0.0298 (17)	0.0296 (16)	-0.0002 (13)	0.0135 (13)	0.0127 (14)
C6B	0.0298 (16)	0.0327 (18)	0.0319 (17)	-0.0022 (13)	-0.0007 (14)	0.0169 (14)

*Geometric parameters (Å, °)*

Fe1—Cl2	2.1862 (10)	C9A—C8A	1.399 (4)
Fe1—Cl1	2.1880 (9)	C2B—C1B	1.386 (4)
Fe1—Cl4	2.1901 (10)	C2B—H2B	0.93
Fe1—Cl3	2.2013 (10)	C6A—C5A	1.356 (4)
N1A—C1A	1.317 (3)	C6A—C7A	1.408 (4)
N1A—C9A	1.367 (3)	C6A—H6A	0.93
N1A—H1A	0.86	C7A—C8A	1.360 (4)
N1B—C1B	1.319 (4)	C7A—H7A	0.93
N1B—C9B	1.370 (3)	C7B—C6B	1.398 (4)
N1B—H1B	0.86	C7B—H7B	0.93
C4A—C3A	1.404 (4)	C4B—C9B	1.407 (4)
C4A—C9A	1.412 (4)	C4B—C5B	1.411 (4)
C4A—C5A	1.419 (4)	C5A—H5A	0.93
C3B—C2B	1.359 (4)	C1B—H1B1	0.93
C3B—C4B	1.410 (4)	C5B—C6B	1.358 (4)
C3B—H3B	0.93	C5B—H5B	0.93
C3A—C2A	1.362 (4)	C8A—H8A	0.93
C3A—H3A	0.93	C1A—C2A	1.388 (4)
C8B—C7B	1.361 (4)	C1A—H1A1	0.93
C8B—C9B	1.400 (4)	C2A—H2A	0.93
C8B—H8B	0.93	C6B—H6B	0.93
Cl2—Fe1—Cl1	108.97 (4)	C8A—C7A—C6A	120.6 (3)
Cl2—Fe1—Cl4	110.06 (4)	C8A—C7A—H7A	119.7
Cl1—Fe1—Cl4	110.70 (4)	C6A—C7A—H7A	119.7
Cl2—Fe1—Cl3	108.87 (4)	C8B—C7B—C6B	120.7 (3)
Cl1—Fe1—Cl3	109.03 (4)	C8B—C7B—H7B	119.6
Cl4—Fe1—Cl3	109.18 (4)	C6B—C7B—H7B	119.6
C1A—N1A—C9A	123.3 (2)	C9B—C4B—C3B	118.3 (2)
C1A—N1A—H1A	118.3	C9B—C4B—C5B	117.5 (2)
C9A—N1A—H1A	118.3	C3B—C4B—C5B	124.2 (2)
C1B—N1B—C9B	122.9 (2)	N1B—C9B—C8B	120.6 (2)
C1B—N1B—H1B	118.5	N1B—C9B—C4B	118.2 (2)



C9B—N1B—H1B	118.5	C8B—C9B—C4B	121.2 (2)
C3A—C4A—C9A	118.6 (2)	C6A—C5A—C4A	120.1 (3)
C3A—C4A—C5A	123.9 (3)	C6A—C5A—H5A	119.9
C9A—C4A—C5A	117.6 (2)	C4A—C5A—H5A	119.9
C2B—C3B—C4B	120.7 (3)	N1B—C1B—C2B	120.6 (3)
C2B—C3B—H3B	119.7	N1B—C1B—H1B1	119.7
C4B—C3B—H3B	119.7	C2B—C1B—H1B1	119.7
C2A—C3A—C4A	120.6 (3)	C6B—C5B—C4B	120.8 (3)
C2A—C3A—H3A	119.7	C6B—C5B—H5B	119.6
C4A—C3A—H3A	119.7	C4B—C5B—H5B	119.6
C7B—C8B—C9B	119.1 (3)	C7A—C8A—C9A	118.9 (3)
C7B—C8B—H8B	120.4	C7A—C8A—H8A	120.5
C9B—C8B—H8B	120.4	C9A—C8A—H8A	120.5
N1A—C9A—C8A	120.6 (2)	N1A—C1A—C2A	120.5 (3)
N1A—C9A—C4A	117.8 (2)	N1A—C1A—H1A1	119.7
C8A—C9A—C4A	121.6 (2)	C2A—C1A—H1A1	119.7
C3B—C2B—C1B	119.3 (3)	C3A—C2A—C1A	119.2 (3)
C3B—C2B—H2B	120.4	C3A—C2A—H2A	120.4
C1B—C2B—H2B	120.4	C1A—C2A—H2A	120.4
C5A—C6A—C7A	121.1 (3)	C5B—C6B—C7B	120.6 (3)
C5A—C6A—H6A	119.4	C5B—C6B—H6B	119.7
C7A—C6A—H6A	119.4	C7B—C6B—H6B	119.7
C9A—C4A—C3A—C2A	1.8 (4)	C5B—C4B—C9B—N1B	179.7 (2)
C5A—C4A—C3A—C2A	-179.2 (3)	C3B—C4B—C9B—C8B	-179.3 (2)
C1A—N1A—C9A—C8A	-179.4 (2)	C5B—C4B—C9B—C8B	-0.2 (4)
C1A—N1A—C9A—C4A	0.2 (4)	C7A—C6A—C5A—C4A	1.6 (4)
C3A—C4A—C9A—N1A	-1.4 (4)	C3A—C4A—C5A—C6A	-179.3 (3)
C5A—C4A—C9A—N1A	179.6 (2)	C9A—C4A—C5A—C6A	-0.3 (4)
C3A—C4A—C9A—C8A	178.2 (2)	C9B—N1B—C1B—C2B	1.3 (4)
C5A—C4A—C9A—C8A	-0.8 (4)	C3B—C2B—C1B—N1B	0.1 (4)
C4B—C3B—C2B—C1B	-1.1 (4)	C9B—C4B—C5B—C6B	-0.4 (4)
C5A—C6A—C7A—C8A	-1.8 (4)	C3B—C4B—C5B—C6B	178.6 (3)
C9B—C8B—C7B—C6B	0.4 (4)	C6A—C7A—C8A—C9A	0.7 (4)
C2B—C3B—C4B—C9B	0.7 (4)	N1A—C9A—C8A—C7A	-179.8 (2)
C2B—C3B—C4B—C5B	-178.3 (3)	C4A—C9A—C8A—C7A	0.6 (4)
C1B—N1B—C9B—C8B	178.2 (3)	C9A—N1A—C1A—C2A	0.6 (4)
C1B—N1B—C9B—C4B	-1.7 (4)	C4A—C3A—C2A—C1A	-1.0 (4)
C7B—C8B—C9B—N1B	-179.7 (2)	N1A—C1A—C2A—C3A	-0.1 (4)
C7B—C8B—C9B—C4B	0.2 (4)	C4B—C5B—C6B—C7B	1.0 (4)
C3B—C4B—C9B—N1B	0.6 (4)	C8B—C7B—C6B—C5B	-1.0 (4)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
N1A—H1A $\cdots$ Cl5 <sup>i</sup>	0.86	2.16	3.014 (3)	174



---

N1B—H1B···Cl5	0.86	2.21	3.043 (3)	163
---------------	------	------	-----------	-----

---

Symmetry code: (i)  $-x+1, -y+1, -z+2$ .