



# Crystal structure of 1,3-bis[[4-(acetylsulfanyl)phenyl]ethynyl]azulene

Sebastian Förster, Wilhelm Seichter and Edwin Weber\*

Institut für Organische Chemie, TU Bergakademie Freiberg, Leipziger Strasse, 29, D-09596 Freiberg/Sachsen, Germany. \*Correspondence e-mail: edwin.weber@chemie.tu-freiberg.de

Received 19 November 2015; accepted 7 January 2016

Edited by W. T. A. Harrison, University of Aberdeen, Scotland


In the title compound,  $C_{30}H_{20}O_2S_2$ , the dihedral angles between the central azulene ring system (r.m.s. deviation = 0.039 Å) and the pendant benzene rings are 28.96 (7) and 55.15 (7)°. The dihedral angles between the benzene rings and their attached acetylsulfanyl groups are 59.60 (10) and 84.79 (10)°. The expected  $\pi$ - $\pi$  stacking interactions are not observed in the crystal structure; instead, the packing features C—H...O hydrogen bonds, which link the molecules into C(12) [010] chains, which are supported by weak C—H... $\pi$  contacts.

**Keywords:** crystal structure; azulene; 1,3-disubstitution; C—H...O hydrogen bond; C—H... $\pi$  interaction.

**CCDC reference:** 1445850

## 1. Related literature

For background to this work, see: Wang *et al.* (2009); Puodziukynaite *et al.* (2014); Xia *et al.* (2014). For the synthesis and related structures, see: Förster *et al.* (2012, 2014).



## 2. Experimental

### 2.1. Crystal data

$C_{30}H_{20}O_2S_2$	$a = 13.7674$ (3) Å
$M_r = 476.58$	$b = 8.9849$ (2) Å
Monoclinic, $P2_1/n$	$c = 19.7586$ (4) Å

$\beta = 104.022$  (1)°  
 $V = 2371.28$  (9) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation

$\mu = 0.25$  mm<sup>-1</sup>  
 $T = 100$  K  
 $0.24 \times 0.23 \times 0.15$  mm

### 2.2. Data collection

Bruker Kappa APEX CCD diffractometer  
 Absorption correction: multi-scan (SADABS; Bruker, 2008)  
 $T_{\min} = 0.942$ ,  $T_{\max} = 0.963$

36340 measured reflections  
 5898 independent reflections  
 4422 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.039$

### 2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$   
 $wR(F^2) = 0.130$   
 $S = 1.03$   
 5898 reflections

309 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.44$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.27$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

Cg1 is the mid-point of the C11—C12 bond and Cg2 is the centroid of the C1—C4/C10 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C2—H2...O1 <sup>i</sup>	0.95	2.40	3.285 (3)	155
C17—H1...Cg1	0.95	2.69	3.612 (3)	165
C20—H20A...Cg2 <sup>ii</sup>	0.98	2.89	3.835 (2)	162

Symmetry code: (i)  $-x + \frac{5}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$ ; (ii)  $x + \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$ .

Data collection: APEX2 (Bruker AXS); cell refinement: SAINT (Sheldrick, 2008); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL2015 (Sheldrick, 2015); molecular graphics: ORTEP-3 for Windows (Farrugia, 2012); software used to prepare material for publication: SHELXTL (Sheldrick, 2008).

## Acknowledgements

This work has been performed within the 'Cluster of Excellence Structure Design of Novel High-Performance Materials via Atomic Design and Defect Engineering' (ADDE), which was supported financially by the European Union (European Regional Development Fund) and by the Ministry of Science and Art of Saxony (SMWK).

Supporting information for this paper is available from the IUCr electronic archives (Reference: HB7546).

## References

- Bruker (2008). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.  
 Förster, S., Hahn, T., Loose, C., Röder, C., Liebing, S., Seichter, W., Eissmann, F., Kortus, J. & Weber, E. (2012). *J. Phys. Org. Chem.* **25**, 856–863.  
 Förster, S., Seichter, W., Kuhnert, R. & Weber, E. (2014). *J. Mol. Struct.* **1075**, 63–70.  
 Puodziukynaite, E., Wang, H.-W., Lawrence, J., Wise, A., Russell, T. P., Barnes, M. D. & Emrick, T. (2014). *J. Am. Chem. Soc.* **136**, 11043–11049.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.

Wang, X., Ng, J. K.-P., Jia, P., Lin, T., Cho, C. M., Xu, J., Lu, X. & He, C. (2009). *Macromolecules*, **42**, 5534–5544.

Xia, J., Capozzi, B., Wei, S., Strange, M., Batra, A., Moreno, J. R., Amir, E., Amir, R., Solomon, G. C., Venkataraman, L. & Campos, L. M. (2014). *Nano Lett.* **14**, 2941–2945.

## supporting information

*Acta Cryst.* (2015). E71, o1099–o1100 [https://doi.org/10.1107/S2056989016000323]

## Crystal structure of 1,3-bis{[4-(acetylsulfanyl)phenyl]ethynyl}azulene

Sebastian Förster, Wilhelm Seichter and Edwin Weber

### S1. Comment

Azulene derivatives offer a number of interesting applications especially in the field of molecular electronics (Wang *et al.*, 2009; Puodziukynaite *et al.*, 2014). It ties up with the fact that the non-alternating azulene possesses remarkable electronic and optical properties (Xia *et al.*, 2014). Although the title compound, C<sub>30</sub>H<sub>20</sub>O<sub>2</sub>S<sub>2</sub>, (I), is fully conjugated, no flat molecular structure can be observed. Both phenyl rings, fig 1, are rotated out of the plane containing the azulene core [*phenyl*(C13—C18) 29.0°, *phenyl*(C23—C28) 55.2°]. The C—S—C angle of the acetyl protected thiol is slightly smaller compared to that found in an analogous compound, 1,3-bis[4-(*tert*-butylsulfanyl)phenylethynyl]azulene, featuring a *tert*-butyl protection group at the sulfur atom (Förster *et al.*, 2012). Unlike the previous case, no  $\pi\cdots\pi$  interactions are present in the title compound. In all probability, this uncommon phenomenon within the substance class of azulenes (Förster *et al.*, 2014) is related to the non-planar molecular structure and may be caused from packing effects. However, the crystal structure is based on C—H $\cdots$ O hydrogen bonds [C2—H2 $\cdots$ O1 (2.5-x, -0.5+y, 1.5-z; 2.40 Å, 155.0°)] and C—H $\cdots\pi$  interactions [C17—H17 $\cdots$ Cg(1) 2.69 Å, 164.9°; C20—H20A $\cdots$ Cg(2) 2.89 Å, 159.0°].

### S2. Experimental

#### S2.1. Synthesis and crystallization

The synthesis of the title compound, (I), has already been described (Förster *et al.* 2012). The crystals were grown from toluene solution by slow evaporation.

#### S2.2. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 1. The hydrogen atoms attached to C were fixed geometrically and treated as riding atoms, with d(C—H) = 0.93 and U<sub>iso</sub>(H) = 1.2 U<sub>eq</sub>(C) for aromatic and U<sub>iso</sub>(H) = 1.5 U<sub>eq</sub>(C) for methyl groups.




Figure 1  
Ellipsoid plot.




Figure 2  
Packing diagram.

### 1,3-Bis[4-(acetysulfanyl)phenyl]ethynyl]azulene

#### Crystal data

$C_{30}H_{20}O_2S_2$   
 $M_r = 476.58$   
 Monoclinic,  $P2_1/n$   
 $a = 13.7674 (3) \text{ \AA}$   
 $b = 8.9849 (2) \text{ \AA}$   
 $c = 19.7586 (4) \text{ \AA}$   
 $\beta = 104.022 (1)^\circ$   
 $V = 2371.28 (9) \text{ \AA}^3$   
 $Z = 4$

$F(000) = 992$   
 $D_x = 1.335 \text{ Mg m}^{-3}$   
 Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$   
 Cell parameters from 8057 reflections  
 $\theta = 2.5\text{--}28.0^\circ$   
 $\mu = 0.25 \text{ mm}^{-1}$   
 $T = 100 \text{ K}$   
 Irregular, green  
 $0.24 \times 0.23 \times 0.15 \text{ mm}$

#### Data collection

Bruker Kappa APEX CCD  
 diffractometer  
 phi and  $\omega$  scans  
 Absorption correction: multi-scan  
 (SADABS; Bruker, 2008)  
 $T_{\min} = 0.942$ ,  $T_{\max} = 0.963$   
 36340 measured reflections

5898 independent reflections  
 4422 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.039$   
 $\theta_{\max} = 28.3^\circ$ ,  $\theta_{\min} = 2.5^\circ$   
 $h = -18 \rightarrow 18$   
 $k = -12 \rightarrow 11$   
 $l = -26 \rightarrow 26$

*Refinement*Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.046$  $wR(F^2) = 0.130$  $S = 1.03$ 

5898 reflections

309 parameters

0 restraints

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0658P)^2 + 1.347P]$ where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\max} = 0.001$  $\Delta\rho_{\max} = 0.44 \text{ e } \text{\AA}^{-3}$  $\Delta\rho_{\min} = -0.27 \text{ e } \text{\AA}^{-3}$ *Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	1.37569 (4)	0.11544 (6)	0.58441 (3)	0.03405 (15)
S2	0.55913 (4)	-0.42293 (6)	1.10153 (3)	0.03144 (14)
O1	1.43557 (12)	0.39449 (19)	0.60090 (9)	0.0418 (4)
O2	0.52513 (12)	-0.17076 (16)	1.16136 (8)	0.0361 (4)
C1	0.93413 (13)	0.2985 (2)	0.81129 (9)	0.0229 (4)
C2	0.91025 (14)	0.1731 (2)	0.84616 (10)	0.0248 (4)
H2	0.9425	0.0791	0.8485	0.030*
C3	0.83172 (14)	0.2075 (2)	0.87699 (10)	0.0235 (4)
C4	0.80547 (13)	0.3585 (2)	0.86356 (9)	0.0224 (4)
C5	0.73471 (14)	0.4346 (2)	0.88954 (10)	0.0258 (4)
H5	0.7021	0.3782	0.9182	0.031*
C6	0.70505 (15)	0.5828 (2)	0.87924 (11)	0.0286 (4)
H6	0.6574	0.6156	0.9036	0.034*
C7	0.73635 (14)	0.6887 (2)	0.83792 (10)	0.0281 (4)
H7	0.7048	0.7831	0.8366	0.034*
C8	0.80679 (15)	0.6777 (2)	0.79820 (10)	0.0279 (4)
H8	0.8147	0.7643	0.7725	0.033*
C9	0.86748 (14)	0.5575 (2)	0.79079 (10)	0.0244 (4)
H9	0.9129	0.5744	0.7623	0.029*
C10	0.86986 (13)	0.4173 (2)	0.81964 (9)	0.0217 (4)
C11	1.01077 (14)	0.2995 (2)	0.77358 (10)	0.0235 (4)
C12	1.07506 (14)	0.2876 (2)	0.74377 (10)	0.0252 (4)
C13	1.15323 (13)	0.2606 (2)	0.70827 (10)	0.0222 (4)
C14	1.15694 (14)	0.3358 (2)	0.64730 (10)	0.0261 (4)
H14	1.1120	0.4155	0.6312	0.031*
C15	1.22577 (14)	0.2948 (2)	0.61000 (10)	0.0272 (4)
H15	1.2263	0.3440	0.5675	0.033*
C16	1.29404 (13)	0.1821 (2)	0.63447 (10)	0.0242 (4)
C17	1.29573 (14)	0.1142 (2)	0.69783 (10)	0.0270 (4)
H17	1.3454	0.0420	0.7163	0.032*

C18	1.22522 (14)	0.1517 (2)	0.73400 (10)	0.0269 (4)
H18	1.2256	0.1031	0.7768	0.032*
C19	1.44675 (15)	0.2760 (3)	0.57564 (11)	0.0328 (5)
C20	1.52157 (17)	0.2459 (3)	0.53316 (12)	0.0483 (7)
H20A	1.4865	0.2112	0.4866	0.072*
H20B	1.5688	0.1692	0.5562	0.072*
H20C	1.5582	0.3375	0.5289	0.072*
C21	0.78423 (15)	0.1054 (2)	0.91530 (9)	0.0243 (4)
C22	0.74797 (15)	0.0199 (2)	0.94649 (10)	0.0275 (4)
C23	0.70236 (14)	-0.0862 (2)	0.98362 (10)	0.0246 (4)
C24	0.76034 (15)	-0.1665 (2)	1.03950 (10)	0.0284 (4)
H24	0.8306	-0.1504	1.0535	0.034*
C25	0.71595 (15)	-0.2691 (2)	1.07442 (11)	0.0289 (4)
H25	0.7557	-0.3248	1.1119	0.035*
C26	0.61271 (15)	-0.2908 (2)	1.05464 (10)	0.0268 (4)
C27	0.55486 (15)	-0.2111 (2)	1.00026 (11)	0.0318 (5)
H27	0.4845	-0.2261	0.9870	0.038*
C28	0.59931 (15)	-0.1085 (2)	0.96468 (11)	0.0308 (4)
H28	0.5592	-0.0532	0.9272	0.037*
C29	0.52713 (14)	-0.3041 (2)	1.16535 (10)	0.0261 (4)
C30	0.50479 (17)	-0.3905 (2)	1.22472 (11)	0.0341 (5)
H30A	0.5636	-0.3895	1.2643	0.051*
H30B	0.4883	-0.4935	1.2100	0.051*
H30C	0.4479	-0.3451	1.2387	0.051*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0321 (3)	0.0331 (3)	0.0435 (3)	0.0021 (2)	0.0220 (2)	-0.0023 (2)
S2	0.0425 (3)	0.0203 (2)	0.0379 (3)	-0.0023 (2)	0.0221 (2)	0.0025 (2)
O1	0.0341 (8)	0.0477 (10)	0.0446 (9)	-0.0138 (7)	0.0117 (7)	-0.0013 (8)
O2	0.0478 (9)	0.0248 (8)	0.0381 (8)	0.0010 (7)	0.0152 (7)	0.0001 (6)
C1	0.0234 (9)	0.0222 (9)	0.0244 (9)	-0.0025 (7)	0.0085 (7)	-0.0004 (7)
C2	0.0276 (9)	0.0225 (9)	0.0260 (9)	-0.0003 (7)	0.0097 (7)	-0.0001 (8)
C3	0.0246 (9)	0.0238 (9)	0.0233 (9)	-0.0034 (7)	0.0083 (7)	0.0004 (7)
C4	0.0224 (8)	0.0239 (9)	0.0217 (9)	-0.0023 (7)	0.0069 (7)	-0.0007 (7)
C5	0.0232 (9)	0.0301 (10)	0.0266 (9)	-0.0025 (8)	0.0108 (7)	-0.0010 (8)
C6	0.0259 (9)	0.0312 (11)	0.0309 (10)	0.0015 (8)	0.0114 (8)	-0.0043 (8)
C7	0.0273 (9)	0.0266 (10)	0.0293 (10)	0.0039 (8)	0.0048 (8)	-0.0027 (8)
C8	0.0314 (10)	0.0241 (10)	0.0277 (10)	0.0006 (8)	0.0062 (8)	0.0041 (8)
C9	0.0259 (9)	0.0261 (10)	0.0224 (9)	-0.0024 (7)	0.0082 (7)	0.0002 (7)
C10	0.0208 (8)	0.0237 (9)	0.0213 (8)	-0.0023 (7)	0.0067 (7)	-0.0012 (7)
C11	0.0265 (9)	0.0180 (9)	0.0269 (9)	-0.0045 (7)	0.0083 (7)	-0.0018 (7)
C12	0.0272 (9)	0.0179 (9)	0.0309 (10)	-0.0011 (7)	0.0078 (8)	-0.0020 (7)
C13	0.0222 (8)	0.0193 (9)	0.0267 (9)	-0.0021 (7)	0.0093 (7)	-0.0034 (7)
C14	0.0234 (9)	0.0230 (10)	0.0331 (10)	0.0033 (7)	0.0093 (8)	0.0033 (8)
C15	0.0283 (9)	0.0271 (10)	0.0284 (10)	0.0010 (8)	0.0111 (8)	0.0059 (8)
C16	0.0217 (8)	0.0241 (10)	0.0293 (9)	-0.0010 (7)	0.0113 (7)	-0.0016 (8)

C17	0.0244 (9)	0.0240 (10)	0.0335 (10)	0.0033 (7)	0.0085 (8)	0.0027 (8)
C18	0.0293 (10)	0.0256 (10)	0.0279 (10)	0.0009 (8)	0.0108 (8)	0.0023 (8)
C19	0.0232 (9)	0.0482 (14)	0.0265 (10)	-0.0047 (9)	0.0051 (8)	0.0029 (9)
C20	0.0271 (11)	0.084 (2)	0.0374 (12)	-0.0038 (12)	0.0152 (9)	0.0081 (13)
C21	0.0329 (10)	0.0213 (9)	0.0204 (9)	0.0035 (8)	0.0095 (7)	-0.0006 (7)
C22	0.0301 (10)	0.0246 (10)	0.0304 (10)	0.0002 (8)	0.0126 (8)	-0.0050 (8)
C23	0.0321 (10)	0.0181 (9)	0.0284 (9)	-0.0015 (7)	0.0169 (8)	-0.0030 (7)
C24	0.0273 (9)	0.0285 (10)	0.0318 (10)	-0.0006 (8)	0.0115 (8)	0.0010 (8)
C25	0.0325 (10)	0.0262 (10)	0.0304 (10)	0.0028 (8)	0.0124 (8)	0.0046 (8)
C26	0.0357 (10)	0.0184 (9)	0.0314 (10)	-0.0013 (8)	0.0181 (8)	0.0002 (8)
C27	0.0253 (9)	0.0294 (11)	0.0424 (12)	-0.0021 (8)	0.0117 (8)	0.0057 (9)
C28	0.0309 (10)	0.0271 (10)	0.0358 (11)	0.0019 (8)	0.0110 (8)	0.0096 (9)
C29	0.0248 (9)	0.0233 (10)	0.0311 (10)	-0.0001 (7)	0.0085 (8)	0.0012 (8)
C30	0.0416 (12)	0.0305 (11)	0.0355 (11)	0.0012 (9)	0.0195 (9)	0.0028 (9)

*Geometric parameters (Å, °)*

S1—C16	1.7716 (19)	C14—C15	1.384 (3)
S1—C19	1.775 (2)	C14—H14	0.9500
S2—C26	1.7727 (19)	C15—C16	1.387 (3)
S2—C29	1.787 (2)	C15—H15	0.9500
O1—C19	1.201 (3)	C16—C17	1.387 (3)
O2—C29	1.201 (2)	C17—C18	1.380 (3)
C1—C2	1.401 (3)	C17—H17	0.9500
C1—C10	1.421 (3)	C18—H18	0.9500
C1—C11	1.432 (2)	C19—C20	1.503 (3)
C2—C3	1.398 (3)	C20—H20A	0.9800
C2—H2	0.9500	C20—H20B	0.9800
C3—C4	1.413 (3)	C20—H20C	0.9800
C3—C21	1.443 (3)	C21—C22	1.170 (3)
C4—C5	1.386 (3)	C22—C23	1.437 (3)
C4—C10	1.480 (2)	C23—C28	1.391 (3)
C5—C6	1.393 (3)	C23—C24	1.396 (3)
C5—H5	0.9500	C24—C25	1.380 (3)
C6—C7	1.388 (3)	C24—H24	0.9500
C6—H6	0.9500	C25—C26	1.394 (3)
C7—C8	1.391 (3)	C25—H25	0.9500
C7—H7	0.9500	C26—C27	1.374 (3)
C8—C9	1.394 (3)	C27—C28	1.388 (3)
C8—H8	0.9500	C27—H27	0.9500
C9—C10	1.380 (3)	C28—H28	0.9500
C9—H9	0.9500	C29—C30	1.500 (3)
C11—C12	1.181 (3)	C30—H30A	0.9800
C12—C13	1.440 (3)	C30—H30B	0.9800
C13—C14	1.393 (3)	C30—H30C	0.9800
C13—C18	1.398 (3)		
C16—S1—C19	102.66 (10)	C17—C16—S1	118.76 (15)

C26—S2—C29	100.01 (9)	C18—C17—C16	120.01 (18)
C2—C1—C10	108.67 (16)	C18—C17—H17	120.0
C2—C1—C11	123.48 (17)	C16—C17—H17	120.0
C10—C1—C11	127.85 (17)	C17—C18—C13	120.56 (18)
C3—C2—C1	109.57 (17)	C17—C18—H18	119.7
C3—C2—H2	125.2	C13—C18—H18	119.7
C1—C2—H2	125.2	O1—C19—C20	124.5 (2)
C2—C3—C4	108.68 (16)	O1—C19—S1	123.40 (16)
C2—C3—C21	125.77 (18)	C20—C19—S1	112.13 (18)
C4—C3—C21	125.55 (17)	C19—C20—H20A	109.5
C5—C4—C3	125.09 (17)	C19—C20—H20B	109.5
C5—C4—C10	128.00 (18)	H20A—C20—H20B	109.5
C3—C4—C10	106.88 (16)	C19—C20—H20C	109.5
C4—C5—C6	128.65 (18)	H20A—C20—H20C	109.5
C4—C5—H5	115.7	H20B—C20—H20C	109.5
C6—C5—H5	115.7	C22—C21—C3	178.1 (2)
C7—C6—C5	128.24 (18)	C21—C22—C23	178.9 (2)
C7—C6—H6	115.9	C28—C23—C24	119.25 (17)
C5—C6—H6	115.9	C28—C23—C22	120.10 (18)
C6—C7—C8	129.88 (19)	C24—C23—C22	120.64 (18)
C6—C7—H7	115.1	C25—C24—C23	120.20 (18)
C8—C7—H7	115.1	C25—C24—H24	119.9
C7—C8—C9	129.18 (19)	C23—C24—H24	119.9
C7—C8—H8	115.4	C24—C25—C26	119.92 (19)
C9—C8—H8	115.4	C24—C25—H25	120.0
C10—C9—C8	128.20 (18)	C26—C25—H25	120.0
C10—C9—H9	115.9	C27—C26—C25	120.30 (18)
C8—C9—H9	115.9	C27—C26—S2	121.48 (15)
C9—C10—C1	126.22 (17)	C25—C26—S2	118.21 (15)
C9—C10—C4	127.60 (17)	C26—C27—C28	119.96 (19)
C1—C10—C4	106.17 (16)	C26—C27—H27	120.0
C12—C11—C1	174.3 (2)	C28—C27—H27	120.0
C11—C12—C13	175.5 (2)	C27—C28—C23	120.36 (19)
C14—C13—C18	118.85 (17)	C27—C28—H28	119.8
C14—C13—C12	121.86 (17)	C23—C28—H28	119.8
C18—C13—C12	119.23 (17)	O2—C29—C30	124.21 (19)
C15—C14—C13	120.34 (18)	O2—C29—S2	123.75 (16)
C15—C14—H14	119.8	C30—C29—S2	112.03 (14)
C13—C14—H14	119.8	C29—C30—H30A	109.5
C14—C15—C16	120.16 (17)	C29—C30—H30B	109.5
C14—C15—H15	119.9	H30A—C30—H30B	109.5
C16—C15—H15	119.9	C29—C30—H30C	109.5
C15—C16—C17	119.82 (17)	H30A—C30—H30C	109.5
C15—C16—S1	121.35 (15)	H30B—C30—H30C	109.5
C10—C1—C2—C3	0.1 (2)	C13—C14—C15—C16	-2.4 (3)
C11—C1—C2—C3	-179.26 (17)	C14—C15—C16—C17	-2.5 (3)
C1—C2—C3—C4	-1.4 (2)	C14—C15—C16—S1	174.31 (15)



C1—C2—C3—C21	177.71 (18)	C19—S1—C16—C15	61.42 (18)
C2—C3—C4—C5	-175.89 (18)	C19—S1—C16—C17	-121.78 (17)
C21—C3—C4—C5	5.0 (3)	C15—C16—C17—C18	4.4 (3)
C2—C3—C4—C10	2.2 (2)	S1—C16—C17—C18	-172.40 (15)
C21—C3—C4—C10	-176.99 (17)	C16—C17—C18—C13	-1.6 (3)
C3—C4—C5—C6	179.6 (2)	C14—C13—C18—C17	-3.1 (3)
C10—C4—C5—C6	1.9 (3)	C12—C13—C18—C17	174.31 (18)
C4—C5—C6—C7	3.0 (4)	C16—S1—C19—O1	0.8 (2)
C5—C6—C7—C8	-2.4 (4)	C16—S1—C19—C20	-179.13 (15)
C6—C7—C8—C9	-2.1 (4)	C28—C23—C24—C25	1.4 (3)
C7—C8—C9—C10	2.4 (4)	C22—C23—C24—C25	-179.43 (18)
C8—C9—C10—C1	-179.16 (19)	C23—C24—C25—C26	-1.1 (3)
C8—C9—C10—C4	2.5 (3)	C24—C25—C26—C27	0.3 (3)
C2—C1—C10—C9	-177.39 (18)	C24—C25—C26—S2	-179.55 (15)
C11—C1—C10—C9	1.9 (3)	C29—S2—C26—C27	-86.86 (18)
C2—C1—C10—C4	1.2 (2)	C29—S2—C26—C25	92.98 (17)
C11—C1—C10—C4	-179.46 (18)	C25—C26—C27—C28	0.1 (3)
C5—C4—C10—C9	-5.5 (3)	S2—C26—C27—C28	179.98 (16)
C3—C4—C10—C9	176.53 (18)	C26—C27—C28—C23	0.2 (3)
C5—C4—C10—C1	175.89 (19)	C24—C23—C28—C27	-1.0 (3)
C3—C4—C10—C1	-2.1 (2)	C22—C23—C28—C27	179.86 (19)
C18—C13—C14—C15	5.1 (3)	C26—S2—C29—O2	14.4 (2)
C12—C13—C14—C15	-172.25 (18)	C26—S2—C29—C30	-164.72 (15)

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

Cg1 is the mid-point of C11—C12 and Cg2 is the centroid of the C1—C4/C10 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C2—H2 $\cdots$ O1 <sup>i</sup>	0.95	2.40	3.285 (3)	155
C17—H1 $\cdots$ Cg1	0.95	2.69	3.612 (3)	165
C20—H20A $\cdots$ Cg2 <sup>ii</sup>	0.98	2.89	3.835 (2)	162

Symmetry codes: (i)  $-x+5/2, y-1/2, -z+3/2$ ; (ii)  $x+1/2, -y+1/2, z-1/2$ .