

# Crystal structures of four chiral imine-substituted thiophene derivatives

Guadalupe Hernández-Téllez,<sup>a</sup> Sylvain Bernès,<sup>b\*</sup> Angel Mendoza,<sup>c</sup> Francisco Javier Ríos-Merino,<sup>c</sup> Gloria E. Moreno,<sup>a</sup> Oscar Portillo<sup>a</sup> and René Gutiérrez<sup>a</sup>

<sup>a</sup>Laboratorio de Síntesis de Complejos, Facultad de Ciencias Químicas, Universidad Autónoma de Puebla, A.P. 1067, 72001 Puebla, Pue., Mexico, <sup>b</sup>Instituto de Física, Universidad Autónoma de Puebla, Av. San Claudio y 18 Sur, 72570 Puebla, Pue., Mexico, and <sup>c</sup>Centro de Química, Instituto de Ciencias, Universidad Autónoma de Puebla, 72570 Puebla, Pue., Mexico. \*Correspondence e-mail: sylvain\_bernes@hotmail.com

Received 1 February 2016

Accepted 10 February 2016

Edited by J. Simpson, University of Otago, New Zealand

**Keywords:** crystal structure; Schiff base; bis-imine; thiophene.


**CCDC references:** 1452795; 1452794; 1452793; 1452792


**Supporting information:** this article has supporting information at journals.iucr.org/e

A series of thiophenes substituted in positions 2 and 5 by imine groups have been synthesized using a solvent-free approach, and their crystal structures determined. The substituents are chiral groups, and the expected absolute configuration for each molecule was confirmed by refinement of the Flack parameter. The compounds are 2,5-bis[(*S*)-(+)-(1,2,3,4-tetrahydronaphthalen-1-yl)imino]thiophene, C<sub>26</sub>H<sub>26</sub>N<sub>2</sub>S, (I), 2,5-bis[[(*R*)-(–)-1-(4-methoxyphenyl)ethyl]iminomethyl]thiophene, C<sub>24</sub>H<sub>26</sub>N<sub>2</sub>O<sub>2</sub>S, (II), 2,5-bis[[(*R*)-(–)-1-(4-fluorophenyl)ethyl]iminomethyl]thiophene, C<sub>22</sub>H<sub>20</sub>F<sub>2</sub>N<sub>2</sub>S, (III), and 2,5-bis[[(*S*)-(+)-1-(4-chlorophenyl)ethyl]iminomethyl]thiophene, C<sub>22</sub>H<sub>20</sub>Cl<sub>2</sub>N<sub>2</sub>S, (IV). A common feature of all four molecules is the presence of twofold symmetry. For (I), which crystallizes in the triclinic space group *P*1, this symmetry is non-crystallographic, but for (II) in *C*2 and the isomorphous structures (III) and (IV) that crystallize in *P*2<sub>1</sub>2<sub>1</sub>2, the twofold symmetry is crystallographically imposed with one half of each molecule in the asymmetric unit. The comparable molecular symmetry in the four structures is also reflected in similar packing, with molecules aggregated to form chains through weak C–H···S interactions.

## 1. Chemical context

Thiophenedicarbaldehydes have a variety of applications (Dean, 1982*a,b*), for instance in the synthesis of annulenones and polyenyl-substituted thiophenes (Sargent & Cresp, 1975), in the preparation of macrocyclic ligands for bimetallic complexes that are able to mimic enzymes (Nelson *et al.*, 1983), in crown ether chemistry (Cram & Trueblood, 1981) and, more recently, in the preparation of azomethines for photovoltaic applications (Bolduc *et al.*, 2013*a,b*; Petrus *et al.*, 2014). In regard to this latter application, most of the conjugated materials used in organic electronics are synthesized using time-consuming Suzuki-, Wittig-, or Heck-type coupling reactions that require expensive catalysts, stringent reaction conditions, and tedious purification processes. In order to afford a more economic route towards organic photovoltaic materials, Schiff bases derived from 2,5-thiophenedicarbaldehyde as the conjugated linker unit have recently been used. The azomethine bond, which is isoelectronic with the vinyl bond and possesses similar optoelectronic and thermal properties, is easily accessible through the Schiff condensation under near ambient reaction conditions (Morgan *et al.*, 1987; Pérez Guarín *et al.*, 2007; Sicard *et al.*, 2013).






We report here the synthesis and X-ray characterization of such thiophene derivatives, as a continuation of a partially published record (Bernès *et al.*, 2013; Mendoza *et al.*, 2014). We are improving a general solvent-free approach for these syntheses, recognising that ecological aspects in organic chemistry have become a priority, in order to minimize the quantity of toxic waste and by-products, and to decrease the amount of solvent in the reaction media or during work-up (Tanaka & Toda, 2000; Noyori, 2005).


In the synthesis of the thiophenes reported here, the Schiff condensation generates a single by-product, water, and a one-step recrystallization affords the pure substituted thiophene in nearly quantitative yields. Our protocol may be readily extended to any low molecular weight 2,5-substituted thiophene, providing that a liquid amine is used for the condensation. In the present work, the starting material is 2,5-thiophenedicarbaldehyde, a low melting-point compound (m.p. = 388–390 K), and four chiral amines were used. We took advantage of the anomalous dispersion of the sulfur sites to confirm that the configuration of the chiral amine is retained during the condensation.

## 2. Structural commentary

The first compound was synthesized using (*S*)-(+)-1-amino-tetraline. The Schiff base (I),  $C_{26}H_{26}N_2S$ , crystallizes in the space group *P*1, with the expected absolute configuration




**Figure 1**  
The molecular structure of (I), with displacement ellipsoids for non-H atoms at the 30% probability level.



**Figure 2**  
The molecular structure of (II), with displacement ellipsoids for non-H atoms at the 30% probability level. Non-labeled atoms are generated by symmetry code  $(1 - x, y, 1 - z)$ .

(Fig. 1). The general shape of the molecule displays a pseudo-twofold axis, passing through the S atom and the midpoint of the thiophene C–C  $\sigma$ -bond. As a consequence, the independent benzene rings are placed above and below the thiophene ring, and are inclined to one another at a dihedral angle of  $73.76(15)^\circ$ . The central core containing the thiophene ring and the imine bonds is virtually planar, and the imine bonds are substituted by the tetralin ring systems, which present the same conformation. The aliphatic rings C9–C13/C18 and C19–C23/C28 each have a half-chair conformation.

Compound (II),  $C_{24}H_{26}N_2O_2S$ , was obtained using (*R*)-(+)-(4-methoxy)phenylethylamine as the chiral component in the Schiff condensation. The twofold molecular axis, which was a latent symmetry in the case of (I), is a true crystallographic symmetry in (II), and this compound crystallizes in the space group *C*2 (Fig. 2). The asymmetric unit thus contains half a



**Figure 3**  
The molecular structures of isomorphous compounds (III) and (IV), with displacement ellipsoids for non-H atoms at the 30% probability level. Notice the different configuration for chiral center C5 in (III) and (IV). Non-labeled atoms are generated by symmetry codes  $(1 - x, -y, z)$  and  $(1 - x, 2 - y, z)$  for (III) and (IV), respectively.

Table 1

Comparison of C—H...S hydrogen bonds (Å, °) in compounds (I)–(IV).

Compound	Contact	C—H	H...S	C...S	C—H...S
(I)	C4—H4A...S1 <sup>i</sup>	0.93	3.00	3.562 (5)	121
(I)	C5—H5A...S1 <sup>i</sup>	0.93	2.97	3.547 (5)	122
(II)	C4—H4A...S1 <sup>ii</sup>	0.93	2.99	3.572 (3)	122
(III)	C4—H4A...S1 <sup>iii</sup>	0.93	3.15	3.743 (3)	124
(IV)	C4—H4A...S1 <sup>iv</sup>	0.93	3.23	3.828 (4)	124

 Symmetry codes: (i)  $x + 1, y, z$ ; (ii)  $x, y + 1, z$ ; (iii)  $x, y, z + 1$ ; (iv)  $x, y, z - 1$ .

molecule, and the molecular conformation for the complete molecule is similar to that of (I). The benzene rings have a free relative orientation, since these rings are not fused in a bicyclic system, as in (I); the dihedral angle between symmetry-related rings is 61.30 (7)°.

Compounds (III) and (IV), synthesized with enantiomerically pure (4-halogen)phenylethylamines (halogen = F, Cl) are isomorphous and crystallize with orthorhombic unit cells. The latent twofold symmetry of (I) is again observed, since both molecules lie on the crystallographic twofold axes of the space group  $P2_12_12$  (Fig. 3). The dihedral angle between the benzene rings is close to that observed for (II): 64.18 (8)° for (III) and 62.03 (9)° for (IV). The same Schiff base but with Br as the halogen substituent has been published previously (Mendoza *et al.*, 2014), but is not isomorphous with (III) and (IV). Instead, this molecule was found to crystallize in the space group  $C2$ , with unit-cell parameters and a crystal structure very similar to those of (II). A systematic trend is thus emerging for these 2,5-substituted thiophenes, related to the potential twofold molecular symmetry: they have a strong tendency to crystallize in space groups that include at least one  $C_2$  axis, such as  $C2$  and  $P2_12_12$  for the chiral crystals. This trend extends to achiral molecules, which also have twofold crystallographic symmetry in the space group  $C2/c$  (Kudiyakova *et al.*, 2011; Suganya *et al.*, 2014; Boyle *et al.*, 2015; Moussallem *et al.*, 2015). The features shared by these related compounds could also be a signature of a propensity towards polymorphism between monoclinic and orthorhombic systems.

The difference between non-crystallographic symmetry in (I) and exact  $C_2$  molecular symmetry in (II)–(IV) is also reflected in the degree of conjugation between thiophene rings and imine bonds. For (I), dihedral angles between the thiophene and C=N—C\* mean planes (C\* is the chiral C atom bonded to the imine functionality) are 6.9 (7) and 1.9 (6)°. Other crystals have a symmetry restriction, inducing a small deconjugation of the imine bonds. The corresponding dihedral angles with the thiophene rings are 8.5 (4), 10.1 (3), and 9.8 (3)°, for (II), (III) and (IV), respectively.

### 3. Supramolecular features

Although all compounds have benzene rings, neither  $\pi$ – $\pi$  nor C—H... $\pi$  contacts stabilize the crystal structures. However, these compounds share a common supramolecular feature. Lone pairs of S atoms interact with thiophenic CH groups of a




Figure 4

Part of the crystal structure of (II), showing C—H...S hydrogen bonds (dashed lines) linking molecules along [010]. [Symmetry codes: (i)  $1 - x, y, 1 - z$ ; (ii)  $x, 1 + y, z$ .]

neighboring molecule in the crystal, forming chains along the short cell axes: [100] for (I), [010] for (II) and [001] for (III) and (IV). An example is presented in Fig. 4, for compound (II). These bifurcated S...C—H contacts have a significant strength for (I), perhaps as a consequence of the relaxed molecular symmetry in space group  $P1$ . The contacts are weaker for (II), (III) and (IV), which have a geometry restrained by the crystallographic symmetry (Table 1).

### 4. Database survey

Many thiophenes substituted in the 2 and 5 positions by imine groups have been characterized; however, almost all were achiral compounds. X-ray structures have been reported mostly in space group  $C2/c$  (Suganya *et al.*, 2014; Kudiyakova *et al.*, 2011, 2012; Bolduc *et al.*, 2013b). Other represented space groups for achiral molecules are  $P2_1$  (Skene & Dufresne, 2006) and  $P2_1/c$  (Wiedermann *et al.*, 2005). Finally, a single case of a molecule presenting mirror symmetry has been described (Fridman & Kaftory, 2007), in space group  $Pnma$ .

The group of chiral molecules belonging to this family is much less populated, with two examples reported by our group in this journal. Both are molecules with the  $C_2$  point group and crystallize in space groups  $C2$  (Mendoza *et al.*, 2014) and  $P2_12_12$  (Bernès *et al.*, 2013).

### 5. Synthesis and crystallization

**Synthesis.** The chiral amines used for the Schiff condensation were obtained directly from suppliers: (*S*)-(+)-1,2,3,4-tetrahydro-1-naphthylamine for (I), (*R*)-(+)-1-(4-methoxyphenyl)ethylamine for (II), (*R*)-(+)-1-(4-fluorophenyl)ethylamine for (III) and (*S*)-(–)-1-(4-chlorophenyl)ethylamine for (IV). 2,5-Thiophenedicarbaldehyde (100 mg, 0.71 mmol) and the chiral amine (1.4 mmol) in a 1:2 molar ratio were mixed at room temperature under solvent-free conditions, giving light-yellow (II and IV), colorless (III) or light-brown (IV) solids, in

**Table 2**  
Experimental details.

	(I)	(II)	(III)	(IV)
<b>Crystal data</b>				
Chemical formula	C <sub>26</sub> H <sub>26</sub> N <sub>2</sub> S	C <sub>24</sub> H <sub>26</sub> N <sub>2</sub> O <sub>2</sub> S	C <sub>22</sub> H <sub>20</sub> F <sub>2</sub> N <sub>2</sub> S	C <sub>22</sub> H <sub>20</sub> Cl <sub>2</sub> N <sub>2</sub> S
<i>M<sub>r</sub></i>	398.55	406.53	382.46	415.36
Crystal system, space group	Triclinic, <i>P</i> 1	Monoclinic, <i>C</i> 2	Orthorhombic, <i>P</i> 2 <sub>1</sub> 2 <sub>1</sub> 2	Orthorhombic, <i>P</i> 2 <sub>1</sub> 2 <sub>1</sub> 2
Temperature (K)	298	298	298	298
<i>a</i> , <i>b</i> , <i>c</i> (Å)	5.9093 (4), 7.6258 (5), 12.6570 (8)	25.3917 (13), 5.9488 (3), 7.5623 (4)	21.1153 (16), 7.7846 (6), 6.1343 (5)	21.893 (2), 7.9212 (6), 6.2315 (4)
$\alpha$ , $\beta$ , $\gamma$ (°)	87.802 (5), 78.329 (5), 87.427 (5)	90, 97.174 (4), 90	90, 90, 90	90, 90, 90
<i>V</i> (Å <sup>3</sup> )	557.76 (6)	1133.34 (10)	1008.32 (14)	1080.66 (15)
<i>Z</i>	1	2	2	2
Radiation type	Mo <i>K</i> $\alpha$	Mo <i>K</i> $\alpha$	Mo <i>K</i> $\alpha$	Mo <i>K</i> $\alpha$
$\mu$ (mm <sup>-1</sup> )	0.16	0.16	0.19	0.41
Crystal size (mm)	0.34 × 0.12 × 0.06	0.45 × 0.33 × 0.12	0.89 × 0.47 × 0.33	0.52 × 0.40 × 0.07
<b>Data collection</b>				
Diffractometer	Agilent Xcalibur (Atlas, Gemini)	Agilent Xcalibur (Atlas, Gemini)	Agilent Xcalibur (Atlas, Gemini)	Agilent Xcalibur (Atlas, Gemini)
Absorption correction	Analytical <i>CrysAlis PRO</i> , (Agilent, 2013)	Analytical ( <i>CrysAlis PRO</i> ; Agilent, 2013)	Analytical <i>CrysAlis PRO</i> , (Agilent, 2013)	Multi-scan <i>CrysAlis PRO</i> , (Agilent, 2013)
<i>T</i> <sub>min</sub> , <i>T</i> <sub>max</sub>	0.969, 0.992	0.973, 0.993	0.904, 0.958	0.692, 1.000
No. of measured, independent and observed [ <i>I</i> > 2 $\sigma$ ( <i>I</i> )] reflections	6689, 4036, 2958	6341, 2221, 1892	12336, 2067, 1591	14195, 2743, 1534
<i>R</i> <sub>int</sub> (sin $\theta/\lambda$ ) <sub>max</sub> (Å <sup>-1</sup> )	0.040 0.618	0.027 0.618	0.058 0.625	0.058 0.692
<b>Refinement</b>				
<i>R</i> [ <i>F</i> <sup>2</sup> > 2 $\sigma$ ( <i>F</i> <sup>2</sup> )], <i>wR</i> ( <i>F</i> <sup>2</sup> ), <i>S</i>	0.058, 0.127, 1.02	0.036, 0.085, 1.02	0.044, 0.092, 1.06	0.052, 0.117, 1.01
No. of reflections	4036	2221	2067	2743
No. of parameters	262	134	124	124
No. of restraints	3	1	0	0
H-atom treatment	H-atom parameters constrained	H-atom parameters constrained	H-atom parameters constrained	H-atom parameters constrained
$\Delta\rho_{\max}$ , $\Delta\rho_{\min}$ (e Å <sup>-3</sup> )	0.31, -0.19	0.11, -0.17	0.15, -0.25	0.13, -0.17
Absolute structure	Flack <i>x</i> determined using 962 quotients [( <i>I</i> <sup>+</sup> ) - ( <i>I</i> <sup>-</sup> )] / [( <i>I</i> <sup>+</sup> ) + ( <i>I</i> <sup>-</sup> )] (Parsons <i>et al.</i> , 2013)	Flack <i>x</i> determined using 708 quotients [( <i>I</i> <sup>+</sup> ) - ( <i>I</i> <sup>-</sup> )] / [( <i>I</i> <sup>+</sup> ) + ( <i>I</i> <sup>-</sup> )] (Parsons <i>et al.</i> , 2013)	Flack <i>x</i> determined using 518 quotients [( <i>I</i> <sup>+</sup> ) - ( <i>I</i> <sup>-</sup> )] / [( <i>I</i> <sup>+</sup> ) + ( <i>I</i> <sup>-</sup> )] (Parsons <i>et al.</i> , 2013)	Flack <i>x</i> determined using 465 quotients [( <i>I</i> <sup>+</sup> ) - ( <i>I</i> <sup>-</sup> )] / [( <i>I</i> <sup>+</sup> ) + ( <i>I</i> <sup>-</sup> )] (Parsons <i>et al.</i> , 2013)
Absolute structure parameter	-0.12 (7)	-0.02 (4)	0.07 (6)	0.10 (6)

Computer programs: *CrysAlis PRO* (Agilent, 2013), *SHELXS97* (Sheldrick, 2008), *SHELXT* (Sheldrick, 2015a), *SHELXL2014* (Sheldrick, 2015b) and *Mercury* (Macrae *et al.*, 2008).

95–97% yields. The crude solids were recrystallized from CH<sub>2</sub>Cl<sub>2</sub>, affording colorless crystals of (I)–(IV).

**Spectroscopy.** (I): m.p. 437–438 K. [ $\alpha$ ]<sub>D</sub><sup>20</sup> = +655.4 (*c* = 1, CHCl<sub>3</sub>). FTIR: 1616 cm<sup>-1</sup> (C=N). <sup>1</sup>H NMR (500 MHz, CHCl<sub>3</sub>/TMS):  $\delta$  = 1.76–1.86 (*m*, 2H; *H*-al), 1.96–2.06 (*m*, 6H; *H*-al), 2.74–2.90 (*m*, 4H; *H*-al), 4.51 (*t*, 2H; *H*-al), 6.98–7.02 (*m*, 2H; *H*-ar), 7.09–7.15 (*m*, 6H; *H*-ar), 7.28 (*s*, 2H; *H*-ar), 8.36 (*s*, 2H; HC=N). <sup>13</sup>C NMR:  $\delta$  = 19.7, 29.3, 31.1, 67.7 (*C*-al), 125.7, 126.9, 128.7, 129.1, 129.6, 136.8, 137.1, 145.1 (*C*-ar), 153.1 (HC=N). MS–EI: *m/z* = 398 (*M*<sup>+</sup>).

(II): m.p. 405–406 K. [ $\alpha$ ]<sub>D</sub><sup>20</sup> = -626.8 (*c* = 1, CHCl<sub>3</sub>). FTIR: 1631 cm<sup>-1</sup> (C=N). <sup>1</sup>H NMR (500 MHz, CHCl<sub>3</sub>/TMS):  $\delta$  = 1.53 (*d*, 6H; CHCH<sub>3</sub>), 3.78 (*s*, 6H; OCH<sub>3</sub>), 4.47 (*q*, 2H; CHCH<sub>3</sub>), 6.85–6.88 (*m*, 4H; *H*-ar), 7.19 (*s*, 2H; *H*-ar), 7.29–7.32 (*m*, 4H; *H*-ar), 8.33 (*s*, 2H; HC=N). <sup>13</sup>C NMR:  $\delta$  = 24.8 (CHCH<sub>3</sub>), 55.2 (OCH<sub>3</sub>), 68.1 (CHCH<sub>3</sub>), 113.7, 127.6, 129.6, 137.1, 145.2, 152.1 (*C*-ar), 158.5 (HC=N). MS–EI: *m/z* = 406 (*M*<sup>+</sup>).

(III): m.p. 420–421 K. [ $\alpha$ ]<sub>D</sub><sup>20</sup> = -542.5 (*c* = 1, CHCl<sub>3</sub>). FTIR: 1621 cm<sup>-1</sup> (C=N). <sup>1</sup>H NMR (500 MHz, CHCl<sub>3</sub>/TMS):  $\delta$  = 1.53 (*d*, 6H; CHCH<sub>3</sub>), 4.49 (*q*, 2H; CHCH<sub>3</sub>), 7.00–7.38 (*m*, 10H;

*H*-ar), 8.37 (*s*, 2H; HC=N). <sup>13</sup>C NMR:  $\delta$  = 25.2 (CHCH<sub>3</sub>), 68.7 (CHCH<sub>3</sub>), 115.2 (*d*, *J*<sub>F-C</sub> = 21.2 Hz; *C*-ar), 128.1 (*d*, *J*<sub>F-C</sub> = 8.7 Hz; *C*-ar), 130.1 (*C*-ar), 140.7 (*d*, *J*<sub>F-C</sub> = 2.5 Hz; *C*-ar), 145.1 (*C*-ar), 161.1 (*d*, *J*<sub>F-C</sub> = 242.5 Hz; *C*-ar), 152.5 (HC=N). MS–EI: *m/z* = 382 (*M*<sup>+</sup>).

(IV): m.p. 434–435 K. [ $\alpha$ ]<sub>D</sub><sup>20</sup> = +726.5 (*c* = 1, CHCl<sub>3</sub>). FTIR: 1623 cm<sup>-1</sup> (C=N). <sup>1</sup>H NMR (500 MHz, CHCl<sub>3</sub>/TMS):  $\delta$  = 1.53 (*d*, 6H; CHCH<sub>3</sub>), 4.48 (*q*, 2H; CHCH<sub>3</sub>), 7.23–7.35 (*m*, 10H; *H*-ar), 8.37 (*s*, 2H; HC=N). <sup>13</sup>C NMR:  $\delta$  = 25.2 (CHCH<sub>3</sub>), 68.7 (CHCH<sub>3</sub>), 128.0, 128.6, 130.2, 132.5, 143.5, 145.1 (*C*-ar), 152.7 (HC=N).

## 6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. No unusual issues appeared, and refinements were carried out on non-restricted models. All H atoms were placed in calculated positions, and refined as riding on their carrier C atoms, with C–H bond lengths fixed to 0.93 (aromatic CH), 0.96 (methyl CH<sub>3</sub>), 0.97 (methylene

CH<sub>2</sub>), or 0.98 Å (methine CH). Isotropic displacement parameters were calculated as  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$  for methyl H atoms and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  for other H atoms. For all compounds, the absolute configuration was based on the refinement of the Flack parameter (Parsons *et al.*, 2013), confirming that the configuration of the chiral amine used as the starting material was retained during the Schiff condensation.

### Acknowledgements

Support from VIEP–UAP is acknowledged.

### References

- Agilent (2013). *CrysAlis PRO*. Agilent Technologies Inc., Santa Clara, CA, USA.
- Bernès, S., Hernández-Téllez, G., Sharma, M., Portillo-Moreno, O. & Gutiérrez, R. (2013). *Acta Cryst.* **E69**, o1428.
- Bolduc, A., Al Ouahabi, A., Mallet, C. & Skene, W. G. (2013*a*). *J. Org. Chem.* **78**, 9258–9269.
- Bolduc, A., Dufresne, S. & Skene, W. G. (2013*b*). *Acta Cryst.* **C69**, 1196–1199.
- Boyle, R., Crundwell, G. & Glagovich, N. M. (2015). *Acta Cryst.* **E71**, o403.
- Cram, D. J. & Trueblood, K. N. (1981). *Top. Curr. Chem.* **98**, 43–106.
- Dean, F. M. (1982*a*). *Adv. Heterocycl. Chem.* **30**, 167–238.
- Dean, F. M. (1982*b*). *Adv. Heterocycl. Chem.* **31**, 237–344.
- Fridman, N. & Kaftory, M. (2007). *Pol. J. Chem.* **81**, 825–832.
- Kudyakova, Yu. S., Burgart, Ya. V. & Saloutin, V. I. (2011). *Chem. Heterocycl. Compd.* **47**, 558–563.
- Kudyakova, Y. S., Burgart, Y. V., Slepukhin, P. A. & Saloutin, V. I. (2012). *Mendeleev Commun.* **22**, 284–286.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Mendoza, A., Bernès, S., Hernández-Téllez, G., Portillo-Moreno, O. & Gutiérrez, R. (2014). *Acta Cryst.* **E70**, o345.
- Morgan, P. W., Kwolek, S. L. & Pletcher, T. C. (1987). *Macromolecules*, **20**, 729–739.
- Moussallem, C., Allain, M., Mallet, C., Gohier, F. & Frère, P. (2015). *J. Fluor. Chem.* **178**, 34–39.
- Nelson, S. M., Esho, F., Lavery, A. & Drew, M. G. B. (1983). *J. Am. Chem. Soc.* **105**, 5693–5695.
- Noyori, R. (2005). *Chem. Commun.* pp. 1807–1811.
- Parsons, S., Flack, H. D. & Wagner, T. (2013). *Acta Cryst.* **B69**, 249–259.
- Pérez Guarín, S. A., Bourgeaux, M., Dufresne, S. & Skene, W. G. (2007). *J. Org. Chem.* **72**, 2631–2643.
- Petrus, M. L., Bouwer, R. K. M., Lafont, U., Athanasopoulos, S., Greenham, N. C. & Dingemans, T. J. (2014). *J. Mater. Chem. A*, **2**, 9474–9477.
- Sargent, M. V. & Cresp, T. M. (1975). *Fortschritte Chem. Forschung*, **57**, 111–143.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sheldrick, G. M. (2015*a*). *Acta Cryst.* **A71**, 3–8.
- Sheldrick, G. M. (2015*b*). *Acta Cryst.* **C71**, 3–8.
- Sicard, L., Navarathne, D., Skalski, T. & Skene, W. G. (2013). *Adv. Funct. Mater.* **23**, 3549–3559.
- Skene, W. G. & Dufresne, S. (2006). *Acta Cryst.* **E62**, o1116–o1117.
- Suganya, S., Velmathi, S. & MubarakAli, D. (2014). *Dyes Pigments*, **104**, 116–122.
- Tanaka, K. & Toda, F. (2000). *Chem. Rev.* **100**, 1025–1074.
- Wiedermann, J., Kirchner, K. & Mereiter, K. (2005). Private communication (refcode NAWMAA). CCDC, Cambridge, England.



## supporting information

*Acta Cryst.* (2016). E72, 350-354 [doi:10.1107/S2056989016002516]

## Crystal structures of four chiral imine-substituted thiophene derivatives

Guadalupe Hernández-Téllez, Sylvain Bernès, Angel Mendoza, Francisco Javier Ríos-Merino, Gloria E. Moreno, Oscar Portillo and René Gutiérrez

## Computing details

For all compounds, data collection: *CrysAlis PRO* (Agilent, 2013); cell refinement: *CrysAlis PRO* (Agilent, 2013); data reduction: *CrysAlis PRO* (Agilent, 2013). Program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008) for (I); *SHELXT* (Sheldrick, 2015a) for (II), (III), (IV). For all compounds, program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015b); molecular graphics: *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXL2014* (Sheldrick, 2015b).

## (I) 2,5-Bis[(S)-(+)-(1,2,3,4-tetrahydro-1-naphthyl)imino]thiophene

## Crystal data

$C_{26}H_{26}N_2S$	$F(000) = 212$
$M_r = 398.55$	$D_x = 1.187 \text{ Mg m}^{-3}$
Triclinic, $P1$	Melting point: 437 K
$a = 5.9093 (4) \text{ \AA}$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$b = 7.6258 (5) \text{ \AA}$	Cell parameters from 2148 reflections
$c = 12.6570 (8) \text{ \AA}$	$\theta = 3.3\text{--}22.6^\circ$
$\alpha = 87.802 (5)^\circ$	$\mu = 0.16 \text{ mm}^{-1}$
$\beta = 78.329 (5)^\circ$	$T = 298 \text{ K}$
$\gamma = 87.427 (5)^\circ$	Plate, colorless
$V = 557.76 (6) \text{ \AA}^3$	$0.34 \times 0.12 \times 0.06 \text{ mm}$
$Z = 1$	

## Data collection

Agilent Xcalibur (Atlas, Gemini) diffractometer	6689 measured reflections
Radiation source: Enhance (Mo) X-ray Source	4036 independent reflections
Graphite monochromator	2958 reflections with $I > 2\sigma(I)$
Detector resolution: $10.5564 \text{ pixels mm}^{-1}$	$R_{\text{int}} = 0.040$
$\omega$ scans	$\theta_{\text{max}} = 26.1^\circ$ , $\theta_{\text{min}} = 3.1^\circ$
Absorption correction: analytical <i>CrysAlis PRO</i> , (Agilent, 2013)	$h = -7 \rightarrow 7$
$T_{\text{min}} = 0.969$ , $T_{\text{max}} = 0.992$	$k = -9 \rightarrow 9$
	$l = -15 \rightarrow 15$

## Refinement

Refinement on $F^2$	262 parameters
Least-squares matrix: full	3 restraints
$R[F^2 > 2\sigma(F^2)] = 0.058$	0 constraints
$wR(F^2) = 0.127$	Primary atom site location: structure-invariant
$S = 1.02$	direct methods
4036 reflections	

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0525P)^2]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.31 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.19 \text{ e } \text{\AA}^{-3}$$

Absolute structure: Flack  $x$  determined using

962 quotients  $[(I^-)-(I^+)]/[(I^+)+(I^-)]$  (Parsons *et al.*, 2013)

Absolute structure parameter:  $-0.12$  (7)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.66581 (19)	0.49640 (17)	0.11819 (12)	0.0488 (4)
N1	0.5097 (7)	0.7980 (6)	0.2625 (3)	0.0507 (12)
C2	0.7239 (10)	0.7657 (7)	0.2474 (4)	0.0490 (13)
H2A	0.8132	0.8300	0.2834	0.059*
C3	0.8355 (8)	0.6297 (7)	0.1747 (4)	0.0469 (13)
C4	1.0622 (9)	0.5892 (7)	0.1402 (5)	0.0575 (15)
H4A	1.1792	0.6461	0.1628	0.069*
C5	1.1042 (8)	0.4510 (7)	0.0658 (5)	0.0595 (15)
H5A	1.2513	0.4075	0.0348	0.071*
C6	0.9068 (8)	0.3894 (6)	0.0450 (4)	0.0425 (12)
C7	0.8786 (9)	0.2528 (7)	-0.0268 (4)	0.0503 (14)
H7A	1.0094	0.1943	-0.0651	0.060*
N8	0.6816 (8)	0.2106 (6)	-0.0390 (3)	0.0518 (12)
C9	0.4190 (9)	0.9453 (7)	0.3325 (4)	0.0518 (13)
H9A	0.5365	0.9745	0.3730	0.062*
C10	0.3728 (12)	1.1032 (8)	0.2631 (5)	0.0772 (18)
H10A	0.2751	1.0713	0.2143	0.093*
H10B	0.5174	1.1431	0.2201	0.093*
C11	0.2537 (13)	1.2501 (8)	0.3345 (5)	0.0802 (19)
H11A	0.3449	1.2749	0.3875	0.096*
H11B	0.2407	1.3560	0.2909	0.096*
C12	0.0161 (11)	1.1958 (8)	0.3911 (5)	0.0682 (18)
H12A	-0.0468	1.2796	0.4462	0.082*
H12B	-0.0847	1.1991	0.3393	0.082*
C13	0.0174 (9)	1.0143 (7)	0.4429 (4)	0.0486 (14)
C14	-0.1721 (10)	0.9610 (9)	0.5196 (5)	0.0620 (16)
H14A	-0.2950	1.0410	0.5406	0.074*
C15	-0.1846 (11)	0.7955 (9)	0.5651 (5)	0.0749 (18)
H15A	-0.3143	0.7635	0.6159	0.090*
C16	-0.0009 (13)	0.6756 (9)	0.5347 (6)	0.080 (2)
H16A	-0.0068	0.5621	0.5644	0.095*
C17	0.1892 (11)	0.7268 (8)	0.4602 (5)	0.0665 (16)
H17A	0.3134	0.6471	0.4414	0.080*
C18	0.2020 (8)	0.8935 (7)	0.4123 (4)	0.0465 (12)
C19	0.6721 (9)	0.0655 (6)	-0.1121 (4)	0.0498 (13)
H19A	0.8294	0.0400	-0.1523	0.060*
C20	0.5911 (13)	-0.0955 (8)	-0.0465 (5)	0.0728 (17)
H20A	0.4515	-0.0668	0.0058	0.087*

H20B	0.7086	-0.1390	-0.0075	0.087*
C21	0.5425 (13)	-0.2380 (8)	-0.1206 (5)	0.0755 (19)
H21A	0.6802	-0.2628	-0.1750	0.091*
H21B	0.5024	-0.3453	-0.0786	0.091*
C22	0.3465 (11)	-0.1769 (9)	-0.1746 (5)	0.0688 (18)
H22A	0.3350	-0.2584	-0.2300	0.083*
H22B	0.2028	-0.1782	-0.1216	0.083*
C23	0.3768 (9)	0.0051 (8)	-0.2248 (4)	0.0503 (14)
C24	0.2515 (10)	0.0601 (9)	-0.3022 (4)	0.0624 (15)
H24A	0.1532	-0.0175	-0.3233	0.075*
C25	0.2684 (12)	0.2252 (10)	-0.3484 (5)	0.079 (2)
H25A	0.1830	0.2591	-0.4004	0.095*
C26	0.4143 (14)	0.3418 (9)	-0.3167 (6)	0.086 (2)
H26A	0.4269	0.4550	-0.3469	0.103*
C27	0.5398 (11)	0.2877 (8)	-0.2403 (5)	0.0671 (17)
H27A	0.6391	0.3653	-0.2199	0.080*
C28	0.5226 (8)	0.1213 (7)	-0.1928 (4)	0.0484 (13)

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0351 (7)	0.0574 (8)	0.0551 (7)	-0.0009 (5)	-0.0088 (5)	-0.0201 (6)
N1	0.045 (3)	0.058 (3)	0.049 (3)	0.004 (2)	-0.006 (2)	-0.026 (2)
C2	0.049 (3)	0.055 (3)	0.047 (3)	-0.007 (3)	-0.013 (2)	-0.016 (3)
C3	0.039 (3)	0.058 (3)	0.048 (3)	-0.002 (2)	-0.016 (2)	-0.014 (3)
C4	0.036 (3)	0.071 (4)	0.070 (4)	0.000 (3)	-0.017 (3)	-0.027 (3)
C5	0.033 (3)	0.075 (4)	0.072 (4)	0.005 (3)	-0.010 (2)	-0.031 (3)
C6	0.031 (3)	0.050 (3)	0.046 (3)	0.004 (2)	-0.006 (2)	-0.011 (2)
C7	0.046 (3)	0.057 (3)	0.048 (3)	0.010 (3)	-0.007 (2)	-0.018 (3)
N8	0.048 (3)	0.059 (3)	0.050 (3)	-0.003 (2)	-0.008 (2)	-0.023 (2)
C9	0.049 (3)	0.053 (3)	0.054 (3)	-0.001 (3)	-0.009 (3)	-0.022 (3)
C10	0.102 (5)	0.061 (4)	0.060 (4)	-0.002 (3)	0.005 (3)	-0.015 (3)
C11	0.109 (5)	0.050 (4)	0.070 (4)	0.005 (3)	0.010 (4)	-0.007 (3)
C12	0.082 (5)	0.062 (4)	0.060 (4)	0.024 (3)	-0.015 (3)	-0.022 (3)
C13	0.048 (3)	0.057 (3)	0.043 (3)	0.002 (3)	-0.012 (3)	-0.016 (3)
C14	0.056 (3)	0.075 (4)	0.056 (3)	0.009 (3)	-0.009 (3)	-0.027 (3)
C15	0.075 (4)	0.082 (5)	0.062 (4)	-0.011 (4)	0.006 (3)	-0.022 (4)
C16	0.104 (5)	0.061 (4)	0.066 (4)	-0.012 (4)	0.001 (4)	-0.002 (4)
C17	0.072 (4)	0.059 (4)	0.064 (4)	0.007 (3)	-0.003 (3)	-0.011 (3)
C18	0.045 (3)	0.054 (3)	0.043 (3)	0.001 (3)	-0.012 (2)	-0.019 (3)
C19	0.052 (3)	0.053 (3)	0.044 (3)	0.000 (3)	-0.007 (2)	-0.017 (3)
C20	0.114 (5)	0.060 (4)	0.051 (3)	-0.010 (4)	-0.030 (3)	-0.011 (3)
C21	0.119 (5)	0.055 (4)	0.056 (4)	-0.012 (4)	-0.021 (4)	-0.009 (3)
C22	0.074 (4)	0.078 (5)	0.056 (4)	-0.028 (4)	-0.008 (3)	-0.017 (3)
C23	0.048 (3)	0.057 (3)	0.043 (3)	-0.007 (3)	0.001 (3)	-0.019 (3)
C24	0.056 (3)	0.079 (4)	0.054 (3)	0.003 (3)	-0.013 (3)	-0.028 (3)
C25	0.096 (5)	0.084 (5)	0.066 (4)	0.026 (4)	-0.037 (4)	-0.028 (4)
C26	0.129 (6)	0.057 (4)	0.080 (5)	0.016 (4)	-0.044 (5)	-0.010 (4)



C27	0.084 (4)	0.059 (4)	0.064 (4)	-0.003 (3)	-0.022 (4)	-0.019 (3)
C28	0.050 (3)	0.046 (3)	0.049 (3)	0.006 (2)	-0.006 (2)	-0.018 (3)

*Geometric parameters (Å, °)*

S1—C6	1.724 (5)	C14—H14A	0.9300
S1—C3	1.728 (5)	C15—C16	1.390 (9)
N1—C2	1.255 (6)	C15—H15A	0.9300
N1—C9	1.471 (6)	C16—C17	1.373 (9)
C2—C3	1.458 (7)	C16—H16A	0.9300
C2—H2A	0.9300	C17—C18	1.386 (8)
C3—C4	1.348 (7)	C17—H17A	0.9300
C4—C5	1.420 (7)	C19—C20	1.496 (7)
C4—H4A	0.9300	C19—C28	1.518 (7)
C5—C6	1.355 (6)	C19—H19A	0.9800
C5—H5A	0.9300	C20—C21	1.536 (8)
C6—C7	1.445 (7)	C20—H20A	0.9700
C7—N8	1.263 (6)	C20—H20B	0.9700
C7—H7A	0.9300	C21—C22	1.508 (9)
N8—C19	1.479 (6)	C21—H21A	0.9700
C9—C10	1.512 (8)	C21—H21B	0.9700
C9—C18	1.520 (7)	C22—C23	1.507 (9)
C9—H9A	0.9800	C22—H22A	0.9700
C10—C11	1.522 (8)	C22—H22B	0.9700
C10—H10A	0.9700	C23—C24	1.384 (8)
C10—H10B	0.9700	C23—C28	1.389 (7)
C11—C12	1.510 (9)	C24—C25	1.367 (9)
C11—H11A	0.9700	C24—H24A	0.9300
C11—H11B	0.9700	C25—C26	1.390 (10)
C12—C13	1.509 (8)	C25—H25A	0.9300
C12—H12A	0.9700	C26—C27	1.374 (8)
C12—H12B	0.9700	C26—H26A	0.9300
C13—C14	1.390 (8)	C27—C28	1.382 (7)
C13—C18	1.398 (7)	C27—H27A	0.9300
C14—C15	1.366 (9)		
C6—S1—C3	91.5 (2)	C16—C15—H15A	120.4
C2—N1—C9	116.5 (4)	C17—C16—C15	119.2 (7)
N1—C2—C3	121.5 (5)	C17—C16—H16A	120.4
N1—C2—H2A	119.3	C15—C16—H16A	120.4
C3—C2—H2A	119.3	C16—C17—C18	122.1 (6)
C4—C3—C2	129.7 (5)	C16—C17—H17A	119.0
C4—C3—S1	111.3 (4)	C18—C17—H17A	119.0
C2—C3—S1	119.1 (4)	C17—C18—C13	118.7 (5)
C3—C4—C5	113.2 (5)	C17—C18—C9	120.0 (5)
C3—C4—H4A	123.4	C13—C18—C9	121.2 (5)
C5—C4—H4A	123.4	N8—C19—C20	109.4 (4)
C6—C5—C4	112.6 (5)	N8—C19—C28	110.1 (4)

C6—C5—H5A	123.7	C20—C19—C28	113.3 (4)
C4—C5—H5A	123.7	N8—C19—H19A	108.0
C5—C6—C7	129.0 (5)	C20—C19—H19A	108.0
C5—C6—S1	111.4 (4)	C28—C19—H19A	108.0
C7—C6—S1	119.6 (4)	C19—C20—C21	109.9 (4)
N8—C7—C6	121.9 (5)	C19—C20—H20A	109.7
N8—C7—H7A	119.1	C21—C20—H20A	109.7
C6—C7—H7A	119.1	C19—C20—H20B	109.7
C7—N8—C19	117.5 (4)	C21—C20—H20B	109.7
N1—C9—C10	109.1 (4)	H20A—C20—H20B	108.2
N1—C9—C18	110.3 (4)	C22—C21—C20	109.9 (5)
C10—C9—C18	111.6 (5)	C22—C21—H21A	109.7
N1—C9—H9A	108.6	C20—C21—H21A	109.7
C10—C9—H9A	108.6	C22—C21—H21B	109.7
C18—C9—H9A	108.6	C20—C21—H21B	109.7
C9—C10—C11	109.6 (5)	H21A—C21—H21B	108.2
C9—C10—H10A	109.7	C23—C22—C21	112.9 (5)
C11—C10—H10A	109.7	C23—C22—H22A	109.0
C9—C10—H10B	109.7	C21—C22—H22A	109.0
C11—C10—H10B	109.7	C23—C22—H22B	109.0
H10A—C10—H10B	108.2	C21—C22—H22B	109.0
C12—C11—C10	109.6 (5)	H22A—C22—H22B	107.8
C12—C11—H11A	109.7	C24—C23—C28	119.1 (5)
C10—C11—H11A	109.7	C24—C23—C22	119.5 (5)
C12—C11—H11B	109.7	C28—C23—C22	121.4 (5)
C10—C11—H11B	109.7	C25—C24—C23	121.8 (6)
H11A—C11—H11B	108.2	C25—C24—H24A	119.1
C13—C12—C11	112.9 (5)	C23—C24—H24A	119.1
C13—C12—H12A	109.0	C24—C25—C26	119.3 (6)
C11—C12—H12A	109.0	C24—C25—H25A	120.4
C13—C12—H12B	109.0	C26—C25—H25A	120.4
C11—C12—H12B	109.0	C27—C26—C25	119.1 (6)
H12A—C12—H12B	107.8	C27—C26—H26A	120.4
C14—C13—C18	118.4 (5)	C25—C26—H26A	120.4
C14—C13—C12	120.1 (5)	C26—C27—C28	121.9 (6)
C18—C13—C12	121.5 (5)	C26—C27—H27A	119.1
C15—C14—C13	122.5 (6)	C28—C27—H27A	119.1
C15—C14—H14A	118.8	C27—C28—C23	118.8 (5)
C13—C14—H14A	118.8	C27—C28—C19	119.8 (5)
C14—C15—C16	119.1 (6)	C23—C28—C19	121.3 (5)
C14—C15—H15A	120.4		
C9—N1—C2—C3	-176.4 (5)	C12—C13—C18—C17	-177.8 (5)
N1—C2—C3—C4	172.4 (6)	C14—C13—C18—C9	-177.3 (5)
N1—C2—C3—S1	-6.2 (7)	C12—C13—C18—C9	5.4 (7)
C6—S1—C3—C4	-1.4 (5)	N1—C9—C18—C17	39.8 (6)
C6—S1—C3—C2	177.5 (4)	C10—C9—C18—C17	161.1 (5)
C2—C3—C4—C5	-177.8 (5)	N1—C9—C18—C13	-143.5 (4)

S1—C3—C4—C5	0.9 (6)	C10—C9—C18—C13	-22.1 (6)
C3—C4—C5—C6	0.2 (7)	C7—N8—C19—C20	105.5 (6)
C4—C5—C6—C7	178.9 (5)	C7—N8—C19—C28	-129.5 (5)
C4—C5—C6—S1	-1.3 (6)	N8—C19—C20—C21	170.8 (5)
C3—S1—C6—C5	1.5 (4)	C28—C19—C20—C21	47.7 (7)
C3—S1—C6—C7	-178.7 (4)	C19—C20—C21—C22	-64.0 (7)
C5—C6—C7—N8	-179.1 (6)	C20—C21—C22—C23	49.0 (7)
S1—C6—C7—N8	1.1 (7)	C21—C22—C23—C24	161.6 (5)
C6—C7—N8—C19	-178.2 (5)	C21—C22—C23—C28	-20.5 (8)
C2—N1—C9—C10	102.3 (6)	C28—C23—C24—C25	0.4 (8)
C2—N1—C9—C18	-134.9 (5)	C22—C23—C24—C25	178.4 (6)
N1—C9—C10—C11	173.7 (5)	C23—C24—C25—C26	-0.3 (9)
C18—C9—C10—C11	51.6 (7)	C24—C25—C26—C27	0.5 (10)
C9—C10—C11—C12	-66.2 (7)	C25—C26—C27—C28	-0.9 (10)
C10—C11—C12—C13	48.1 (7)	C26—C27—C28—C23	1.0 (8)
C11—C12—C13—C14	164.1 (5)	C26—C27—C28—C19	177.5 (6)
C11—C12—C13—C18	-18.7 (8)	C24—C23—C28—C27	-0.7 (7)
C18—C13—C14—C15	-0.5 (8)	C22—C23—C28—C27	-178.7 (6)
C12—C13—C14—C15	176.8 (5)	C24—C23—C28—C19	-177.2 (5)
C13—C14—C15—C16	0.5 (9)	C22—C23—C28—C19	4.9 (8)
C14—C15—C16—C17	0.5 (10)	N8—C19—C28—C27	41.7 (6)
C15—C16—C17—C18	-1.6 (10)	C20—C19—C28—C27	164.5 (5)
C16—C17—C18—C13	1.6 (9)	N8—C19—C28—C23	-141.9 (5)
C16—C17—C18—C9	178.4 (6)	C20—C19—C28—C23	-19.1 (7)
C14—C13—C18—C17	-0.5 (7)		

## Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C4—H4 <i>A</i> ...S1 <sup>i</sup>	0.93	3.00	3.562 (5)	121
C5—H5 <i>A</i> ...S1 <sup>i</sup>	0.93	2.97	3.547 (5)	122

Symmetry code: (i) *x*+1, *y*, *z*.(II) 2,5-Bis{[(*R*)-(-)-1-(4-methoxyphenyl)ethyl]iminomethyl}thiophene

## Crystal data

C<sub>24</sub>H<sub>26</sub>N<sub>2</sub>O<sub>2</sub>S*M<sub>r</sub>* = 406.53Monoclinic, *C*2*a* = 25.3917 (13) Å*b* = 5.9488 (3) Å*c* = 7.5623 (4) Å $\beta$  = 97.174 (4)°*V* = 1133.34 (10) Å<sup>3</sup>*Z* = 2*F*(000) = 432*D<sub>x</sub>* = 1.191 Mg m<sup>-3</sup>

Melting point: 405 K

Mo *K*α radiation,  $\lambda$  = 0.71073 Å

Cell parameters from 2504 reflections

 $\theta$  = 3.0–24.2° $\mu$  = 0.16 mm<sup>-1</sup>*T* = 298 K

Prism, colourless

0.45 × 0.33 × 0.12 mm

Data collection

Agilent Xcalibur (Atlas, Gemini)  
diffractometer

Radiation source: Enhance (Mo) X-ray Source

Graphite monochromator

$\omega$  scans

Absorption correction: analytical  
(*CrysAlis PRO*; Agilent, 2013)

$T_{\min} = 0.973$ ,  $T_{\max} = 0.993$

6341 measured reflections

2221 independent reflections

1892 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.027$

$\theta_{\max} = 26.1^\circ$ ,  $\theta_{\min} = 3.0^\circ$

$h = -31 \rightarrow 31$

$k = -7 \rightarrow 7$

$l = -9 \rightarrow 9$

Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.036$

$wR(F^2) = 0.085$

$S = 1.02$

2221 reflections

134 parameters

1 restraint

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0393P)^2 + 0.1801P]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.11 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.17 \text{ e } \text{\AA}^{-3}$

Absolute structure: Flack  $x$  determined using

708 quotients  $[(F^-)-(F^+)]/[(F^-)+(F^+)]$  (Parsons *et al.*,  
2013)

Absolute structure parameter:  $-0.02$  (4)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.5000	0.37429 (14)	0.5000	0.0490 (3)
N1	0.56565 (9)	0.3213 (4)	0.1855 (3)	0.0471 (6)
C2	0.55195 (11)	0.5176 (5)	0.2189 (4)	0.0469 (7)
H2A	0.5608	0.6324	0.1445	0.056*
C3	0.52314 (10)	0.5774 (5)	0.3665 (4)	0.0449 (6)
C4	0.51313 (13)	0.7856 (5)	0.4229 (4)	0.0596 (8)
H4A	0.5225	0.9159	0.3665	0.072*
C5	0.59848 (11)	0.2933 (4)	0.0386 (4)	0.0484 (7)
H5A	0.5949	0.4291	-0.0354	0.058*
C6	0.57863 (13)	0.0963 (6)	-0.0754 (4)	0.0658 (8)
H6A	0.5416	0.1162	-0.1164	0.099*
H6B	0.5835	-0.0394	-0.0065	0.099*
H6C	0.5981	0.0861	-0.1759	0.099*
C7	0.65613 (11)	0.2719 (4)	0.1189 (3)	0.0441 (6)
C8	0.69277 (11)	0.4349 (4)	0.0871 (4)	0.0494 (7)
H8A	0.6817	0.5585	0.0168	0.059*
C9	0.74515 (11)	0.4171 (5)	0.1576 (4)	0.0571 (8)
H9A	0.7692	0.5278	0.1343	0.069*
C10	0.76215 (11)	0.2354 (6)	0.2628 (4)	0.0546 (7)
C11	0.72642 (12)	0.0745 (6)	0.2994 (4)	0.0593 (8)
H11A	0.7375	-0.0469	0.3722	0.071*
C12	0.67374 (12)	0.0936 (6)	0.2275 (4)	0.0551 (8)
H12A	0.6497	-0.0161	0.2528	0.066*
O1	0.81556 (9)	0.2307 (5)	0.3227 (3)	0.0781 (7)
C13	0.83539 (15)	0.0345 (9)	0.4189 (5)	0.1010 (15)

H13A	0.8734	0.0434	0.4432	0.152*
H13B	0.8257	-0.0970	0.3487	0.152*
H13C	0.8204	0.0258	0.5292	0.152*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0540 (6)	0.0354 (5)	0.0603 (6)	0.000	0.0183 (4)	0.000
N1	0.0442 (13)	0.0512 (16)	0.0477 (13)	0.0014 (10)	0.0129 (10)	0.0058 (10)
C2	0.0460 (15)	0.0443 (18)	0.0507 (16)	-0.0021 (13)	0.0071 (13)	0.0121 (13)
C3	0.0421 (14)	0.0392 (14)	0.0534 (17)	-0.0002 (12)	0.0062 (13)	0.0052 (12)
C4	0.074 (2)	0.0365 (16)	0.071 (2)	0.0007 (13)	0.0193 (16)	0.0074 (13)
C5	0.0497 (16)	0.0537 (18)	0.0434 (15)	0.0015 (12)	0.0120 (12)	0.0097 (12)
C6	0.0626 (19)	0.079 (2)	0.0551 (19)	-0.0010 (18)	0.0058 (15)	-0.0036 (17)
C7	0.0468 (15)	0.0492 (15)	0.0387 (14)	0.0022 (13)	0.0147 (12)	0.0018 (12)
C8	0.0559 (17)	0.0500 (17)	0.0451 (14)	-0.0025 (13)	0.0169 (13)	0.0045 (12)
C9	0.0528 (17)	0.067 (2)	0.0548 (17)	-0.0139 (16)	0.0177 (14)	-0.0004 (16)
C10	0.0455 (16)	0.078 (2)	0.0414 (16)	0.0001 (15)	0.0080 (13)	-0.0080 (15)
C11	0.0575 (18)	0.068 (2)	0.0524 (18)	0.0087 (17)	0.0086 (14)	0.0150 (15)
C12	0.0527 (17)	0.0569 (17)	0.0574 (19)	-0.0031 (14)	0.0138 (14)	0.0147 (15)
O1	0.0478 (13)	0.121 (2)	0.0642 (14)	-0.0021 (14)	0.0011 (10)	-0.0011 (14)
C13	0.063 (2)	0.161 (4)	0.075 (3)	0.022 (3)	-0.0086 (19)	0.022 (3)

*Geometric parameters (Å, °)*

S1—C3 <sup>i</sup>	1.724 (3)	C7—C12	1.381 (4)
S1—C3	1.724 (3)	C7—C8	1.385 (3)
N1—C2	1.253 (3)	C8—C9	1.374 (4)
N1—C5	1.480 (3)	C8—H8A	0.9300
C2—C3	1.453 (4)	C9—C10	1.379 (4)
C2—H2A	0.9300	C9—H9A	0.9300
C3—C4	1.345 (4)	C10—C11	1.371 (4)
C4—C4 <sup>i</sup>	1.413 (6)	C10—O1	1.375 (3)
C4—H4A	0.9300	C11—C12	1.384 (4)
C5—C6	1.504 (4)	C11—H11A	0.9300
C5—C7	1.519 (4)	C12—H12A	0.9300
C5—H5A	0.9800	O1—C13	1.433 (5)
C6—H6A	0.9600	C13—H13A	0.9600
C6—H6B	0.9600	C13—H13B	0.9600
C6—H6C	0.9600	C13—H13C	0.9600
C3 <sup>i</sup> —S1—C3	91.01 (19)	C12—C7—C5	121.8 (2)
C2—N1—C5	116.9 (2)	C8—C7—C5	120.4 (2)
N1—C2—C3	124.2 (3)	C9—C8—C7	121.2 (3)
N1—C2—H2A	117.9	C9—C8—H8A	119.4
C3—C2—H2A	117.9	C7—C8—H8A	119.4
C4—C3—C2	127.1 (3)	C8—C9—C10	120.1 (3)
C4—C3—S1	111.6 (2)	C8—C9—H9A	119.9

C2—C3—S1	121.3 (2)	C10—C9—H9A	119.9
C3—C4—C4 <sup>i</sup>	112.91 (17)	C11—C10—O1	124.7 (3)
C3—C4—H4A	123.5	C11—C10—C9	119.8 (3)
C4 <sup>i</sup> —C4—H4A	123.5	O1—C10—C9	115.5 (3)
N1—C5—C6	109.7 (2)	C10—C11—C12	119.7 (3)
N1—C5—C7	108.3 (2)	C10—C11—H11A	120.2
C6—C5—C7	113.7 (2)	C12—C11—H11A	120.2
N1—C5—H5A	108.3	C7—C12—C11	121.5 (3)
C6—C5—H5A	108.3	C7—C12—H12A	119.3
C7—C5—H5A	108.3	C11—C12—H12A	119.3
C5—C6—H6A	109.5	C10—O1—C13	117.0 (3)
C5—C6—H6B	109.5	O1—C13—H13A	109.5
H6A—C6—H6B	109.5	O1—C13—H13B	109.5
C5—C6—H6C	109.5	H13A—C13—H13B	109.5
H6A—C6—H6C	109.5	O1—C13—H13C	109.5
H6B—C6—H6C	109.5	H13A—C13—H13C	109.5
C12—C7—C8	117.8 (3)	H13B—C13—H13C	109.5
C5—N1—C2—C3	-175.3 (2)	C12—C7—C8—C9	1.5 (4)
N1—C2—C3—C4	171.0 (3)	C5—C7—C8—C9	-179.4 (3)
N1—C2—C3—S1	-5.7 (4)	C7—C8—C9—C10	-0.3 (4)
C3 <sup>i</sup> —S1—C3—C4	-0.22 (17)	C8—C9—C10—C11	-1.1 (4)
C3 <sup>i</sup> —S1—C3—C2	176.9 (3)	C8—C9—C10—O1	178.3 (2)
C2—C3—C4—C4 <sup>i</sup>	-176.3 (3)	O1—C10—C11—C12	-178.1 (3)
S1—C3—C4—C4 <sup>i</sup>	0.6 (4)	C9—C10—C11—C12	1.2 (4)
C2—N1—C5—C6	-136.4 (3)	C8—C7—C12—C11	-1.4 (4)
C2—N1—C5—C7	99.0 (3)	C5—C7—C12—C11	179.6 (3)
N1—C5—C7—C12	63.9 (3)	C10—C11—C12—C7	0.0 (5)
C6—C5—C7—C12	-58.3 (3)	C11—C10—O1—C13	4.8 (4)
N1—C5—C7—C8	-115.0 (3)	C9—C10—O1—C13	-174.6 (3)
C6—C5—C7—C8	122.7 (3)		

Symmetry code: (i)  $-x+1, y, -z+1$ .

#### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C4—H4A $\cdots$ S1 <sup>ii</sup>	0.93	2.99	3.572 (3)	122

Symmetry code: (ii)  $x, y+1, z$ .

#### (III) 2,5-Bis{[(*R*)-(-)-1-(4-fluorophenyl)ethyl]iminomethyl}thiophene

##### Crystal data

$C_{22}H_{20}F_2N_2S$

$M_r = 382.46$

Orthorhombic,  $P2_12_12$

$a = 21.1153$  (16)  $\text{\AA}$

$b = 7.7846$  (6)  $\text{\AA}$

$c = 6.1343$  (5)  $\text{\AA}$

$V = 1008.32$  (14)  $\text{\AA}^3$

$Z = 2$

$F(000) = 400$

$D_x = 1.260$   $\text{Mg m}^{-3}$

Melting point: 420 K

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$   $\text{\AA}$

Cell parameters from 2744 reflections

$\theta = 3.8\text{--}23.2^\circ$



$\mu = 0.19 \text{ mm}^{-1}$   
 $T = 298 \text{ K}$

Prism, colourless  
 $0.89 \times 0.47 \times 0.33 \text{ mm}$

*Data collection*

Agilent Xcalibur (Atlas, Gemini)  
 diffractometer  
 Radiation source: Enhance (Mo) X-ray Source  
 Graphite monochromator  
 Detector resolution:  $10.5564 \text{ pixels mm}^{-1}$   
 $\omega$  scans  
 Absorption correction: analytical  
*CrysAlis PRO*, (Agilent, 2013)  
 $T_{\text{min}} = 0.904$ ,  $T_{\text{max}} = 0.958$

12336 measured reflections  
 2067 independent reflections  
 1591 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.058$   
 $\theta_{\text{max}} = 26.4^\circ$ ,  $\theta_{\text{min}} = 3.8^\circ$   
 $h = -26 \rightarrow 26$   
 $k = -9 \rightarrow 9$   
 $l = -7 \rightarrow 7$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.044$   
 $wR(F^2) = 0.092$   
 $S = 1.06$   
 2067 reflections  
 124 parameters  
 0 restraints  
 Primary atom site location: structure-invariant  
 direct methods  
 Secondary atom site location: difference Fourier  
 map

Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0384P)^2 + 0.0613P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\text{max}} < 0.001$   
 $\Delta\rho_{\text{max}} = 0.15 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.25 \text{ e } \text{\AA}^{-3}$   
 Absolute structure: Flack  $x$  determined using  
 518 quotients  $[(I^-)-(I)]/[(I^+)+(I)]$  (Parsons *et al.*,  
 2013)  
 Absolute structure parameter: 0.07 (6)

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.5000	0.0000	1.06817 (15)	0.0479 (3)
F1	0.85802 (10)	0.3203 (3)	0.5731 (5)	0.1163 (9)
N1	0.58698 (11)	0.3119 (3)	1.0120 (4)	0.0498 (6)
C2	0.56853 (13)	0.2830 (4)	1.2046 (5)	0.0479 (8)
H2A	0.5795	0.3610	1.3130	0.057*
C3	0.53102 (13)	0.1344 (4)	1.2646 (4)	0.0461 (8)
C4	0.51751 (14)	0.0774 (4)	1.4690 (4)	0.0556 (8)
H4A	0.5300	0.1342	1.5953	0.067*
C5	0.62513 (14)	0.4679 (4)	0.9770 (5)	0.0568 (8)
H5A	0.6336	0.5216	1.1185	0.068*
C6	0.58661 (16)	0.5931 (5)	0.8368 (7)	0.0829 (12)
H6A	0.5476	0.6200	0.9094	0.124*
H6B	0.5777	0.5412	0.6983	0.124*
H6C	0.6105	0.6967	0.8149	0.124*
C7	0.68777 (14)	0.4223 (3)	0.8702 (5)	0.0462 (7)
C8	0.74419 (15)	0.4799 (4)	0.9563 (5)	0.0577 (8)
H8A	0.7436	0.5427	1.0852	0.069*
C9	0.80144 (15)	0.4470 (4)	0.8567 (7)	0.0697 (10)
H9A	0.8391	0.4873	0.9163	0.084*
C10	0.80134 (17)	0.3551 (5)	0.6707 (7)	0.0689 (10)
C11	0.74733 (18)	0.2932 (4)	0.5777 (5)	0.0654 (9)

H11A	0.7489	0.2301	0.4491	0.078*
C12	0.69042 (15)	0.3265 (4)	0.6789 (5)	0.0543 (8)
H12A	0.6532	0.2843	0.6184	0.065*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0499 (6)	0.0559 (6)	0.0379 (5)	-0.0056 (6)	0.000	0.000
F1	0.0687 (14)	0.1167 (19)	0.164 (2)	-0.0047 (14)	0.0485 (15)	-0.030 (2)
N1	0.0452 (14)	0.0502 (15)	0.0540 (16)	-0.0082 (12)	0.0032 (11)	-0.0028 (11)
C2	0.0405 (16)	0.054 (2)	0.0489 (18)	-0.0028 (15)	-0.0043 (14)	-0.0073 (16)
C3	0.0388 (15)	0.0544 (19)	0.0452 (17)	-0.0011 (15)	-0.0013 (13)	-0.0024 (14)
C4	0.056 (2)	0.071 (2)	0.0396 (15)	-0.0115 (15)	-0.0020 (13)	-0.0040 (14)
C5	0.0545 (18)	0.0513 (19)	0.0647 (18)	-0.0104 (16)	0.0112 (15)	-0.0111 (16)
C6	0.065 (2)	0.060 (2)	0.123 (3)	0.0111 (19)	0.023 (2)	0.016 (2)
C7	0.0468 (17)	0.0403 (15)	0.0514 (17)	-0.0063 (14)	0.0002 (14)	0.0001 (13)
C8	0.0576 (19)	0.0492 (17)	0.0662 (19)	-0.0090 (17)	0.0002 (16)	-0.0104 (18)
C9	0.047 (2)	0.064 (2)	0.099 (3)	-0.0114 (17)	-0.0023 (19)	-0.007 (2)
C10	0.054 (2)	0.058 (2)	0.095 (3)	-0.0009 (18)	0.021 (2)	0.001 (2)
C11	0.075 (2)	0.063 (2)	0.0577 (19)	0.000 (2)	0.011 (2)	-0.0053 (18)
C12	0.0499 (18)	0.0573 (19)	0.0557 (18)	-0.0058 (17)	-0.0063 (16)	-0.0020 (16)

*Geometric parameters (Å, °)*

S1—C3 <sup>i</sup>	1.725 (3)	C6—H6A	0.9600
S1—C3	1.725 (3)	C6—H6B	0.9600
F1—C10	1.365 (4)	C6—H6C	0.9600
N1—C2	1.264 (4)	C7—C8	1.378 (4)
N1—C5	1.473 (4)	C7—C12	1.392 (4)
C2—C3	1.450 (4)	C8—C9	1.378 (4)
C2—H2A	0.9300	C8—H8A	0.9300
C3—C4	1.360 (4)	C9—C10	1.347 (5)
C4—C4 <sup>i</sup>	1.414 (6)	C9—H9A	0.9300
C4—H4A	0.9300	C10—C11	1.363 (5)
C5—C7	1.518 (4)	C11—C12	1.377 (4)
C5—C6	1.533 (5)	C11—H11A	0.9300
C5—H5A	0.9800	C12—H12A	0.9300
C3 <sup>i</sup> —S1—C3	91.4 (2)	H6A—C6—H6C	109.5
C2—N1—C5	116.8 (3)	H6B—C6—H6C	109.5
N1—C2—C3	123.2 (3)	C8—C7—C12	117.6 (3)
N1—C2—H2A	118.4	C8—C7—C5	120.8 (3)
C3—C2—H2A	118.4	C12—C7—C5	121.6 (3)
C4—C3—C2	127.5 (3)	C7—C8—C9	121.9 (3)
C4—C3—S1	111.5 (2)	C7—C8—H8A	119.1
C2—C3—S1	120.9 (2)	C9—C8—H8A	119.1
C3—C4—C4 <sup>i</sup>	112.81 (18)	C10—C9—C8	118.2 (3)
C3—C4—H4A	123.6	C10—C9—H9A	120.9

C4 <sup>i</sup> —C4—H4A	123.6	C8—C9—H9A	120.9
N1—C5—C7	110.3 (2)	C9—C10—C11	122.9 (3)
N1—C5—C6	108.4 (2)	C9—C10—F1	118.4 (3)
C7—C5—C6	111.6 (2)	C11—C10—F1	118.7 (3)
N1—C5—H5A	108.8	C10—C11—C12	118.4 (3)
C7—C5—H5A	108.8	C10—C11—H11A	120.8
C6—C5—H5A	108.8	C12—C11—H11A	120.8
C5—C6—H6A	109.5	C11—C12—C7	121.1 (3)
C5—C6—H6B	109.5	C11—C12—H12A	119.5
H6A—C6—H6B	109.5	C7—C12—H12A	119.5
C5—C6—H6C	109.5		
C5—N1—C2—C3	-179.6 (2)	C6—C5—C7—C12	-67.4 (4)
N1—C2—C3—C4	168.6 (3)	C12—C7—C8—C9	1.0 (5)
N1—C2—C3—S1	-8.7 (4)	C5—C7—C8—C9	-176.9 (3)
C3 <sup>i</sup> —S1—C3—C4	-0.29 (16)	C7—C8—C9—C10	-0.4 (5)
C3 <sup>i</sup> —S1—C3—C2	177.4 (3)	C8—C9—C10—C11	-0.2 (5)
C2—C3—C4—C4 <sup>i</sup>	-176.7 (3)	C8—C9—C10—F1	-179.1 (3)
S1—C3—C4—C4 <sup>i</sup>	0.8 (4)	C9—C10—C11—C12	0.0 (6)
C2—N1—C5—C7	124.5 (3)	F1—C10—C11—C12	179.0 (3)
C2—N1—C5—C6	-113.0 (3)	C10—C11—C12—C7	0.6 (5)
N1—C5—C7—C8	-129.0 (3)	C8—C7—C12—C11	-1.1 (4)
C6—C5—C7—C8	110.4 (3)	C5—C7—C12—C11	176.7 (3)
N1—C5—C7—C12	53.2 (4)		

Symmetry code: (i)  $-x+1, -y, z$ .

#### (IV) 2,5-Bis[(S)-(+)-1-(4-chlorophenyl)ethyl]iminomethyl]thiophene

##### Crystal data

C<sub>22</sub>H<sub>20</sub>Cl<sub>2</sub>N<sub>2</sub>S

*M<sub>r</sub>* = 415.36

Orthorhombic, *P*2<sub>1</sub>2<sub>1</sub>2

*a* = 21.893 (2) Å

*b* = 7.9212 (6) Å

*c* = 6.2315 (4) Å

*V* = 1080.66 (15) Å<sup>3</sup>

*Z* = 2

*F*(000) = 432

*D<sub>x</sub>* = 1.276 Mg m<sup>-3</sup>

Melting point: 434 K

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 2744 reflections

θ = 3.7–21.5°

μ = 0.41 mm<sup>-1</sup>

*T* = 298 K

Prism, colorless

0.52 × 0.40 × 0.07 mm

##### Data collection

Agilent Xcalibur (Atlas, Gemini)  
diffractometer

Radiation source: Enhance (Mo) X-ray Source

Graphite monochromator

Detector resolution: 10.5564 pixels mm<sup>-1</sup>

ω scans

Absorption correction: multi-scan

*CrysAlis PRO*, (Agilent, 2013)

*T<sub>min</sub>* = 0.692, *T<sub>max</sub>* = 1.000

14195 measured reflections

2743 independent reflections

1534 reflections with *I* > 2σ(*I*)

*R<sub>int</sub>* = 0.058

θ<sub>max</sub> = 29.5°, θ<sub>min</sub> = 3.3°

*h* = -28→27

*k* = -10→9

*l* = -8→8

Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.052$

$wR(F^2) = 0.117$

$S = 1.01$

2743 reflections

124 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0483P)^2]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.13 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.17 \text{ e } \text{\AA}^{-3}$

Absolute structure: Flack  $x$  determined using  
465 quotients  $[(F^-)-(F)]/[(F^+)+(F)]$  (Parsons *et al.*,  
2013)

Absolute structure parameter: 0.10 (6)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.5000	1.0000	-0.13764 (18)	0.0590 (4)
Cl1	0.15354 (6)	0.67176 (16)	0.4970 (2)	0.1059 (5)
N1	0.41391 (14)	0.6958 (3)	-0.0839 (5)	0.0612 (8)
C2	0.43185 (16)	0.7244 (4)	-0.2735 (6)	0.0589 (10)
H2A	0.4204	0.6496	-0.3812	0.071*
C3	0.46969 (15)	0.8689 (4)	-0.3311 (5)	0.0544 (9)
C4	0.48309 (16)	0.9247 (4)	-0.5338 (5)	0.0640 (10)
H4A	0.4711	0.8688	-0.6581	0.077*
C5	0.37559 (18)	0.5438 (4)	-0.0520 (7)	0.0677 (11)
H5A	0.3625	0.5021	-0.1927	0.081*
C6	0.4148 (2)	0.4087 (5)	0.0553 (9)	0.0986 (17)
H6A	0.4484	0.3803	-0.0369	0.148*
H6B	0.3905	0.3098	0.0810	0.148*
H6C	0.4302	0.4509	0.1892	0.148*
C7	0.31964 (17)	0.5839 (4)	0.0796 (5)	0.0555 (9)
C8	0.26313 (19)	0.5202 (5)	0.0255 (7)	0.0701 (10)
H8A	0.2593	0.4575	-0.0999	0.084*
C9	0.21198 (19)	0.5464 (5)	0.1512 (7)	0.0735 (11)
H9A	0.1744	0.5020	0.1109	0.088*
C10	0.21746 (18)	0.6384 (5)	0.3351 (6)	0.0640 (10)
C11	0.2728 (2)	0.7062 (5)	0.3945 (6)	0.0691 (10)
H11A	0.2761	0.7692	0.5199	0.083*
C12	0.32346 (18)	0.6802 (4)	0.2665 (5)	0.0623 (9)
H12A	0.3607	0.7276	0.3056	0.075*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0694 (9)	0.0596 (7)	0.0482 (6)	-0.0040 (7)	0.000	0.000
Cl1	0.0878 (9)	0.1061 (9)	0.1238 (10)	0.0069 (7)	0.0384 (8)	0.0026 (9)
N1	0.062 (2)	0.0569 (18)	0.0643 (19)	-0.0089 (15)	0.0064 (15)	0.0009 (14)
C2	0.055 (2)	0.060 (2)	0.061 (2)	0.0015 (17)	-0.0031 (19)	-0.0056 (18)

C3	0.050 (2)	0.060 (2)	0.0523 (19)	0.0004 (17)	-0.0008 (16)	-0.0008 (17)
C4	0.063 (3)	0.080 (2)	0.0487 (18)	-0.0130 (18)	-0.0011 (17)	-0.0059 (17)
C5	0.071 (3)	0.057 (2)	0.075 (2)	-0.0098 (18)	0.015 (2)	-0.0050 (18)
C6	0.085 (3)	0.064 (2)	0.147 (4)	0.011 (2)	0.040 (3)	0.019 (3)
C7	0.061 (2)	0.0442 (17)	0.061 (2)	-0.0052 (17)	-0.0039 (18)	0.0020 (16)
C8	0.073 (3)	0.061 (2)	0.076 (2)	-0.017 (2)	-0.002 (2)	-0.014 (2)
C9	0.063 (3)	0.067 (3)	0.091 (3)	-0.0168 (19)	-0.003 (2)	-0.007 (2)
C10	0.065 (3)	0.056 (2)	0.072 (2)	0.0010 (19)	0.007 (2)	0.007 (2)
C11	0.081 (3)	0.068 (2)	0.059 (2)	-0.002 (2)	-0.003 (2)	-0.0067 (18)
C12	0.059 (2)	0.065 (2)	0.063 (2)	-0.004 (2)	-0.010 (2)	-0.002 (2)

*Geometric parameters (Å, °)*

S1—C3	1.724 (3)	C6—H6A	0.9600
S1—C3 <sup>i</sup>	1.724 (3)	C6—H6B	0.9600
C11—C10	1.746 (4)	C6—H6C	0.9600
N1—C2	1.265 (4)	C7—C8	1.378 (5)
N1—C5	1.481 (4)	C7—C12	1.394 (4)
C2—C3	1.458 (5)	C8—C9	1.382 (5)
C2—H2A	0.9300	C8—H8A	0.9300
C3—C4	1.370 (4)	C9—C10	1.363 (5)
C4—C4 <sup>i</sup>	1.404 (7)	C9—H9A	0.9300
C4—H4A	0.9300	C10—C11	1.376 (5)
C5—C7	1.508 (5)	C11—C12	1.382 (5)
C5—C6	1.527 (5)	C11—H11A	0.9300
C5—H5A	0.9800	C12—H12A	0.9300
C3—S1—C3 <sup>i</sup>	91.3 (2)	H6A—C6—H6C	109.5
C2—N1—C5	116.6 (3)	H6B—C6—H6C	109.5
N1—C2—C3	123.2 (3)	C8—C7—C12	117.3 (4)
N1—C2—H2A	118.4	C8—C7—C5	121.3 (3)
C3—C2—H2A	118.4	C12—C7—C5	121.4 (3)
C4—C3—C2	127.0 (3)	C7—C8—C9	122.2 (4)
C4—C3—S1	111.6 (3)	C7—C8—H8A	118.9
C2—C3—S1	121.3 (2)	C9—C8—H8A	118.9
C3—C4—C4 <sup>i</sup>	112.8 (2)	C10—C9—C8	119.0 (4)
C3—C4—H4A	123.6	C10—C9—H9A	120.5
C4 <sup>i</sup> —C4—H4A	123.6	C8—C9—H9A	120.5
N1—C5—C7	111.2 (3)	C9—C10—C11	120.8 (4)
N1—C5—C6	108.1 (3)	C9—C10—C11	119.8 (3)
C7—C5—C6	111.5 (3)	C11—C10—C11	119.4 (3)
N1—C5—H5A	108.7	C10—C11—C12	119.5 (3)
C7—C5—H5A	108.7	C10—C11—H11A	120.2
C6—C5—H5A	108.7	C12—C11—H11A	120.2
C5—C6—H6A	109.5	C11—C12—C7	121.0 (4)
C5—C6—H6B	109.5	C11—C12—H12A	119.5
H6A—C6—H6B	109.5	C7—C12—H12A	119.5
C5—C6—H6C	109.5		

---

C5—N1—C2—C3	179.8 (3)	C6—C5—C7—C12	74.9 (4)
N1—C2—C3—C4	-168.8 (4)	C12—C7—C8—C9	-1.2 (5)
N1—C2—C3—S1	7.3 (5)	C5—C7—C8—C9	175.8 (3)
C3 <sup>i</sup> —S1—C3—C4	0.42 (19)	C7—C8—C9—C10	0.0 (6)
C3 <sup>i</sup> —S1—C3—C2	-176.3 (4)	C8—C9—C10—C11	0.8 (6)
C2—C3—C4—C4 <sup>i</sup>	175.3 (4)	C8—C9—C10—C11	-179.6 (3)
S1—C3—C4—C4 <sup>i</sup>	-1.1 (5)	C9—C10—C11—C12	-0.3 (6)
C2—N1—C5—C7	-132.2 (3)	C11—C10—C11—C12	-179.9 (3)
C2—N1—C5—C6	105.1 (4)	C10—C11—C12—C7	-1.0 (6)
N1—C5—C7—C8	137.3 (4)	C8—C7—C12—C11	1.7 (5)
C6—C5—C7—C8	-102.0 (4)	C5—C7—C12—C11	-175.3 (3)
N1—C5—C7—C12	-45.8 (4)		

---

Symmetry code: (i)  $-x+1, -y+2, z$ .