

# *mer*-Trichloridotris(tetrahydrothiophene- $\kappa$ S)- iridium(III): preparation and comparison with other *mer*-trichloridotris(tetrahydrothiophene- $\kappa$ S)metal complexes

Loren C. Brown, Christine M. DuChane and Joseph S. Merola\*

Received 27 July 2016  
Accepted 9 August 2016

Department of Chemistry, Virginia Tech, Blacksburg, VA 24061, USA. \*Correspondence e-mail: jmerola@vt.edu

Edited by S. Parkin, University of Kentucky, USA

**Keywords:** crystal structure; iridium; tetrahydrothiophene; conformers.

**CCDC reference:** 1495966

**Supporting information:** this article has supporting information at journals.iucr.org/e


The title complex,  $[\text{IrCl}_3(\text{C}_4\text{H}_8\text{S})_3]$ , was prepared according to a literature method. A suitable crystal was obtained by diffusion of pentane into a dichloromethane solution and analyzed by single-crystal X-ray diffraction at 100 K. The title complex is isotypic with *mer*-trichloridotris(tetrahydrothiophene- $\kappa$ S)rhodium(III). However, the orientation of the tetrahydrothiophene rings is different from an earlier report of *mer*-trichloridotris(tetrahydrothiophene- $\kappa$ S)iridium(III) deposited in the Cambridge Structural Database. The  $\text{IrS}_3\text{Cl}_3$  core shows a nearly octahedral structure with various bond angles within  $1\text{--}2^\circ$  of the perfect  $90$  or  $180^\circ$  expected for an octahedron. The structure of the title compound is compared with the previous iridium complex as well as the rhodium and other octahedral metal tris-tetrahydrothiophene compounds previously structurally characterized. DFT calculations were performed, which indicate the *mer* isomer is significantly lower in energy than the *fac* isomer by  $50.1\text{ kJ mol}^{-1}$ , thereby accounting for all compounds in the CSD being of the *mer* geometry. Powder X-ray diffraction of the bulk material showed that the preparation method yielded only the isomorph reported in this communication.

## 1. Chemical context


We have been engaged in various studies of iridium chemistry for many years (Merola, 1997; Merola & Franks, 2015; Merola *et al.*, 2013) and recently had need to find alternate routes to some iridium(III) complexes for our research. An examination of the literature led to the title compound as a possible anhydrous source of iridium(III) that we could use as a starting material (Allen & Wilkinson, 1972). *mer*-Trichloridotris(tetrahydrothiophene- $\kappa$ S)iridium(III) has been mentioned in the literature as a starting material for other organometallic iridium complexes (Hay-Motherwell *et al.*, 1989, 1992, 1990; John *et al.*, 2000, 2001, 2014), and most recently has been the starting material of choice for new emissive materials (Chang *et al.*, 2008, 2011, 2013; Chiu *et al.*, 2009; Hung *et al.*, 2010; Lin, Chang *et al.*, 2011; Lin, Chi *et al.*, 2011; Lin *et al.*, 2012). However, no crystallographic studies had been published on this compound. Given its increasing importance, we decided that a single crystal structure determination of the title compound would be worthwhile.

## 2. Structural commentary

*mer*-Trichloridotris(tetrahydrothiophene- $\kappa$ S)iridium(III) (CCDC refcode 1495966) crystallizes in the  $P2_1/n$  space group




with one molecule in the asymmetric unit (Fig. 1). The core structure (heavy atoms around the iridium) is very close to rigorous octahedral geometry with the largest angular variation [ $\text{Cl1}-\text{Ir1}-\text{Cl33}$ ,  $177.35(3)^\circ$ ] being less than  $2.7^\circ$  from ideal linearity.



The Ir–Cl bond lengths [range 2.3648 (8)–2.3774 (9) Å] are somewhat longer than the Ir–S bonds [range 2.3279 (9)–2.3575 (9) Å], as expected from the slightly larger radius of Cl. A search for Ir–S bonds in the CSD (Groom *et al.*, 2016) and analyzed with *Mercury* (Macrae *et al.*, 2008) found 2566 instances with distances ranging from 2.134 to 2.633 Å and a mean value of 2.358 Å. That places the bond lengths for the title compound slightly above the mean value. Similarly, a *Mercury* data analysis of the CSD for Ir–Cl bond lengths found 3965 instances with distances ranging from 2.121 to 2.816 Å and a mean value of 2.413 Å, which places the Ir–Cl distances for the title compound lower than the mean. This comparison should not be considered as too significant since it was not possible to compare bond lengths only for iridium(III) compounds and the analysis includes quite a few iridium(I) complexes. The tetrahydrothiophene rings are well ordered in the title structure, adopting a puckered conformation consistent with trying to minimize ring strain. Two of the rings are positioned with the center of the ring aligned over a chlorine atom in the structure, while the third is aligned over a sulfur atom of another ring. More will be said about the ring conformations in the *Database survey* section.

### 3. Supramolecular features


An examination of the packing diagrams for the title compound shows no unusual intermolecular features other than van der Waals interactions.



**Figure 1**  
Displacement ellipsoid plot (50% probability) of *mer*-trichloridotris-(tetrahydrothiophene- $\kappa$ S)iridium(III) (CCDC 1495966).

### 4. Database survey

A survey of the CCDC database (Groom *et al.*, 2016) uncovered a number of metal *mer*-tris(THT- $\kappa$ S)metal complexes (THT = tetrahydrothiophene), including one iridium structure deposited as a private communication (CCDC 1438699; Rheingold & Donovan-Merkert, 2015). The deposited structure (CCDC 1438699) packs with very different unit-cell parameters but the overall molecular structure is substantially the same. The results of the different packing, however, are slightly different conformations of two of the three THT ligands, as shown in Fig. 2, a structure overlay calculated in *Mercury* (Macrae *et al.*, 2008). On the other hand, the rhodium(III) complex is isotopic with the title complex with similar unit-cell parameters (CCDC refcode GEZHUU; Clark *et al.*, 1988). Fig. 3 shows an overlay calculated with *Mercury* (Macrae *et al.*, 2008) of the title complex with the rhodium compound, showing the nearly perfect atomic overlay. Ruthenium(III) (VIJYAO; Yapp *et al.*, 1990) and molybdenum(III) (REDXIH; Boorman *et al.*, 1996) complexes were also found in the database, with all showing the same meridional arrangement of ligands with the exception that the ruthenium complex displays disorder from overlapping conformations of one of the THT ligands.




**Figure 2**  
Calculated overlay of two polymorphs of *mer*-trichloridotris(tetrahydrothiophene- $\kappa$ S)iridium(III) (CCDC 1438699 and CCDC 1495966). Structure from this paper shown in yellow.

## 5. Theoretical calculations

We were interested in determining if the bulk material synthesized by this process is of a single polymorph or if both of the iridium structures reported (CCDC 1495966, this report, and CCDC 1438699, Rheingold & Donovan-Merkert, 2015) were present. Fig. 4 shows an overlay of the powder X-ray diffraction pattern for the complex reported here with the powder pattern predicted by *Mercury* (Macrae *et al.*, 2008). The match is very good and quite distinct from the pattern predicted for CCDC 1438699, indicating that the bulk material formed in this process is a single polymorph matching the structure reported here.

One feature that stands out in all cases is that the  $MCl_3(\text{THT})_3$  compounds found in the database adopt the *mer* configuration. Calculations were performed using density functional theory with *Gaussian 09* (Frisch *et al.*, 2009). Full geometry optimization of both the *mer* and *fac* isomers was carried out *via* density functional theory (DFT) with the Becke-3-parameter exchange functional (Becke, 1993) and the Lee–Yang–Parr correlation functional (Lee *et al.*, 1988). Because iridium is not covered in the cc-PVDZ basis set used,




**Figure 3**  
Calculated overlay of *mer*-trichloridotris(tetrahydrothiophene- $\kappa$ S)iridium(III) (CCDC 1495966) in yellow with the isotypical rhodium complex (CCDC GEZHUO) in blue.

computations involving Ir employed Stuttgart/Dresden quasi-relativistic pseudopotentials (Andrae *et al.*, 1990). The difference between the two isomers was quite large with the *mer* isomer being more stable than the *fac* by  $50.1 \text{ kJ mol}^{-1}$ , suggesting the occurrence of only the *mer* isomer for the small set of compounds surveyed may be due to thermodynamic stability.

## 6. Synthesis and crystallization

The title compound was synthesized using a slight modification of a literature procedure (John *et al.*, 2014).  $\text{IrCl}_3 \cdot 3\text{H}_2\text{O}$  (1.00 g, 2.84 mmol) and 2-methoxyethanol (50 mL) were added to a 250 mL round-bottomed flask fitted with a magnetic stir bar and a reflux condenser. Tetrahydrothiophene (1.25 mL, 14.2 mmol) was added all at once with stirring. The resulting suspension was refluxed for 18 h, providing a clear orange solution that gave a yellow precipitate upon cooling to room temperature. Deionized water (75 mL) was added and the suspension was cooled overnight (273 K) before collection on a fine-porosity sintered glass frit. The resulting yellow powder was washed with deionized water (3 x 15 mL) then



**Figure 4**  
Powder X-ray diffraction pattern of title compound collected on a Rigaku Miniflex 600 Powder X-ray diffractometer compared with pattern simulated by Mercury (Macrae *et al.*, 2008). Experimental and simulated patterns scaled to highest intensity peak in each.

cold ethanol (3 x 15 mL). After vacuum drying overnight the yellow powder (1.40 g, 88%) was characterized by  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectroscopy. As the NMR spectra were in agreement with previously reported data, no further purification was necessary. Single crystals for X-ray diffraction were grown by slow diffusion of *n*-pentane into a dichloromethane solution of *mer*- $\text{IrCl}_3(\text{THT})_3$ .

## 7. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 1.

## References

Allen, E. A. & Wilkinson, W. (1972). *J. Chem. Soc. Dalton Trans.* pp. 613–617.  
 Andrae, D., Häussermann, U., Dolg, M., Stoll, H. & Preuss, H. (1990). *Theor. Chim. Acta*, **77**, 123–141.  
 Becke, A. D. (1993). *J. Chem. Phys.* **98**, 5648–5652.  
 Boorman, P. M., Wang, M. & Parvez, M. (1996). *J. Chem. Soc. Dalton Trans.* pp. 4533–4542.  
 Chang, C.-F., Cheng, Y.-M., Chi, Y., Chiu, Y.-C., Lin, C.-C., Lee, G.-H., Chou, P.-T., Chen, C.-C., Chang, C.-H. & Wu, C.-C. (2008). *Angew. Chem. Int. Ed.* **47**, 4542–4545.

**Table 1**  
Experimental details.

Crystal data	
Chemical formula	$[\text{IrCl}_3(\text{C}_4\text{H}_8\text{S})_3]$
$M_r$	563.04
Crystal system, space group	Monoclinic, $P2_1/n$
Temperature (K)	100
$a, b, c$ (Å)	11.9160 (3), 10.2528 (2), 14.9434 (4)
$\beta$ (°)	107.202 (3)
$V$ (Å <sup>3</sup> )	1744.00 (7)
$Z$	4
Radiation type	Mo $K\alpha$
$\mu$ (mm <sup>-1</sup> )	8.46
Crystal size (mm)	0.51 × 0.43 × 0.32
Data collection	
Diffractometer	Rigaku OD Xcalibur Eos Gemini ultra
Absorption correction	Analytical [ <i>CrysAlis PRO</i> (Rigaku Oxford Diffraction, 2015) based on expressions derived by Clark & Reid (1995)]
$T_{\min}, T_{\max}$	0.064, 0.155
No. of measured, independent and observed [ $I > 2\sigma(I)$ ] reflections	19537, 5773, 5062
$R_{\text{int}}$	0.042
$(\sin \theta/\lambda)_{\text{max}}$ (Å <sup>-1</sup> )	0.751
Refinement	
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.029, 0.063, 1.05
No. of reflections	5773
No. of parameters	172
H-atom treatment	H-atom parameters constrained
$\Delta\rho_{\text{max}}, \Delta\rho_{\text{min}}$ (e Å <sup>-3</sup> )	1.54, -1.46

Computer programs: *CrysAlis PRO* (Rigaku Oxford Diffraction, 2015), *SHELXT* (Sheldrick, 2015a), *SHELXL2016* (Sheldrick, 2015b), *OLEX2* (Dolomanov *et al.*, 2009) and *Mercury* (Macrae *et al.*, 2008).

Chang, C.-H., Ho, C.-L., Chang, Y.-S., Lien, I. C., Lin, C.-H., Yang, Y.-W., Liao, J.-L. & Chi, Y. (2013). *J. Mater. Chem. C*, **1**, 2639–2647.  
 Chang, Y.-Y., Hung, J.-Y., Chi, Y., Chyn, J.-P., Chung, M.-W., Lin, C.-L., Chou, P.-T., Lee, G.-H., Chang, C.-H. & Lin, W.-C. (2011). *Inorg. Chem.* **50**, 5075–5084.  
 Chiu, Y.-C., Hung, J.-Y., Chi, Y., Chen, C.-C., Chang, C.-H., Wu, C.-C., Cheng, Y.-M., Yu, Y.-C., Lee, G.-H. & Chou, P.-T. (2009). *Adv. Mater.* **21**, 2221–2225.  
 Clark, R. C. & Reid, J. S. (1995). *Acta Cryst.* **A51**, 887–897.  
 Clark, P. D., Machin, J. H., Richardson, J. F., Dowling, N. I. & Hyne, J. B. (1988). *Inorg. Chem.* **27**, 3526–3529.  
 Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.  
 Frisch, M. J., Trucks, G. W., Schlegel, H. B., Scuseria, G. E., Robb, M. A., Cheeseman, J. R., Scalmani, G., Barone, V., Mennucci, B., Petersson, G. A., Nakatsuji, H., Caricato, M., Li, X., Hratchian, H. P., Izmaylov, A. F., Bloino, J., Zheng, G., Sonnenberg, J. L., Hada, M., Ehara, M., Toyota, K., Fukuda, R., Hasegawa, J., Ishida, M., Nakajima, T., Honda, Y., Kitao, O., Nakai, H., Vreven, T., Montgomery, J. J. A., Peralta, J. E., Ogliaro, F., Bearpark, M., Heyd, J. J., Brothers, E., Kudin, K. N., Staroverov, V. N., Kobayashi, R., Normand, J., Raghavachari, K., Rendell, A., Burant, J. C., Iyengar, S. S., Tomasi, J., Cossi, M., Rega, N., Millam, J. M., Klene, M., Knox, J. E., Cross, J. B., Bakken, V., Adamo, C., Jaramillo, J., Gomperts, R., Stratmann, R. E., Yazyev, O., Austin, A. J., Cammi, R., Pomelli, C., Ochterski, J. W., Martin, R. L., Morokuma, K., Zakrzewski, V. G., Voth, G. A., Salvador, P., Dannenberg, J. J., Dapprich, S., Daniels, A. D., Farkas, Ö., Foresman, J. B., Ortiz, J. V., Cioslowski, J. & Fox, D. J. (2009). *Gaussian 09*, Revision A. 1. Gaussian Inc., Pittsburgh, Pennsylvania, USA.

- Groom, C. R., Bruno, I. J., Lightfoot, M. P. & Ward, S. C. (2016). *Acta Cryst.* **B72**, 171–179.
- Hay-Motherwell, R. S., Wilkinson, G., Hussain, B. & Hursthouse, M. B. (1989). *J. Chem. Soc. Chem. Commun.* pp. 1436–1437.
- Hay-Motherwell, R. S., Wilkinson, G., Hussain-Bates, B. & Hursthouse, M. B. (1990). *Polyhedron*, **9**, 2071–2080.
- Hay-Motherwell, R. S., Wilkinson, G., Hussain-Bates, B. & Hursthouse, M. B. (1992). *J. Chem. Soc. Dalton Trans.* pp. 3477–3482.
- Hung, J.-Y., Lin, C.-H., Chi, Y., Chung, M.-W., Chen, Y.-J., Lee, G.-H., Chou, P.-T., Chen, C.-C. & Wu, C.-C. (2010). *J. Mater. Chem.* **20**, 7682–7693.
- John, K. D., Eglin, J. L., Salazar, K. V., Baker, R. T., Sattelberger, A. P., Serra, D. & White, L. M. (2014). *Inorg. Synth.* **36**, 165–171.
- John, K. D., Salazar, K. V., Scott, B. L., Baker, R. T. & Sattelberger, A. P. (2001). *Organometallics*, **20**, 296–304.
- John, K. D., Scott, B. L., Baker, R. T., Sattelberger, A. P. & Salazar, K. V. (2000). *Chem. Commun.* pp. 581–582.
- Lee, C., Yang, W. & Parr, R. G. (1988). *Phys. Rev. B*, **37**, 785–789.
- Lin, C.-H., Chang, Y.-Y., Hung, J.-Y., Lin, C.-Y., Chi, Y., Chung, M.-W., Lin, C.-L., Chou, P.-T., Lee, G.-H., Chang, C.-H. & Lin, W.-C. (2011). *Angew. Chem. Int. Ed.* **50**, 3182–3186.
- Lin, C.-H., Chi, Y., Chung, M.-W., Chen, Y.-J., Wang, K.-W., Lee, G.-H., Chou, P.-T., Hung, W.-Y. & Chiu, H.-C. (2011). *Dalton Trans.* **40**, 1132–1143.
- Lin, C.-H., Chiu, Y.-C., Chi, Y., Tao, Y.-T., Liao, L.-S., Tseng, M.-R. & Lee, G.-H. (2012). *Organometallics*, **31**, 4349–4355.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Merola, J. S. (1997). *Curr. Org. Chem.* **1**, 235–248.
- Merola, J. S. & Franks, M. A. (2015). *Acta Cryst.* **E71**, 226–230.
- Merola, J. S., Franks, M. A. & Frazier, J. F. (2013). *Polyhedron*, **54**, 67–73.
- Rheingold, A. L. & Donovan-Merkert, B. (2015). Private communication (refcode 1438699). CCDC, Cambridge, England.
- Rigaku Oxford Diffraction (2015). *CrysAlis PRO*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2015a). *Acta Cryst.* **A71**, 3–8.
- Sheldrick, G. M. (2015b). *Acta Cryst.* **C71**, 3–8.
- Yapp, D. T. T., Jaswal, J., Rettig, S. J., James, B. R. & Skov, K. A. (1990). *Inorg. Chim. Acta*, **177**, 199–208.

## supporting information

*Acta Cryst.* (2016). E72, 1305-1309 [https://doi.org/10.1107/S2056989016012883]

## ***mer*-Trichloridotris(tetrahydrothiophene- $\kappa$ S)iridium(III): preparation and comparison with other *mer*-trichloridotris(tetrahydrothiophene- $\kappa$ S)metal complexes**

**Loren C. Brown, Christine M. DuChane and Joseph S. Merola**

### Computing details

Data collection: *CrysAlis PRO* (Rigaku Oxford Diffraction, 2015); cell refinement: *CrysAlis PRO* (Rigaku Oxford Diffraction, 2015); data reduction: *CrysAlis PRO* (Rigaku Oxford Diffraction, 2015); program(s) used to solve structure: SHELXT (Sheldrick, 2015a); program(s) used to refine structure: *SHELXL2016* (Sheldrick, 2015b); molecular graphics: *OLEX2* (Dolomanov *et al.*, 2009); software used to prepare material for publication: *OLEX2* (Dolomanov *et al.*, 2009) and *Mercury* (Macrae *et al.*, 2008).

### *mer*-Trichloridotris(tetrahydrothiophene- $\kappa$ S)iridium(III)

#### Crystal data

[IrCl<sub>3</sub>(C<sub>4</sub>H<sub>8</sub>S)<sub>3</sub>]  
 $M_r = 563.04$   
 Monoclinic,  $P2_1/n$   
 $a = 11.9160$  (3) Å  
 $b = 10.2528$  (2) Å  
 $c = 14.9434$  (4) Å  
 $\beta = 107.202$  (3)°  
 $V = 1744.00$  (7) Å<sup>3</sup>  
 $Z = 4$

$F(000) = 1088$   
 $D_x = 2.144$  Mg m<sup>-3</sup>  
 Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å  
 Cell parameters from 9679 reflections  
 $\theta = 4.1$ – $32.2^\circ$   
 $\mu = 8.46$  mm<sup>-1</sup>  
 $T = 100$  K  
 Cube, yellow  
 $0.51 \times 0.43 \times 0.32$  mm

#### Data collection

Rigaku OD Xcalibur Eos Gemini ultra  
 diffractometer  
 Radiation source: fine-focus sealed X-ray tube,  
 Enhance (Mo) X-ray Source  
 Graphite monochromator  
 Detector resolution: 8.0061 pixels mm<sup>-1</sup>  
 $\omega$  scans

Absorption correction: analytical  
 [CrysAlis PRO (Rigaku Oxford Diffraction,  
 2015) based on expressions derived by Clark &  
 Reid (1995)]  
 $T_{\min} = 0.064$ ,  $T_{\max} = 0.155$   
 19537 measured reflections  
 5773 independent reflections  
 5062 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.042$   
 $\theta_{\max} = 32.2^\circ$ ,  $\theta_{\min} = 3.6^\circ$   
 $h = -13 \rightarrow 17$   
 $k = -13 \rightarrow 15$   
 $l = -22 \rightarrow 21$



Refinement

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.029$   
 $wR(F^2) = 0.063$   
 $S = 1.05$   
 5773 reflections  
 172 parameters  
 0 restraints

Primary atom site location: structure-invariant  
 direct methods  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0223P)^2 + 0.8135P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.002$   
 $\Delta\rho_{\max} = 1.54 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -1.46 \text{ e } \text{\AA}^{-3}$

Special details

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Ir1	0.45092 (2)	0.70407 (2)	0.73222 (2)	0.01133 (4)
Cl1	0.59718 (7)	0.70422 (8)	0.65431 (6)	0.01922 (17)
Cl2	0.59716 (8)	0.73216 (8)	0.87865 (6)	0.01887 (16)
Cl3	0.30344 (7)	0.71455 (8)	0.80946 (6)	0.01688 (16)
S1	0.46678 (7)	0.47667 (8)	0.74921 (6)	0.01509 (16)
S2	0.30373 (7)	0.68096 (8)	0.59053 (6)	0.01540 (16)
S3	0.44035 (7)	0.93361 (9)	0.72529 (6)	0.01636 (16)
C1	0.6230 (3)	0.4314 (4)	0.7907 (3)	0.0236 (8)
H1A	0.6388	0.3570	0.7540	0.028*
H1B	0.6727	0.5059	0.7837	0.028*
C2	0.6496 (3)	0.3940 (4)	0.8934 (3)	0.0281 (9)
H2A	0.7157	0.3314	0.9114	0.034*
H2B	0.6710	0.4722	0.9337	0.034*
C3	0.5384 (4)	0.3321 (4)	0.9044 (3)	0.0257 (8)
H3A	0.5449	0.3203	0.9715	0.031*
H3B	0.5246	0.2459	0.8732	0.031*
C4	0.4380 (3)	0.4262 (4)	0.8582 (3)	0.0216 (7)
H4A	0.4387	0.5024	0.8991	0.026*
H4B	0.3610	0.3820	0.8448	0.026*
C5	0.3455 (3)	0.5674 (4)	0.5104 (3)	0.0277 (9)
H5A	0.3271	0.6057	0.4469	0.033*
H5B	0.4308	0.5488	0.5331	0.033*
C6	0.2749 (3)	0.4419 (4)	0.5082 (3)	0.0261 (8)
H6A	0.2620	0.3969	0.4474	0.031*
H6B	0.3179	0.3822	0.5588	0.031*
C7	0.1582 (3)	0.4802 (4)	0.5219 (3)	0.0225 (8)
H7A	0.1078	0.5244	0.4654	0.027*
H7B	0.1163	0.4024	0.5348	0.027*
C8	0.1883 (3)	0.5720 (4)	0.6051 (3)	0.0192 (7)

H8A	0.2167	0.5224	0.6643	0.023*
H8B	0.1182	0.6229	0.6065	0.023*
C9	0.3843 (3)	0.9968 (4)	0.6060 (2)	0.0188 (7)
H9A	0.3912	0.9305	0.5597	0.023*
H9B	0.3010	1.0232	0.5920	0.023*
C10	0.4624 (3)	1.1147 (4)	0.6047 (3)	0.0213 (7)
H10A	0.4527	1.1433	0.5396	0.026*
H10B	0.4424	1.1883	0.6400	0.026*
C11	0.5875 (3)	1.0686 (4)	0.6510 (3)	0.0228 (8)
H11A	0.6424	1.1434	0.6631	0.027*
H11B	0.6119	1.0058	0.6099	0.027*
C12	0.5882 (3)	1.0027 (4)	0.7436 (3)	0.0209 (7)
H12A	0.6061	1.0674	0.7951	0.025*
H12B	0.6483	0.9330	0.7598	0.025*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Ir1	0.01159 (6)	0.01222 (7)	0.01049 (6)	0.00034 (4)	0.00376 (5)	-0.00076 (4)
Cl1	0.0165 (4)	0.0225 (4)	0.0222 (4)	-0.0003 (3)	0.0112 (3)	-0.0008 (3)
Cl2	0.0199 (4)	0.0171 (4)	0.0157 (4)	-0.0013 (3)	-0.0007 (3)	-0.0021 (3)
Cl3	0.0183 (4)	0.0179 (4)	0.0177 (4)	0.0004 (3)	0.0102 (3)	-0.0020 (3)
S1	0.0158 (4)	0.0132 (4)	0.0156 (4)	0.0007 (3)	0.0035 (3)	-0.0012 (3)
S2	0.0151 (4)	0.0170 (4)	0.0130 (4)	0.0001 (3)	0.0024 (3)	0.0001 (3)
S3	0.0202 (4)	0.0152 (4)	0.0157 (4)	0.0008 (3)	0.0084 (3)	-0.0004 (3)
C1	0.0173 (17)	0.0225 (18)	0.032 (2)	0.0057 (14)	0.0092 (16)	0.0040 (16)
C2	0.0209 (18)	0.025 (2)	0.030 (2)	0.0032 (15)	-0.0046 (17)	0.0039 (17)
C3	0.031 (2)	0.0182 (18)	0.026 (2)	0.0064 (15)	0.0056 (17)	0.0081 (16)
C4	0.0242 (18)	0.0193 (18)	0.0240 (18)	0.0017 (14)	0.0113 (16)	0.0053 (15)
C5	0.0195 (18)	0.046 (3)	0.0179 (18)	0.0012 (17)	0.0059 (15)	-0.0126 (17)
C6	0.032 (2)	0.0249 (19)	0.0184 (18)	0.0067 (16)	0.0027 (16)	-0.0076 (16)
C7	0.0290 (19)	0.0168 (17)	0.0180 (17)	-0.0032 (14)	0.0015 (16)	-0.0019 (14)
C8	0.0143 (15)	0.0214 (17)	0.0238 (18)	-0.0031 (13)	0.0082 (14)	-0.0051 (15)
C9	0.0191 (17)	0.0207 (17)	0.0149 (16)	0.0011 (13)	0.0023 (14)	0.0031 (13)
C10	0.0262 (18)	0.0181 (17)	0.0212 (18)	0.0021 (14)	0.0097 (16)	0.0024 (15)
C11	0.0222 (18)	0.0221 (18)	0.0260 (19)	-0.0030 (14)	0.0100 (16)	0.0021 (15)
C12	0.0189 (17)	0.0199 (18)	0.0209 (18)	-0.0037 (13)	0.0013 (15)	-0.0008 (14)

*Geometric parameters (Å, °)*

Ir1—C11	2.3648 (8)	C5—H5A	0.9900
Ir1—C12	2.3774 (9)	C5—H5B	0.9900
Ir1—C13	2.3732 (8)	C5—C6	1.533 (6)
Ir1—S1	2.3469 (9)	C6—H6A	0.9900
Ir1—S2	2.3279 (9)	C6—H6B	0.9900
Ir1—S3	2.3575 (9)	C6—C7	1.516 (5)
S1—C1	1.839 (4)	C7—H7A	0.9900
S1—C4	1.835 (4)	C7—H7B	0.9900



S2—C5	1.841 (4)	C7—C8	1.515 (5)
S2—C8	1.834 (3)	C8—H8A	0.9900
S3—C9	1.827 (4)	C8—H8B	0.9900
S3—C12	1.843 (4)	C9—H9A	0.9900
C1—H1A	0.9900	C9—H9B	0.9900
C1—H1B	0.9900	C9—C10	1.529 (5)
C1—C2	1.522 (6)	C10—H10A	0.9900
C2—H2A	0.9900	C10—H10B	0.9900
C2—H2B	0.9900	C10—C11	1.521 (5)
C2—C3	1.521 (6)	C11—H11A	0.9900
C3—H3A	0.9900	C11—H11B	0.9900
C3—H3B	0.9900	C11—C12	1.537 (5)
C3—C4	1.533 (5)	C12—H12A	0.9900
C4—H4A	0.9900	C12—H12B	0.9900
C4—H4B	0.9900		
C11—Ir1—C12	90.39 (3)	S2—C5—H5A	110.3
C11—Ir1—C13	177.35 (3)	S2—C5—H5B	110.3
C13—Ir1—C12	89.63 (3)	H5A—C5—H5B	108.6
S1—Ir1—C11	90.37 (3)	C6—C5—S2	107.0 (2)
S1—Ir1—C12	90.41 (3)	C6—C5—H5A	110.3
S1—Ir1—C13	92.28 (3)	C6—C5—H5B	110.3
S1—Ir1—S3	176.44 (3)	C5—C6—H6A	110.2
S2—Ir1—C11	91.09 (3)	C5—C6—H6B	110.2
S2—Ir1—C12	178.15 (3)	H6A—C6—H6B	108.5
S2—Ir1—C13	88.85 (3)	C7—C6—C5	107.4 (3)
S2—Ir1—S1	90.70 (3)	C7—C6—H6A	110.2
S2—Ir1—S3	92.61 (3)	C7—C6—H6B	110.2
S3—Ir1—C11	90.88 (3)	C6—C7—H7A	110.6
S3—Ir1—C12	86.25 (3)	C6—C7—H7B	110.6
S3—Ir1—C13	86.47 (3)	H7A—C7—H7B	108.8
C1—S1—Ir1	109.15 (13)	C8—C7—C6	105.5 (3)
C4—S1—Ir1	110.23 (12)	C8—C7—H7A	110.6
C4—S1—C1	93.79 (17)	C8—C7—H7B	110.6
C5—S2—Ir1	112.32 (13)	S2—C8—H8A	110.4
C8—S2—Ir1	110.12 (13)	S2—C8—H8B	110.4
C8—S2—C5	92.79 (17)	C7—C8—S2	106.6 (2)
C9—S3—Ir1	113.33 (12)	C7—C8—H8A	110.4
C9—S3—C12	93.79 (16)	C7—C8—H8B	110.4
C12—S3—Ir1	109.93 (12)	H8A—C8—H8B	108.6
S1—C1—H1A	110.3	S3—C9—H9A	110.9
S1—C1—H1B	110.3	S3—C9—H9B	110.9
H1A—C1—H1B	108.6	H9A—C9—H9B	108.9
C2—C1—S1	106.9 (3)	C10—C9—S3	104.2 (2)
C2—C1—H1A	110.3	C10—C9—H9A	110.9
C2—C1—H1B	110.3	C10—C9—H9B	110.9
C1—C2—H2A	110.4	C9—C10—H10A	110.7
C1—C2—H2B	110.4	C9—C10—H10B	110.7

---

H2A—C2—H2B	108.6	H10A—C10—H10B	108.8
C3—C2—C1	106.7 (3)	C11—C10—C9	105.5 (3)
C3—C2—H2A	110.4	C11—C10—H10A	110.7
C3—C2—H2B	110.4	C11—C10—H10B	110.7
C2—C3—H3A	110.5	C10—C11—H11A	110.4
C2—C3—H3B	110.5	C10—C11—H11B	110.4
C2—C3—C4	106.1 (3)	C10—C11—C12	106.8 (3)
H3A—C3—H3B	108.7	H11A—C11—H11B	108.6
C4—C3—H3A	110.5	C12—C11—H11A	110.4
C4—C3—H3B	110.5	C12—C11—H11B	110.4
S1—C4—H4A	110.8	S3—C12—H12A	110.4
S1—C4—H4B	110.8	S3—C12—H12B	110.4
C3—C4—S1	104.6 (2)	C11—C12—S3	106.5 (2)
C3—C4—H4A	110.8	C11—C12—H12A	110.4
C3—C4—H4B	110.8	C11—C12—H12B	110.4
H4A—C4—H4B	108.9	H12A—C12—H12B	108.6
Ir1—S1—C1—C2	106.1 (3)	C2—C3—C4—S1	43.0 (4)
Ir1—S1—C4—C3	-132.6 (2)	C4—S1—C1—C2	-6.9 (3)
Ir1—S2—C5—C6	106.8 (2)	C5—S2—C8—C7	-20.6 (3)
Ir1—S2—C8—C7	-135.6 (2)	C5—C6—C7—C8	-48.0 (4)
Ir1—S3—C9—C10	-138.6 (2)	C6—C7—C8—S2	42.1 (3)
Ir1—S3—C12—C11	114.0 (2)	C8—S2—C5—C6	-6.3 (3)
S1—C1—C2—C3	33.0 (4)	C9—S3—C12—C11	-2.5 (3)
S2—C5—C6—C7	31.9 (4)	C9—C10—C11—C12	-49.9 (4)
S3—C9—C10—C11	46.4 (3)	C10—C11—C12—S3	29.9 (4)
C1—S1—C4—C3	-20.5 (3)	C12—S3—C9—C10	-25.0 (3)
C1—C2—C3—C4	-49.8 (4)		

---