



Ruthenium(II) carbonyl compounds with the 4'-chloro-2,2':6',2''-terpyridine ligand

Rajendhrasrad Tatikonda and Matti Haukka*

University of Jyväskylä, Department of Chemistry, PO Box 35, FI-40014 University of Jyväskylä, Finland.

*Correspondence e-mail: matti.o.haukka@jyu.fi

Received 7 February 2017

Accepted 10 March 2017


Edited by M. Weil, Vienna University of
Technology, Austria**Keywords:** crystal structure; ruthenium; terpyridine ligand; carbonyl ligand.**CCDC references:** 1537190; 1537189**Supporting information:** this article has supporting information at journals.iucr.org/e

Two ruthenium carbonyl complexes with the 4'-chloro-2,2':6',2''-terpyridine ligand (tpy-Cl, C₁₅H₁₀ClN₃), *i.e.* [RuCl(tpy-Cl)(CO)₂][RuCl₃(CO)₃] (I) [systematic name: *cis*-dicarbonylchlorido(4'-chloro-2,2':6',2''-terpyridine-κ³N)-ruthenium(II) *fac*-tricarbonyltrichloridoruthenate(II)], and [RuCl₂(tpy-Cl)(CO)₂] (II) [*cis*-dicarbonyl-*trans*-dichlorido(4'-chloro-2,2':6',2''-terpyridine-κ²N¹,N^{1'})ruthenium(II)], were synthesized and characterized by single-crystal X-ray diffraction. The Ru^{II} atoms in both centrosymmetric structures (I) and (II) display similar, slightly distorted octahedral coordination spheres. The coordination sphere in the complex cation in compound (I) is defined by three N atoms of the tridentate tpy-Cl ligand, two carbonyl carbon atoms and one chlorido ligand; the charge is balanced by an octahedral [Ru(CO)₃Cl₃][−] counteranion. In the neutral compound (II), the tpy-Cl ligand coordinates to the metal only through two of its N atoms. The coordination sphere of the Ru^{II} atom is completed by two carbonyl and two chlorido ligands. In the crystal structures of both (I) and (II), weak C—H...Cl interactions are observed.


1. Chemical context

Ruthenium-carbonyl compounds with polypyridine ligands are known to be active catalysts for several catalytic processes including the reduction of carbon dioxide (Collomb-Dunand-Sauthier *et al.*, 1994; Chardon-Noblat *et al.*, 2002; Kuramochi *et al.*, 2015), water-gas shift reaction (Luukkanen *et al.*, 1999) and hydroformylation (Alvila *et al.*, 1994). Many of these systems are metallopolymers obtained by reducing mononuclear precursors either chemically or electrochemically. The 2,2'-bipyridine ligand or its derivatives are the most commonly used ligand systems in these catalysts. It is also reported that possible substituents on polypyridine rings can have a strong impact on the catalytic behaviour of the compounds (Chardon-Noblat *et al.*, 2001), which could offer a route to tailor the catalytic activity. Compounds with terpyridine and its derivatives as ligands together with carbonyl ligands are less commonly used (Deacon *et al.*, 1984; Gibson *et al.*, 1997; Ziessel *et al.*, 2004), although it has also been shown that these types of compounds can be used to obtain active catalysts. Terpyridines are able to act as strong tridentate ligands because of the arrangement of the pyridine nitrogen atoms. However, bidentate coordination is also known (Deacon *et al.*, 1984; Kooijman *et al.*, 2007; Amoroso *et al.*, 2010).

In this context we report on the two title compounds, [RuCl(tpy-Cl)(CO)₂][Ru(CO)₃Cl₃] (I) and [RuCl₂(tpy-Cl)(CO)₂] (II) with the 4'-chloro-2,2':6',2''-terpyridine ligand (tpy-Cl, C₁₅H₁₀ClN₃), which show both types of coordination, *i.e.* tridentate for (I) and bidentate for (II). The title



compounds were synthesized by adopting a literature procedure (Homanen *et al.*, 1996).



2. Structural commentary

Compound (I) is a salt and crystallizes in the monoclinic space group $P2_1/c$ with four formula units in the unit cell. The coordination sphere of the Ru^{II} atom in the cation is a slightly distorted octahedron. The equatorial positions are occupied by three pyridine N atoms from the Tpy-Cl ligand and by one carbonyl ligand; axial positions are occupied by one chloride and one carbonyl ligand. The charge on the Ru^{II} atom is balanced by an octahedrally shaped $\text{fac}[\text{Ru}(\text{CO})_3\text{Cl}_3]^-$ anion (Fig. 1). As expected, in the cation the Ru1–N5 bond to the central pyridine ring of the tpy-Cl ligand [2.019 (2) Å] is the shortest of the Ru–N bonds (Gibson *et al.*, 1997; Ziessel *et al.*, 2004). The Ru1–N1 [2.097 (2) Å] and Ru1–N15 [2.093 (2) Å] bonds involving the outer pyridine rings are lengthened to relieve strain and to retain a typical terpyridine bite angle of about 79°. Similar structures can be found in other ruthenium(II) complexes containing terpyridine ligands (Gibson *et al.*, 1997). The Ru1–C2 bond of the equatorial carbonyl group [1.918 (3) Å] is longer than the Ru1–C1 bond [1.893 (3) Å] of the axial carbonyl group, indicating a slightly stronger *trans*-influence caused by the pyridine N atom. The Ru1–Cl1 distance [2.4279 (7) Å] is in the range of typical Ru–Cl bond lengths (Deacon *et al.*, 1984; Ziessel *et al.*, 2004). The corresponding Ru–Cl bond lengths in the $[\text{Ru}(\text{CO})_3\text{Cl}_3]^-$ counter-anion [2.4129 (7)–2.4212 (7) Å] also fall into the typical range of Ru–Cl bonds (Table 1).




Figure 1

The molecular structures of the cation and anion in compound (I). Displacement ellipsoids are drawn at the 50% probability level.

Table 1

Selected bond lengths (Å) for (I).

Ru1–C1	1.893 (3)	Ru2–C20	1.902 (3)
Ru1–C2	1.918 (3)	Ru2–C18	1.914 (3)
Ru1–N5	2.019 (2)	Ru2–Cl4	2.4129 (7)
Ru1–N15	2.093 (2)	Ru2–Cl5	2.4199 (7)
Ru1–N1	2.097 (2)	Ru2–Cl3	2.4212 (7)
Ru1–Cl1	2.4279 (7)	N1–C3	1.336 (3)
Ru2–C19	1.893 (3)		

Table 2

Selected bond lengths (Å) for (II).

Ru1–C2	1.877 (3)	Ru1–N2	2.157 (2)
Ru1–C1	1.895 (3)	Ru1–Cl1	2.3762 (8)
Ru1–N1	2.105 (2)	Ru1–Cl2	2.4098 (7)

Compound (II) is a neutral complex and crystallizes in the triclinic space group $P\bar{1}$ with two formula units. The coordination sphere around the Ru^{II} atom is again a slightly distorted octahedron (Fig. 2). The four equatorial positions are occupied by two N atoms [Ru1–N1 = 2.105 (2) and Ru1–N2 = 2.157 (2) Å] from the Tpy-Cl ligand and by two carbonyl ligands [Ru1–C2 = 1.877 (3); Ru1–C1 = 1.895 (3) Å]. The chlorido ligands [Ru1–Cl1 = 2.3762 (8); Ru1–Cl2 = 2.4098 (7) Å] are placed at axial positions of the molecule. The Ru1–N2 and Ru1–C1 bond lengths are slightly longer than Ru1–N1 and Ru1–C2 bond lengths due to the steric strain generated by the non-coordinating pyridine ring (Table 2).

The Tpy-Cl ligand in compound (I) is non-planar, despite coordination of all its three N atoms [dihedral angles between the mean planes of the central pyridine ring and the adjacent pyridine rings are 5.70 (8) and 13.28 (7)°]. In compound (II), the ring with the non-coordinating N atom is inclined considerably relative to the coordination plane of the two pyridine rings [dihedral angle 57.71 (9)°].

3. Supramolecular features

The packing of molecules (I) and (II) are dominated by van der Waals interactions; packing plots are displayed in Fig. 3 for




Figure 2

The molecular structure of compound (II). Displacement ellipsoids are drawn at the 50% probability level.




Figure 3
The crystal packing of (I) in a view along the *b* axis.

(I) and Fig. 4 for (II). Only weak hydrogen bonds and π - π contacts can be found in these structures. In both (I) and (II), some non-conventional hydrogen bonds between the aromatic C-H hydrogen atoms and chlorido ligands of neighboring molecules do exist. The shortest contacts are summarized in Tables 3 and 4. In addition to these hydrogen bonds, the aromatic rings in structure (I) are involved in weak face-to-face π - π -interactions with considerable offsets. The shortest intermolecular C-C distances range from 3.23 to 3.50 Å. In (II), an edge-to-face contact exists between C3-H3 and C16 of the neighboring molecule. The distance between H3 and C16 is 2.89 Å and the angle C3-H3...C16 amounts to 134°.

Table 3
Hydrogen-bond geometry (Å, °) for (I).

<i>D</i> -H... <i>A</i>	<i>D</i> -H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> -H... <i>A</i>
C11-H11...Cl5 ⁱ	0.95	2.76	3.664 (3)	158
C16-H16...Cl1 ⁱⁱ	0.95	2.72	3.515 (3)	142
C5-H5...Cl3 ⁱⁱⁱ	0.95	2.82	3.553 (3)	134

Symmetry codes: (i) $-x, -y + 1, -z$; (ii) $-x, y + \frac{1}{2}, -z + \frac{1}{2}$; (iii) $x, y - 1, z$.

Table 4
Hydrogen-bond geometry (Å, °) for (II).

<i>D</i> -H... <i>A</i>	<i>D</i> -H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> -H... <i>A</i>
C9-H9...Cl2 ⁱ	0.95	2.77	3.687 (3)	163

Symmetry code: (i) $-x + 2, -y + 1, -z + 1$.

All interactions considered, three-dimensional network structures are obtained both for (I) and (II).

4. Synthesis and crystallization

The title compounds were synthesized using a literature procedure (Homanen *et al.*, 1996) and both compounds were obtained in a single pot reaction. A solution of $[\text{Ru}(\text{CO})_3\text{Cl}_2]_2$ (25.6 mg, 0.05 mmol) in 3 ml of THF was refluxed for 1 h under argon gas. After the reaction time, 26.7 mg (0.1 mmol) of tpy-Cl in 3 ml of THF was added to the above reaction mixture. The resulting mixture was refluxed for another 2 h in air with continuous stirring. During the reaction, the pale yellow solution turned to a reddish solution with a colourless precipitate. The precipitate was collected through centrifugation and the filtrate was evaporated for crystallization. Compound (I) was obtained as a major product originating from the precipitate and compound (II) was collected as a minor product from the filtrate. High-quality crystals of the salt (I) for single-crystal X-ray diffraction were obtained from




Figure 4
The crystal packing of (II) in a view along the *b* axis.

Table 5
Experimental details.

	(I)	(II)
Crystal data		
Chemical formula	[RuCl(C ₁₅ H ₁₀ ClN ₃)(CO) ₂][Ru(CO) ₃ Cl ₃]	[RuCl ₂ (C ₁₅ H ₁₀ ClN ₃ (CO) ₂]
<i>M_r</i>	751.70	495.70
Crystal system, space group	Monoclinic, <i>P</i> ₂ ₁ / <i>c</i>	Triclinic, <i>P</i> $\bar{1}$
Temperature (K)	123	123
<i>a</i> , <i>b</i> , <i>c</i> (Å)	14.3578 (4), 13.9158 (2), 13.2220 (3)	7.3019 (3), 8.5080 (3), 14.7702 (6)
α , β , γ (°)	90, 114.080 (3), 90	101.287 (3), 91.835 (3), 98.144 (3)
<i>V</i> (Å ³)	2411.86 (11)	889.09 (6)
<i>Z</i>	4	2
Radiation type	Mo <i>K</i> α	Mo <i>K</i> α
μ (mm ⁻¹)	1.85	1.35
Crystal size (mm)	0.34 × 0.08 × 0.06	0.30 × 0.08 × 0.05
Data collection		
Diffractometer	Agilent SuperNova, Dual, Cu at zero, Atlas	Agilent SuperNova, Dual, Cu at zero, Atlas
Absorption correction	Multi-scan (<i>CrysAlis PRO</i> ; Agilent, 2013)	Multi-scan (<i>CrysAlis PRO</i> ; Agilent, 2013)
<i>T</i> _{min} , <i>T</i> _{max}	0.914, 1.000	0.300, 1.000
No. of measured, independent and observed [<i>I</i> > 2 σ (<i>I</i>)] reflections	11072, 4864, 4264	7508, 3662, 3405
<i>R</i> _{int}	0.023	0.036
(<i>sin</i> θ / λ) _{max} (Å ⁻¹)	0.625	0.630
Refinement		
<i>R</i> [<i>F</i> ² > 2 σ (<i>F</i> ²)], <i>wR</i> (<i>F</i> ²), <i>S</i>	0.024, 0.050, 1.06	0.033, 0.088, 1.07
No. of reflections	4864	3662
No. of parameters	316	235
H-atom treatment	H-atom parameters constrained	H-atom parameters constrained
$\Delta\rho_{\text{max}}$, $\Delta\rho_{\text{min}}$ (e Å ⁻³)	0.43, -0.48	0.74, -1.43

Computer programs: *CrysAlis PRO* (Agilent, 2013), *SUPERFLIP* (Palatinus & Chapuis, 2007), *SHELXL2014* (Sheldrick, 2015) and *UCSF Chimera* (Pettersen *et al.*, 2004).

DMSO solution and those of complex (II) were obtained as brown-coloured crystals from the filtrate.

5. Refinement details

Crystal data, data collection and structure refinement details are summarized in Table 5. All H atoms were positioned in calculated positions and constrained to ride on their parent atoms, with C–H = 0.95 Å and *U*_{iso} = 1.2*U*_{eq}(C). The maximum electron density in complex (I) is located at 0.67 Å from atom C8 and in complex (II) at 1.28 Å from atom N2, respectively. The minimum density in complex (I) is located at 0.77 Å from atom Ru1 and in complex (II) at 0.87 Å from atom Ru1, respectively.

Funding information

Funding for this research was provided by: Academy of Finland (award No. 295581); COST Action 1302, 'Smart Inorganic Polymers'.

References

Agilent (2013). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.
 Alvila, L., Pursiainen, J., Kiviaho, J., Pakkanen, T. A. & Krause, O. (1994). *J. Mol. Catal.* **91**, 335–342.

Amoroso, A. J., Banu, A., Coogan, M. P., Edwards, P. G., Hossain, G. & Malik, K. M. A. (2010). *Dalton Trans.* **39**, 6993–7003.
 Chardon-Noblat, S., Da Costa, P., Deronzier, A., Maniguet, S. & Ziessel, R. (2002). *J. Electroanal. Chem.* **529**, 135–144.
 Chardon-Noblat, S., Deronzier, A. & Ziessel, R. (2001). *Collect. Czech. Chem. Commun.* **66**, 207–227.
 Collomb-Dunand-Sauthier, M.-N., Deronzier, A. & Ziessel, R. (1994). *J. Chem. Soc. Chem. Commun.* pp. 189–191.
 Deacon, G. B., Patrick, J. M., Skelton, B. W., Thomas, N. C. & White, A. H. (1984). *Aust. J. Chem.* **37**, 929–945.
 Gibson, D. H., Sleadd, B. A., Mashuta, M. S. & Richardson, J. F. (1997). *Organometallics*, **16**, 4421–4427.
 Homanen, P., Haukka, M., Pakkanen, T. A., Pursiainen, J. & Laitinen, R. H. (1996). *Organometallics*, **15**, 4081–4084.
 Kooijman, H., Spek, A. L., Mulder, A. & Reinhoudt, D. N. (2007). Private communication (CCDC numbers 666468 and 666615). CCDC, Cambridge, England.
 Kuramochi, Y., Fukaya, K., Yoshida, M. & Ishida, M. (2015). *Chem. Eur. J.* **21**, 10049–10060.
 Luukkanen, S., Homanen, P., Haukka, M., Pakkanen, T. A., Deronzier, A., Chardon-Noblat, S., Zsoldos, D. & Ziessel, R. (1999). *Appl. Catal. Gen.* **185**, 157–164.
 Palatinus, L. & Chapuis, G. (2007). *J. Appl. Cryst.* **40**, 786–790.
 Pettersen, E. F., Goddard, T. D., Huang, C. C., Couch, G. S., Greenblatt, D. M., Meng, E. C. & Ferrin, T. E. (2004). *J. Comput. Chem.* **25**, 1605–1612.
 Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
 Ziessel, R., Grosshenny, V., Hissler, M. & Stroh, C. (2004). *Inorg. Chem.* **43**, 4262–4271.

supporting information

Acta Cryst. (2017). E73, 556-559 [https://doi.org/10.1107/S2056989017003917]

Ruthenium(II) carbonyl compounds with the 4'-chloro-2,2':6',2''-terpyridine ligand

Rajendhraprasad Tatikonda and Matti Haukka

Computing details

For both compounds, data collection: *CrysAlis PRO* (Agilent, 2013); cell refinement: *CrysAlis PRO* (Agilent, 2013); data reduction: *CrysAlis PRO* (Agilent, 2013); program(s) used to solve structure: SUPERFLIP (Palatinus & Chapuis, 2007); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *UCSF Chimera* (Pettersen *et al.*, 2004); software used to prepare material for publication: *SHELXL2014* (Sheldrick, 2015).

(I) *cis*-Dicarbonylchlorido(4'-chloro-2,2':6',2''-terpyridine- κ^3N)ruthenium(II) *fac*-tricarboxyltrichloridoruthenate(II)]

Crystal data

[RuCl(C₁₅H₁₀ClN₃)(CO)₂][RuCl₃(CO)₃]

$M_r = 751.70$

Monoclinic, *P2₁/c*

$a = 14.3578$ (4) Å

$b = 13.9158$ (2) Å

$c = 13.2220$ (3) Å

$\beta = 114.080$ (3)°

$V = 2411.86$ (11) Å³

$Z = 4$

$F(000) = 1456$

$D_x = 2.070$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 6406 reflections

$\theta = 3.5\text{--}28.5^\circ$

$\mu = 1.85$ mm⁻¹

$T = 123$ K

Plate, colourless

$0.34 \times 0.08 \times 0.06$ mm

Data collection

Agilent SuperNova, Dual, Cu at zero, Atlas diffractometer

Radiation source: micro-source

Mirror monochromator

Detector resolution: 10.3953 pixels mm⁻¹

φ scans and ω scans with κ offset

Absorption correction: multi-scan

(*CrysAlisPro*; Agilent, 2013)

$T_{\min} = 0.914$, $T_{\max} = 1.000$

11072 measured reflections

4864 independent reflections

4264 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.023$

$\theta_{\max} = 26.4^\circ$, $\theta_{\min} = 3.1^\circ$

$h = -17 \rightarrow 17$

$k = -16 \rightarrow 17$

$l = -16 \rightarrow 14$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.024$

$wR(F^2) = 0.050$

$S = 1.06$

4864 reflections

316 parameters

0 restraints

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0156P)^2 + 1.2338P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.43$ e Å⁻³

$\Delta\rho_{\min} = -0.48$ e Å⁻³

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ru1	0.17472 (2)	0.27299 (2)	0.26207 (2)	0.01106 (6)
Ru2	0.39629 (2)	0.64980 (2)	0.27157 (2)	0.01345 (6)
Cl1	0.08461 (5)	0.27737 (4)	0.38251 (5)	0.01640 (14)
Cl5	0.37371 (5)	0.70601 (5)	0.08982 (6)	0.02129 (15)
Cl3	0.22742 (5)	0.70409 (5)	0.24405 (6)	0.02024 (15)
Cl2	-0.21948 (5)	0.02482 (5)	-0.03389 (6)	0.02083 (15)
Cl4	0.32731 (5)	0.49515 (5)	0.19478 (6)	0.01848 (15)
N1	0.22418 (16)	0.13257 (15)	0.31413 (17)	0.0124 (5)
O1	0.29403 (15)	0.26878 (13)	0.11931 (16)	0.0217 (4)
N5	0.05630 (16)	0.19549 (15)	0.15594 (17)	0.0122 (5)
C12	-0.0292 (2)	0.24201 (18)	0.0912 (2)	0.0127 (5)
O3	0.42369 (15)	0.57772 (14)	0.49897 (17)	0.0229 (4)
C13	-0.01943 (19)	0.34784 (18)	0.1003 (2)	0.0117 (5)
C18	0.4133 (2)	0.60467 (19)	0.4148 (2)	0.0167 (6)
C7	0.1595 (2)	0.06339 (18)	0.2494 (2)	0.0142 (6)
C2	0.2842 (2)	0.34104 (18)	0.3748 (2)	0.0163 (6)
C8	0.0622 (2)	0.09880 (18)	0.1648 (2)	0.0123 (5)
C20	0.4498 (2)	0.7734 (2)	0.3261 (2)	0.0186 (6)
N15	0.07324 (16)	0.38173 (15)	0.17428 (17)	0.0123 (5)
O4	0.59817 (15)	0.57097 (14)	0.28775 (17)	0.0253 (5)
O5	0.48190 (16)	0.84698 (14)	0.35804 (19)	0.0297 (5)
C19	0.5239 (2)	0.60141 (19)	0.2846 (2)	0.0173 (6)
C10	-0.1111 (2)	0.09151 (19)	0.0372 (2)	0.0141 (6)
O2	0.34998 (15)	0.37495 (13)	0.44648 (16)	0.0227 (4)
C17	0.0862 (2)	0.47673 (18)	0.1892 (2)	0.0140 (6)
H17	0.1505	0.5006	0.2394	0.017*
C9	-0.0227 (2)	0.04364 (19)	0.1039 (2)	0.0136 (6)
H9	-0.0203	-0.0245	0.1077	0.016*
C11	-0.1165 (2)	0.19109 (19)	0.0286 (2)	0.0154 (6)
H11	-0.1775	0.2228	-0.0183	0.018*
C14	-0.0985 (2)	0.40961 (18)	0.0409 (2)	0.0148 (6)
H14	-0.1616	0.3852	-0.0113	0.018*
C3	0.3098 (2)	0.10532 (19)	0.3987 (2)	0.0154 (6)
H3	0.3549	0.1534	0.4432	0.018*
C16	0.0084 (2)	0.54168 (18)	0.1334 (2)	0.0145 (6)
H16	0.0191	0.6087	0.1468	0.017*
C4	0.3355 (2)	0.00955 (19)	0.4243 (2)	0.0181 (6)
H4	0.3956	-0.0077	0.4868	0.022*
C1	0.2473 (2)	0.27202 (18)	0.1705 (2)	0.0157 (6)

C5	0.2717 (2)	-0.06013 (19)	0.3569 (2)	0.0175 (6)
H5	0.2885	-0.1262	0.3713	0.021*
C15	-0.0840 (2)	0.50795 (19)	0.0589 (2)	0.0154 (6)
H15	-0.1377	0.5515	0.0199	0.019*
C6	0.1835 (2)	-0.03308 (19)	0.2687 (2)	0.0163 (6)
H6	0.1394	-0.0804	0.2215	0.020*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ru1	0.01021 (11)	0.00932 (11)	0.01166 (11)	-0.00047 (8)	0.00241 (9)	-0.00009 (9)
Ru2	0.01030 (12)	0.01230 (11)	0.01481 (11)	-0.00009 (8)	0.00210 (9)	-0.00140 (9)
Cl1	0.0184 (4)	0.0147 (3)	0.0170 (3)	-0.0021 (3)	0.0081 (3)	-0.0012 (3)
Cl5	0.0233 (4)	0.0193 (4)	0.0174 (3)	-0.0003 (3)	0.0043 (3)	0.0022 (3)
Cl3	0.0127 (3)	0.0181 (3)	0.0262 (4)	0.0016 (3)	0.0042 (3)	-0.0053 (3)
Cl2	0.0168 (4)	0.0221 (4)	0.0218 (4)	-0.0090 (3)	0.0060 (3)	-0.0057 (3)
Cl4	0.0188 (4)	0.0130 (3)	0.0227 (4)	-0.0025 (3)	0.0074 (3)	-0.0034 (3)
N1	0.0127 (12)	0.0125 (11)	0.0121 (11)	-0.0003 (9)	0.0053 (10)	0.0011 (9)
O1	0.0216 (11)	0.0229 (11)	0.0232 (11)	-0.0031 (9)	0.0119 (10)	-0.0026 (9)
N5	0.0122 (12)	0.0122 (11)	0.0115 (11)	-0.0017 (9)	0.0041 (10)	-0.0005 (9)
C12	0.0126 (14)	0.0155 (13)	0.0106 (13)	0.0004 (11)	0.0055 (11)	-0.0008 (11)
O3	0.0250 (12)	0.0238 (11)	0.0223 (11)	0.0017 (9)	0.0122 (10)	0.0021 (9)
C13	0.0126 (14)	0.0135 (13)	0.0109 (13)	-0.0003 (11)	0.0067 (11)	0.0011 (11)
C18	0.0102 (14)	0.0137 (14)	0.0254 (16)	-0.0001 (11)	0.0063 (12)	-0.0041 (13)
C7	0.0162 (15)	0.0163 (14)	0.0115 (13)	0.0008 (11)	0.0070 (12)	-0.0027 (11)
C2	0.0181 (16)	0.0127 (13)	0.0204 (15)	0.0029 (12)	0.0102 (13)	0.0026 (12)
C8	0.0171 (15)	0.0126 (13)	0.0108 (13)	-0.0003 (11)	0.0093 (12)	-0.0009 (11)
C20	0.0129 (15)	0.0211 (15)	0.0205 (15)	0.0034 (12)	0.0053 (12)	0.0005 (13)
N15	0.0122 (12)	0.0125 (11)	0.0129 (11)	0.0012 (9)	0.0059 (10)	0.0012 (9)
O4	0.0183 (12)	0.0298 (12)	0.0271 (12)	0.0071 (9)	0.0086 (10)	0.0017 (10)
O5	0.0236 (12)	0.0206 (11)	0.0428 (14)	-0.0060 (9)	0.0114 (11)	-0.0129 (10)
C19	0.0196 (16)	0.0160 (14)	0.0130 (13)	-0.0023 (12)	0.0031 (12)	0.0002 (11)
C10	0.0148 (14)	0.0187 (14)	0.0102 (13)	-0.0053 (11)	0.0065 (12)	-0.0024 (11)
O2	0.0194 (11)	0.0216 (10)	0.0205 (11)	-0.0064 (9)	0.0013 (10)	-0.0051 (9)
C17	0.0148 (14)	0.0142 (13)	0.0135 (13)	-0.0024 (11)	0.0064 (12)	-0.0013 (11)
C9	0.0195 (15)	0.0123 (13)	0.0113 (13)	-0.0047 (11)	0.0087 (12)	-0.0030 (11)
C11	0.0133 (14)	0.0192 (14)	0.0113 (13)	-0.0004 (11)	0.0026 (12)	0.0016 (11)
C14	0.0148 (14)	0.0174 (14)	0.0139 (13)	-0.0012 (11)	0.0076 (12)	0.0020 (11)
C3	0.0151 (15)	0.0158 (14)	0.0153 (13)	0.0001 (11)	0.0062 (12)	0.0003 (11)
C16	0.0222 (16)	0.0095 (12)	0.0181 (14)	0.0021 (11)	0.0146 (13)	0.0009 (11)
C4	0.0172 (15)	0.0204 (15)	0.0163 (14)	0.0062 (12)	0.0063 (12)	0.0070 (12)
C1	0.0154 (15)	0.0108 (13)	0.0158 (14)	-0.0017 (11)	0.0011 (12)	-0.0001 (11)
C5	0.0234 (16)	0.0118 (13)	0.0217 (15)	0.0031 (12)	0.0135 (13)	0.0018 (12)
C15	0.0158 (15)	0.0184 (14)	0.0145 (13)	0.0074 (12)	0.0087 (12)	0.0052 (12)
C6	0.0197 (16)	0.0142 (13)	0.0184 (14)	-0.0011 (11)	0.0113 (13)	-0.0015 (12)

Geometric parameters (Å, °)

Ru1—C1	1.893 (3)	C7—C8	1.473 (4)
Ru1—C2	1.918 (3)	C2—O2	1.133 (3)
Ru1—N5	2.019 (2)	C8—C9	1.387 (4)
Ru1—N15	2.093 (2)	C20—O5	1.131 (3)
Ru1—N1	2.097 (2)	N15—C17	1.338 (3)
Ru1—C11	2.4279 (7)	O4—C19	1.132 (3)
Ru2—C19	1.893 (3)	C10—C9	1.386 (4)
Ru2—C20	1.902 (3)	C10—C11	1.390 (4)
Ru2—C18	1.914 (3)	C17—C16	1.392 (4)
Ru2—C14	2.4129 (7)	C17—H17	0.9500
Ru2—C15	2.4199 (7)	C9—H9	0.9500
Ru2—C13	2.4212 (7)	C11—H11	0.9500
C12—C10	1.724 (3)	C14—C15	1.390 (4)
N1—C3	1.336 (3)	C14—H14	0.9500
N1—C7	1.369 (3)	C3—C4	1.387 (4)
O1—C1	1.131 (3)	C3—H3	0.9500
N5—C12	1.342 (3)	C16—C15	1.373 (4)
N5—C8	1.350 (3)	C16—H16	0.9500
C12—C11	1.383 (4)	C4—C5	1.381 (4)
C12—C13	1.480 (3)	C4—H4	0.9500
O3—C18	1.125 (3)	C5—C6	1.379 (4)
C13—N15	1.374 (3)	C5—H5	0.9500
C13—C14	1.385 (4)	C15—H15	0.9500
C7—C6	1.384 (4)	C6—H6	0.9500
C1—Ru1—C2	90.60 (11)	C6—C7—C8	123.5 (2)
C1—Ru1—N5	94.52 (10)	O2—C2—Ru1	174.6 (2)
C2—Ru1—N5	174.14 (10)	N5—C8—C9	119.5 (2)
C1—Ru1—N15	95.21 (10)	N5—C8—C7	114.0 (2)
C2—Ru1—N15	103.79 (10)	C9—C8—C7	126.3 (2)
N5—Ru1—N15	78.61 (8)	O5—C20—Ru2	179.5 (3)
C1—Ru1—N1	90.13 (10)	C17—N15—C13	118.7 (2)
C2—Ru1—N1	98.32 (10)	C17—N15—Ru1	127.51 (18)
N5—Ru1—N1	78.86 (8)	C13—N15—Ru1	113.54 (16)
N15—Ru1—N1	157.17 (8)	O4—C19—Ru2	177.0 (2)
C1—Ru1—C11	178.53 (8)	C9—C10—C11	122.5 (2)
C2—Ru1—C11	87.98 (8)	C9—C10—C12	118.5 (2)
N5—Ru1—C11	86.92 (6)	C11—C10—C12	119.0 (2)
N15—Ru1—C11	84.78 (6)	N15—C17—C16	122.0 (3)
N1—Ru1—C11	90.45 (6)	N15—C17—H17	119.0
C19—Ru2—C20	93.51 (11)	C16—C17—H17	119.0
C19—Ru2—C18	93.76 (11)	C10—C9—C8	117.6 (2)
C20—Ru2—C18	93.03 (11)	C10—C9—H9	121.2
C19—Ru2—C14	86.39 (8)	C8—C9—H9	121.2
C20—Ru2—C14	177.64 (9)	C12—C11—C10	117.1 (2)
C18—Ru2—C14	89.34 (8)	C12—C11—H11	121.5

C19—Ru2—C15	86.44 (8)	C10—C11—H11	121.5
C20—Ru2—C15	87.36 (9)	C13—C14—C15	118.9 (3)
C18—Ru2—C15	179.55 (9)	C13—C14—H14	120.6
C14—Ru2—C15	90.27 (2)	C15—C14—H14	120.6
C19—Ru2—C13	175.98 (8)	N1—C3—C4	122.6 (3)
C20—Ru2—C13	89.96 (8)	N1—C3—H3	118.7
C18—Ru2—C13	88.08 (8)	C4—C3—H3	118.7
C14—Ru2—C13	90.06 (2)	C15—C16—C17	119.3 (2)
C15—Ru2—C13	91.70 (3)	C15—C16—H16	120.3
C3—N1—C7	118.8 (2)	C17—C16—H16	120.3
C3—N1—Ru1	127.74 (18)	C5—C4—C3	118.5 (3)
C7—N1—Ru1	113.43 (17)	C5—C4—H4	120.8
C12—N5—C8	123.0 (2)	C3—C4—H4	120.8
C12—N5—Ru1	118.52 (17)	O1—C1—Ru1	176.8 (2)
C8—N5—Ru1	117.81 (17)	C6—C5—C4	119.5 (2)
N5—C12—C11	120.3 (2)	C6—C5—H5	120.2
N5—C12—C13	113.3 (2)	C4—C5—H5	120.2
C11—C12—C13	126.3 (2)	C16—C15—C14	119.5 (2)
N15—C13—C14	121.5 (2)	C16—C15—H15	120.2
N15—C13—C12	115.6 (2)	C14—C15—H15	120.2
C14—C13—C12	122.9 (2)	C5—C6—C7	119.7 (3)
O3—C18—Ru2	179.5 (3)	C5—C6—H6	120.2
N1—C7—C6	120.8 (2)	C7—C6—H6	120.2
N1—C7—C8	115.5 (2)		
C8—N5—C12—C11	-0.7 (4)	C12—C13—N15—Ru1	-3.3 (3)
Ru1—N5—C12—C11	-171.18 (18)	C13—N15—C17—C16	1.0 (4)
C8—N5—C12—C13	176.3 (2)	Ru1—N15—C17—C16	-172.89 (18)
Ru1—N5—C12—C13	5.8 (3)	C11—C10—C9—C8	1.0 (4)
N5—C12—C13—N15	-1.4 (3)	C12—C10—C9—C8	-176.72 (18)
C11—C12—C13—N15	175.3 (2)	N5—C8—C9—C10	-0.9 (3)
N5—C12—C13—C14	-179.9 (2)	C7—C8—C9—C10	172.8 (2)
C11—C12—C13—C14	-3.2 (4)	N5—C12—C11—C10	0.6 (4)
C3—N1—C7—C6	-2.0 (4)	C13—C12—C11—C10	-175.9 (2)
Ru1—N1—C7—C6	176.53 (18)	C9—C10—C11—C12	-0.8 (4)
C3—N1—C7—C8	174.3 (2)	C12—C10—C11—C12	176.86 (19)
Ru1—N1—C7—C8	-7.2 (3)	N15—C13—C14—C15	-1.6 (4)
C12—N5—C8—C9	0.8 (4)	C12—C13—C14—C15	176.8 (2)
Ru1—N5—C8—C9	171.39 (17)	C7—N1—C3—C4	-0.6 (4)
C12—N5—C8—C7	-173.6 (2)	Ru1—N1—C3—C4	-178.83 (18)
Ru1—N5—C8—C7	-3.1 (3)	N15—C17—C16—C15	-1.5 (4)
N1—C7—C8—N5	6.9 (3)	N1—C3—C4—C5	2.5 (4)
C6—C7—C8—N5	-177.0 (2)	C3—C4—C5—C6	-1.8 (4)
N1—C7—C8—C9	-167.1 (2)	C17—C16—C15—C14	0.4 (4)
C6—C7—C8—C9	9.0 (4)	C13—C14—C15—C16	1.1 (4)
C14—C13—N15—C17	0.6 (4)	C4—C5—C6—C7	-0.6 (4)
C12—C13—N15—C17	-177.9 (2)	N1—C7—C6—C5	2.6 (4)
C14—C13—N15—Ru1	175.26 (18)	C8—C7—C6—C5	-173.4 (2)

Hydrogen-bond geometry (Å, °)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C11—H11 \cdots C15 ⁱ	0.95	2.76	3.664 (3)	158
C16—H16 \cdots C11 ⁱⁱ	0.95	2.72	3.515 (3)	142
C5—H5 \cdots Cl3 ⁱⁱⁱ	0.95	2.82	3.553 (3)	134

Symmetry codes: (i) $-x, -y+1, -z$; (ii) $-x, y+1/2, -z+1/2$; (iii) $x, y-1, z$.

(II) *cis*-Dicarbonyl-*trans*-dichlorido(4'-chloro-2,2':6',2''-terpyridine- $\kappa^2 N^1, N^1$)ruthenium(II)

Crystal data

[RuCl₂(C₁₅H₁₀ClN₃(CO)₂)₂]

$M_r = 495.70$

Triclinic, $P\bar{1}$

$a = 7.3019$ (3) Å

$b = 8.5080$ (3) Å

$c = 14.7702$ (6) Å

$\alpha = 101.287$ (3)°

$\beta = 91.835$ (3)°

$\gamma = 98.144$ (3)°

$V = 889.09$ (6) Å³

$Z = 2$

$F(000) = 488$

$D_x = 1.852$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 5231 reflections

$\theta = 5.4-76.2^\circ$

$\mu = 1.35$ mm⁻¹

$T = 123$ K

Plate, brown

$0.30 \times 0.08 \times 0.05$ mm

Data collection

Agilent SuperNova, Dual, Cu at zero, Atlas diffractometer

Radiation source: micro-source

Mirror monochromator

Detector resolution: 10.3953 pixels mm⁻¹

φ scans and ω scans with κ offset

Absorption correction: multi-scan

(*CrysAlisPro*; Agilent, 2013)

$T_{\min} = 0.300$, $T_{\max} = 1.000$

7508 measured reflections

3662 independent reflections

3405 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.036$

$\theta_{\max} = 26.6^\circ$, $\theta_{\min} = 1.4^\circ$

$h = -8 \rightarrow 9$

$k = -7 \rightarrow 10$

$l = -18 \rightarrow 18$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.033$

$wR(F^2) = 0.088$

$S = 1.07$

3662 reflections

235 parameters

0 restraints

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0453P)^2 + 0.5543P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.74$ e Å⁻³

$\Delta\rho_{\min} = -1.43$ e Å⁻³

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Ru1	0.97989 (3)	0.30538 (2)	0.20165 (2)	0.01402 (9)

C12	1.19336 (10)	0.26275 (9)	0.31916 (5)	0.02134 (16)
C13	0.53931 (11)	0.29015 (9)	0.58174 (5)	0.02370 (16)
C11	0.76545 (11)	0.37329 (9)	0.09657 (5)	0.02563 (17)
O1	0.9433 (4)	-0.0448 (3)	0.10113 (17)	0.0312 (5)
O2	1.2842 (4)	0.3556 (3)	0.07324 (19)	0.0380 (6)
N2	0.7827 (3)	0.2976 (3)	0.30696 (17)	0.0187 (5)
N1	1.0061 (3)	0.5492 (3)	0.27105 (17)	0.0183 (5)
N3	0.6478 (4)	-0.1192 (3)	0.26366 (18)	0.0226 (5)
C2	1.1703 (5)	0.3368 (4)	0.1217 (2)	0.0242 (6)
C13	0.6151 (4)	0.0228 (4)	0.2446 (2)	0.0201 (6)
C12	0.6700 (4)	0.1661 (4)	0.3213 (2)	0.0190 (6)
C10	0.6327 (4)	0.2955 (4)	0.4765 (2)	0.0197 (6)
C16	0.5048 (5)	-0.2483 (4)	0.1144 (2)	0.0280 (7)
H16	0.4700	-0.3452	0.0693	0.034*
C9	0.7419 (4)	0.4346 (4)	0.4621 (2)	0.0204 (6)
H9	0.7662	0.5293	0.5096	0.024*
C11	0.5945 (4)	0.1598 (4)	0.4060 (2)	0.0208 (6)
H11	0.5186	0.0646	0.4152	0.025*
C7	0.9276 (4)	0.5750 (3)	0.3533 (2)	0.0190 (6)
C8	0.8142 (4)	0.4312 (3)	0.3764 (2)	0.0176 (5)
C17	0.5949 (5)	-0.2517 (4)	0.1970 (2)	0.0271 (7)
H17	0.6211	-0.3531	0.2076	0.032*
C1	0.9506 (4)	0.0845 (4)	0.1412 (2)	0.0226 (6)
C14	0.5230 (4)	0.0391 (4)	0.1639 (2)	0.0236 (6)
H14	0.5002	0.1420	0.1543	0.028*
C6	0.9447 (4)	0.7303 (4)	0.4081 (2)	0.0234 (6)
H6	0.8917	0.7472	0.4665	0.028*
C15	0.4656 (5)	-0.1013 (4)	0.0977 (2)	0.0281 (7)
H15	0.4004	-0.0966	0.0418	0.034*
C3	1.0952 (4)	0.6760 (4)	0.2401 (2)	0.0235 (6)
H3	1.1461	0.6571	0.1812	0.028*
C5	1.0398 (5)	0.8593 (4)	0.3764 (2)	0.0261 (6)
H5	1.0533	0.9657	0.4132	0.031*
C4	1.1151 (4)	0.8330 (4)	0.2910 (2)	0.0248 (6)
H4	1.1790	0.9206	0.2678	0.030*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ru1	0.01375 (13)	0.01423 (13)	0.01338 (13)	0.00090 (9)	0.00123 (8)	0.00191 (9)
C12	0.0197 (3)	0.0204 (3)	0.0232 (3)	0.0047 (3)	-0.0033 (3)	0.0023 (3)
C13	0.0266 (4)	0.0286 (4)	0.0179 (3)	0.0064 (3)	0.0063 (3)	0.0069 (3)
C11	0.0278 (4)	0.0256 (4)	0.0223 (4)	0.0028 (3)	-0.0067 (3)	0.0043 (3)
O1	0.0374 (14)	0.0215 (11)	0.0310 (13)	0.0025 (10)	0.0078 (10)	-0.0031 (10)
O2	0.0325 (14)	0.0397 (14)	0.0417 (15)	0.0039 (11)	0.0167 (12)	0.0068 (12)
N2	0.0167 (11)	0.0194 (12)	0.0205 (12)	0.0036 (9)	0.0012 (9)	0.0043 (9)
N1	0.0172 (12)	0.0179 (11)	0.0194 (12)	0.0015 (9)	-0.0012 (9)	0.0042 (9)
N3	0.0228 (13)	0.0226 (12)	0.0224 (13)	0.0033 (10)	0.0023 (10)	0.0044 (10)

C2	0.0289 (16)	0.0192 (14)	0.0238 (15)	0.0035 (12)	0.0008 (12)	0.0030 (12)
C13	0.0175 (13)	0.0229 (14)	0.0191 (14)	-0.0009 (11)	0.0027 (11)	0.0044 (11)
C12	0.0177 (13)	0.0197 (13)	0.0191 (14)	0.0010 (11)	-0.0004 (11)	0.0043 (11)
C10	0.0205 (14)	0.0256 (14)	0.0141 (13)	0.0069 (11)	0.0014 (10)	0.0043 (11)
C16	0.0315 (17)	0.0237 (15)	0.0224 (15)	-0.0075 (13)	0.0072 (13)	-0.0033 (12)
C9	0.0210 (14)	0.0209 (14)	0.0193 (14)	0.0059 (11)	-0.0018 (11)	0.0028 (11)
C11	0.0188 (14)	0.0207 (14)	0.0225 (14)	0.0012 (11)	-0.0001 (11)	0.0050 (11)
C7	0.0164 (13)	0.0193 (13)	0.0209 (14)	0.0026 (11)	-0.0003 (11)	0.0033 (11)
C8	0.0159 (13)	0.0180 (13)	0.0172 (13)	0.0027 (10)	-0.0032 (10)	0.0004 (10)
C17	0.0279 (16)	0.0234 (15)	0.0296 (17)	0.0032 (13)	0.0070 (13)	0.0043 (13)
C1	0.0208 (14)	0.0271 (16)	0.0208 (14)	0.0037 (12)	0.0052 (11)	0.0069 (12)
C14	0.0248 (15)	0.0238 (14)	0.0207 (15)	-0.0008 (12)	-0.0013 (12)	0.0049 (12)
C6	0.0237 (15)	0.0224 (15)	0.0227 (15)	0.0030 (12)	0.0006 (12)	0.0017 (12)
C15	0.0309 (17)	0.0303 (16)	0.0195 (15)	-0.0057 (14)	-0.0022 (12)	0.0040 (12)
C3	0.0204 (14)	0.0259 (15)	0.0246 (15)	0.0018 (12)	0.0004 (12)	0.0074 (12)
C5	0.0246 (15)	0.0210 (15)	0.0305 (17)	0.0022 (12)	-0.0014 (13)	0.0014 (12)
C4	0.0211 (14)	0.0201 (14)	0.0329 (17)	-0.0008 (12)	-0.0015 (12)	0.0078 (12)

Geometric parameters (Å, °)

Ru1—C2	1.877 (3)	C16—C17	1.376 (5)
Ru1—C1	1.895 (3)	C16—C15	1.387 (5)
Ru1—N1	2.105 (2)	C16—H16	0.9500
Ru1—N2	2.157 (2)	C9—C8	1.383 (4)
Ru1—C11	2.3762 (8)	C9—H9	0.9500
Ru1—C12	2.4098 (7)	C11—H11	0.9500
C13—C10	1.723 (3)	C7—C6	1.395 (4)
O1—C1	1.135 (4)	C7—C8	1.481 (4)
O2—C2	1.129 (4)	C17—H17	0.9500
N2—C12	1.348 (4)	C14—C15	1.390 (4)
N2—C8	1.361 (4)	C14—H14	0.9500
N1—C3	1.345 (4)	C6—C5	1.384 (4)
N1—C7	1.352 (4)	C6—H6	0.9500
N3—C13	1.344 (4)	C15—H15	0.9500
N3—C17	1.344 (4)	C3—C4	1.384 (4)
C13—C14	1.391 (4)	C3—H3	0.9500
C13—C12	1.490 (4)	C5—C4	1.383 (5)
C12—C11	1.391 (4)	C5—H5	0.9500
C10—C11	1.384 (4)	C4—H4	0.9500
C10—C9	1.387 (4)		
C2—Ru1—C1	85.52 (13)	C8—C9—C10	117.9 (3)
C2—Ru1—N1	96.08 (11)	C8—C9—H9	121.0
C1—Ru1—N1	178.40 (10)	C10—C9—H9	121.0
C2—Ru1—N2	171.98 (12)	C10—C11—C12	118.4 (3)
C1—Ru1—N2	101.47 (11)	C10—C11—H11	120.8
N1—Ru1—N2	76.94 (10)	C12—C11—H11	120.8
C2—Ru1—C11	90.26 (10)	N1—C7—C6	120.8 (3)

C1—Ru1—C11	93.67 (10)	N1—C7—C8	115.5 (3)
N1—Ru1—C11	86.24 (7)	C6—C7—C8	123.5 (3)
N2—Ru1—C11	93.17 (7)	N2—C8—C9	122.6 (3)
C2—Ru1—C12	92.02 (10)	N2—C8—C7	115.3 (3)
C1—Ru1—C12	92.10 (10)	C9—C8—C7	122.0 (3)
N1—Ru1—C12	87.93 (7)	N3—C17—C16	123.4 (3)
N2—Ru1—C12	83.87 (7)	N3—C17—H17	118.3
C11—Ru1—C12	173.94 (3)	C16—C17—H17	118.3
C12—N2—C8	118.4 (3)	O1—C1—Ru1	175.0 (3)
C12—N2—Ru1	126.9 (2)	C15—C14—C13	117.3 (3)
C8—N2—Ru1	112.57 (19)	C15—C14—H14	121.3
C3—N1—C7	119.5 (3)	C13—C14—H14	121.3
C3—N1—Ru1	125.0 (2)	C5—C6—C7	119.1 (3)
C7—N1—Ru1	115.54 (19)	C5—C6—H6	120.4
C13—N3—C17	116.5 (3)	C7—C6—H6	120.4
O2—C2—Ru1	179.6 (3)	C16—C15—C14	119.1 (3)
N3—C13—C14	124.5 (3)	C16—C15—H15	120.5
N3—C13—C12	114.9 (3)	C14—C15—H15	120.5
C14—C13—C12	120.4 (3)	N1—C3—C4	122.3 (3)
N2—C12—C11	122.1 (3)	N1—C3—H3	118.9
N2—C12—C13	120.7 (3)	C4—C3—H3	118.9
C11—C12—C13	117.1 (3)	C4—C5—C6	119.8 (3)
C11—C10—C9	120.4 (3)	C4—C5—H5	120.1
C11—C10—C13	119.0 (2)	C6—C5—H5	120.1
C9—C10—C13	120.6 (2)	C5—C4—C3	118.4 (3)
C17—C16—C15	119.1 (3)	C5—C4—H4	120.8
C17—C16—H16	120.5	C3—C4—H4	120.8
C15—C16—H16	120.5		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C9—H9 \cdots C12 ⁱ	0.95	2.77	3.687 (3)	163

Symmetry code: (i) $-x+2, -y+1, -z+1$.