



Two closely related 2-(benzofuran-2-yl)-2-oxoethyl benzoates: structural differences and C—H...O hydrogen-bonded supramolecular assemblies

Li Yee Then,^a C. S. Chidan Kumar,^{b*} Huey Chong Kwong,^c Yip-Foo Win,^d Siau Hui Mah,^e Ching Kheng Quah,^a S. Naveen^f and Ismail Warad^{g*}

Received 22 June 2017

Accepted 23 June 2017

Edited by P. McArdle, National University of Ireland, Ireland

Keywords: crystal structure; benzofuran; intramolecular interaction; intermolecular interaction; functional group.

CCDC references: 1449589; 1449587


Supporting information: this article has supporting information at journals.iucr.org/e


^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, ^bDepartment of Engineering Chemistry, Vidya Vikas Institute of Engineering & Technology, Visvesvaraya Technological University, Alanahally, Mysuru 570 028, Karnataka, India, ^cSchool of Chemical Sciences, Universiti Sains Malaysia, Penang 11800 USM, Malaysia, ^dDepartment of Chemical Science, Faculty of Science, Universiti Tunku Abdul Rahman, Perak Campus, Jalan Universiti, Bandar Barat, Perak, Malaysia, ^eSchool of Biosciences, Taylor's University, Lakeside Campus, 47500 Subang Jaya, Selangor, Malaysia, ^fInstitution of Excellence, University of Mysore, Manasagangotri, Mysuru 570 006, India, and ^gDepartment of Chemistry, Science College, An-Najah National University, PO Box 7, Nablus, West Bank, Palestinian Territories. *Correspondence e-mail: chidankumar@gmail.com, khalil.i@najah.edu

The compounds 2-(1-benzofuran-2-yl)-2-oxoethyl 2-nitrobenzoate, C₁₇H₁₁NO₆ (I), and 2-(1-benzofuran-2-yl)-2-oxoethyl 2-aminobenzoate, C₁₇H₁₃NO₄ (II), were synthesized under mild conditions. Their molecular structures were characterized by both spectroscopic and single-crystal X-ray diffraction analysis. The molecular conformations of both title compounds are generally similar. However, different *ortho*-substituted moieties at the phenyl ring of the two compounds cause deviations in the torsion angles between the carbonyl group and the attached phenyl ring. In compound (I), the *ortho*-nitrophenyl ring is twisted away from the adjacent carbonyl group whereas in compound (II), the *ortho*-aminophenyl ring is almost co-planar with the carbonyl group. In the crystal of compound (I), two C—H...O hydrogen bonds link the molecules into chains propagating along the *c*-axis direction and the chains are interdigitated, forming sheets parallel to [20 $\bar{1}$]. Conversely, pairs of N—H...O hydrogen bonds in compound (II) link inversion-related molecules into dimers, which are further extended by C—H...O hydrogen bonds into dimer chains. These chains are interconnected by π – π interactions involving the furan rings, forming sheets parallel to the *ac* plane.

1. Chemical context

Oxygen-containing heterocycles are the basic cores of many bioactive structures. Among these, benzofuran and its derivatives occur frequently in nature because of their stability and ease of generation. Those with substitution(s) at their C-2 and/or C-3 positions are important. Important biological activity such as anticancer (Swamy *et al.*, 2015), anti-acetylcholinesterase (Zhou *et al.*, 2010), antimicrobial (Ugale *et al.*, 2012) and antioxidant (Naik *et al.*, 2013) actions exhibited by this scaffold have attracted the attention of synthetic chemists. Some of the biological and medicinal significance of benzofuran derivatives (Nevagi *et al.*, 2015) have been discussed in review reports. The known potential of benzofuran derivatives has motivated us to synthesise some new compounds incorporating this core structure and we herein report the synthesis and crystal structures of 2-(1-benzofuran-2-yl)-2-oxoethyl 2-nitrobenzoate (I) and 2-(1-benzofuran-2-yl)-2-oxoethyl 2-aminobenzoate (II).





2. Structural commentary

The molecular structures of the title compounds (Fig. 1) contain a benzofuran ring and an *ortho*-substituted [nitro- for compound (I) and amino- for compound (II)] phenyl ring, joined by a C–C(=O)–O–C(=O) carbonyl-connecting bridge. Their molecular conformations can be characterized by three degrees of freedom, as indicated by the O1–C8–C9–O3 (τ_1), C9–C10–O2–C11 (τ_2) and O4–C11–C12–C13 (τ_3) torsion angles, respectively (Fig. 2). The torsion angle τ_1 for compounds (I) and (II) is close to 0° , showing that the benzofuran ring is nearly coplanar with the C–C(=O)–O–




Figure 1
ORTEP diagram of the title compounds, with ellipsoids drawn at the 50% probability level, showing the atomic labelling scheme.

Table 1
Hydrogen-bond geometry (\AA , $^\circ$) for (I).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$C10-H10A\cdots O3^i$	0.99	2.59	3.471 (4)	148
$C15-H15A\cdots O5^{ii}$	0.95	2.58	3.380 (3)	142

Symmetry codes: (i) $x, y, z - 1$; (ii) $x - \frac{1}{2}, -y + \frac{1}{2}, z + 1$.

Table 2
Hydrogen-bond geometry (\AA , $^\circ$) for (II).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$N1-H1A\cdots O4$	0.91 (2)	2.05 (2)	2.700 (3)	127.7 (18)
$N1-H1A\cdots O4^i$	0.91 (2)	2.49 (2)	3.246 (2)	141.4 (18)
$C10-H10A\cdots O3^{ii}$	0.97	2.50	3.444 (2)	165

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $x - 1, y, z$.

C(=O) carbonyl bridge. Torsion angle τ_2 adopts a *syn*-clinal conformation, as both carbonyl groups at the connecting bridges are twisted away from each other forming torsion angles of $-71.43 (3)^\circ$ in (I) and $-70.85 (18)^\circ$ in (II). For compound (I), the substituted *ortho*-nitrophenyl moiety is perpendicular to the adjacent carbonyl group with a τ_3 torsion angle of $-90.2 (4)^\circ$; this may arise from a steric repulsion force between the nitro group and carbonyl group. In contrast, the *ortho*-aminophenyl ring in compound (II) is almost coplanar with its adjacent carbonyl group due to the intramolecular hydrogen bond ($N1-H1A\cdots O4$, Table 2) between the amino and carbonyl groups, which generates an $S(6)$ ring.

3. Supramolecular features

The crystal packing of compound (I) depends mainly on two weak intermolecular hydrogen bonds. Molecules are joined into infinite chains propagating along the c -axis by $C10-H10A\cdots O3$ hydrogen bonds (Table 1, Fig. 3), meanwhile those chains are interdigitated into a fishbone sheet extending along the $[20\bar{1}]$ direction through $C15-H15A\cdots O5$ hydrogen bonds. The fishbone sheets alternate in an up-down manner along the ab plane as shown in Fig. 4.

In compound (II), the molecular interactions are more abundant than in (I) because of the *ortho*-substituted amino




Figure 2
General chemical diagram showing torsion angles τ_1 , τ_2 and τ_3 in compounds (I) and (II).




Figure 3
Molecules in compound (I) joined by intermolecular hydrogen bonds, forming a fishbone chain.

group at its phenyl ring. Pairs of $N1-H1A \cdots O4$ hydrogen bonds link molecules into inversion dimers with an $R_2^2(12)$ graph-set motif (Fig. 5). These dimers are further expanded by $C10-H10A \cdots O3$ hydrogen bonds into infinite chains along the [100] direction (Fig. 6). In addition, neighbouring chains are interconnected by $\pi-\pi$ interactions involving adjacent furan rings [centroid-centroid distance = 3.7982 (15) Å; symmetry code: $-x, -y + 1, -z$], forming a sheet parallel to the ac plane (Fig. 7).

4. Database survey

A survey of the Cambridge Structural Database (Groom *et al.*, 2016) revealed five benzofuran structures (Kumar *et al.*, 2015) similar to the title compounds: ITAXUY, ITAYAF, ITAYEJ, ITAYIN and ITAYOT. The molecular structures of the studied and previous compounds differ only at their substituted phenyl rings. By comparing their torsion angles at the




Figure 4
Fishbone chains in an up-down manner are shown in different colours.




Figure 5
Intramolecular and intermolecular $N1-H1A \cdots O4$ hydrogen bonds.

$C(=O)-O-C(=O)$ carbonyl bridges, the title compounds exhibit a *syn-clinal* conformation similar to ITAXUY, ITAYEJ and ITAYIN with respect to their torsion angles which range from 75 to 80°.




Figure 6
Interactions in the crystal structure of compound (II), showing hydrogen bonds (cyan dotted lines) and $\pi-\pi$ interactions (red dotted lines).




Figure 7
The packing of compound (II), showing the hydrogen bonds (cyan dotted lines) and $\pi-\pi$ interactions (red dotted lines).

Table 3
Experimental details.

	(I)	(II)
Crystal data		
Chemical formula	C ₁₇ H ₁₁ NO ₆	C ₁₇ H ₁₃ NO ₄
<i>M_r</i>	325.27	295.28
Crystal system, space group	Orthorhombic, <i>Pna</i> 2 ₁	Triclinic, <i>P</i> $\bar{1}$
Temperature (K)	100	297
<i>a</i> , <i>b</i> , <i>c</i> (Å)	9.3022 (10), 28.482 (3), 5.5208 (6)	5.1839 (12), 10.853 (3), 12.269 (3)
α , β , γ (°)	90, 90, 90	93.562 (3), 91.167 (3), 98.714 (3)
<i>V</i> (Å ³)	1462.7 (3)	680.6 (3)
<i>Z</i>	4	2
Radiation type	Mo <i>K</i> α	Mo <i>K</i> α
μ (mm ⁻¹)	0.11	0.10
Crystal size (mm)	0.27 × 0.16 × 0.13	0.40 × 0.32 × 0.21
Data collection		
Diffractometer	Bruker APEXII DUO CCD area-detector	Bruker APEXII DUO CCD area-detector
Absorption correction	Multi-scan (<i>SADABS</i> ; Bruker, 2009)	Multi-scan (<i>SADABS</i> ; Bruker, 2009)
<i>T_{min}</i> , <i>T_{max}</i>	0.933, 0.985	0.871, 0.978
No. of measured, independent and observed [<i>I</i> > 2 σ (<i>I</i>)] reflections	15875, 3358, 2915	17052, 3105, 2214
<i>R_{int}</i>	0.037	0.037
(<i>sin</i> θ / λ) _{max} (Å ⁻¹)	0.651	0.650
Refinement		
<i>R</i> [<i>F</i> ² > 2 σ (<i>F</i> ²)], <i>wR</i> (<i>F</i> ²), <i>S</i>	0.038, 0.085, 1.08	0.046, 0.123, 1.08
No. of reflections	3358	3105
No. of parameters	217	207
No. of restraints	1	0
H-atom treatment	H-atom parameters constrained	H atoms treated by a mixture of independent and constrained refinement
$\Delta\rho_{\max}$, $\Delta\rho_{\min}$ (e Å ⁻³)	0.18, -0.17	0.18, -0.18

Computer programs: *APEX2* and *SAINT* (Bruker, 2009), *SHELXT2013* (Sheldrick, 2015a), *SHELXL2013* (Sheldrick, 2015b), *Mercury* (Macrae *et al.*, 2006) and *PLATON* (Spek, 2009).

5. Synthesis and crystallization

The synthesis was carried out by reacting 1-(benzofuran-2-yl)-2-bromoethan-1-one (1 mmol) with 2-nitrobenzoic acid (1 mmol) for compound (I) and 2-aminobenzoic acid (1 mmol) for compound (II) in 8 ml of *N,N*-dimethylformamide in the presence of a catalytic amount of anhydrous potassium carbonate at room temperature. The reaction solution was stirred for about two h and monitored by thin-layer chromatography (TLC). After the reaction was complete, the resultant mixture was then added to a beaker of ice-cooled water to form a precipitate. The precipitate was then filtered, rinsed with distilled water and dried. Crystals suitable for X-ray analysis were obtained by slow evaporation using a suitable solvent.

2-(Benzofuran-2-yl)-2-oxoethyl 2-nitrobenzoate (I):

Solvents used to grow crystal: acetone + methanol 1:1 *v/v*; yield: 80%, m.p. 381–383 K; ¹H NMR (500MHz, CDCl₃) in ppm: δ 8.041–8.025 (*d*, 1H, *J* = 7.9Hz, ¹⁴CH), 7.995–7.980 (*d*, 1H, *J* = 7.9Hz, ¹⁷CH), 7.796–7.763 (*m*, 2H, ²CH, ³CH), 7.726–7.695 (*t*, 1H, *J* = 7.9Hz, ¹⁵CH), 7.673 (*s*, 1H, ⁷CH), 7.644–7.627 (*d*, 1H, *J* = 8.4Hz, ⁵CH), 7.578–7.544 (*t*, 1H, *J* = 8.4Hz, ⁴CH), 7.398–7.366 (*t*, 1H, *J* = 7.9Hz, ¹⁶CH), 5.609 (*s*, 2H, ¹⁰CH₂). ¹³C NMR (125 MHz, CDCl₃) in ppm: 182.94 (C9), 165.67 (C11), 155.80 (C1), 150.25 (C13), 133.31 (C16), 132.04 (C15), 130.39 (C17), 130.10 (C8), 128.97 (C3), 127.23 (C12), 126.70 (C6), 124.34 (C5), 124.11 (C4), 123.60 (C14), 113.75 (C7), 112.57 (C2), 67.10 (C10). FT-IR (ATR (solid) cm⁻¹): 3089 (Ar C–H,

v), 2953 (C–H, *v*), 1744, 1686 (C=O, *v*), 1612 (C=C, *v*), 1554, 1422 (Ar C=C, *v*), 1529, 1344 (N=O, *v*), 1278, 1123 (C–O, *v*).

2-(Benzofuran-2-yl)-2-oxoethyl 2-aminobenzoate (II):

Solvents used to grow crystal: acetone + acetonitrile (1:1 *v/v*); yield: 83%; m.p. 432–434 K; ¹H NMR (500 MHz, DMSO) in ppm: δ 8.083 (*s*, 1H, ⁷CH), 7.907–7.891 (*d*, 1H, *J* = 8.1Hz, ¹⁷CH), 7.848–7.832 (*d*, 1H, *J* = 8.1Hz, ¹⁴CH), 7.787–7.770 (*d*, 1H, *J* = 8.5Hz, ²CH), 7.617–7.583 (*t*, 1H, *J* = 8.5Hz, ³CH), 7.437–7.405 (*t*, 1H, *J* = 8.1Hz, ¹⁵CH), 7.329–7.295 (*t*, 1H, *J* = 8.5Hz, ⁴CH), 6.824–6.807 (*d*, 1H, *J* = 8.5Hz, ⁵CH), 6.669 (*br-s*, 2H, ¹NH₂), 6.607–6.574 (*t*, 1H, *J* = 8.1Hz, ¹⁶CH), 5.591 (*s*, 2H, ¹⁰CH₂). ¹³C NMR (125MHz, DMSO) in ppm: 184.08 (C9), 166.60 (C11), 154.96 (C1), 151.62 (C15), 149.63 (C13), 134.49 (C8), 130.78 (C17), 128.88 (C3), 126.49 (C6), 124.28 (C5), 123.84 (C4), 116.63 (C14), 114.84 (C16), 114.66 (C7), 112.31 (C2), 107.87 (C2), 65.63 (C10). FT-IR (ATR (solid) cm⁻¹): 3473, 3360 (N–H, *v*), 3078 (Ar C–H, *v*), 2942 (C–H, *v*), 1697, 1676 (C=O, *v*), 1615 (C=C, *v*), 1583, 1487 (Ar C=C, *v*), 1244, 1112 (C–O, *v*).

6. Refinement

Crystal data, data collection and structure refinement details for both compounds are summarized in Table 3. All C-bound H atoms were positioned geometrically (C–H = 0.93–0.97 Å) and refined using a riding model with *U*_{iso}(H) = 1.2*U*_{eq}(parent

atom). The N-bound H atoms of compound (II) were located in a difference-Fourier map and refined freely.

Acknowledgements

LYT thanks Universiti Sains Malaysia for the USM Fellowship Scheme and the Malaysian Government for a MyBrain15 (MyMaster) scholarship. HCK thanks the Malaysian Government for a MyBrain15 (MyPhD) scholarship. The authors extend their appreciation to Vidya Vikas Research & Development Center for the facilities and encouragement.

References

- Bruker (2009). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Groom, C. R., Bruno, I. J., Lightfoot, M. P. & Ward, S. C. (2016). *Acta Cryst.* **B72**, 171–179.
- Kumar, C., Then, L., Chia, T., Chandrabu, S., Win, Y.-F., Sulaiman, S., Hashim, N., Ooi, K., Quah, C. & Fun, H.-K. (2015). *Molecules*, **20**, 16566–16581.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Naik, N., Vijay, K., Dias, S. M. & Ranga, S. (2013). *Int. J. Pharm. Pharm. Sci.* **5**, 242–247.
- Nevagi, R. J., Dighe, S. N. & Dighe, S. N. (2015). *Eur. J. Med. Chem.* **97**, 561–581.
- Sheldrick, G. M. (2015a). *Acta Cryst.* **A71**, 3–8.
- Sheldrick, G. M. (2015b). *Acta Cryst.* **C71**, 3–8.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Swamy, P. M. G., Prasad, Y. R., Ashvini, H. M., Giles, D., Shashidhar, B. V. & Agasimundin, Y. S. (2015). *Med. Chem. Res.* **24**, 3437–3452.
- Ugale, V., Patel, H., Patel, B. & Bari, S. (2012). *Arab. J. Chem.* **10**, S389–A396.
- Zhou, X., Li, M., Wang, X.-B., Wang, T. & Kong, L.-Y. (2010). *Molecules*, **15**, 8593–8601.

supporting information

Acta Cryst. (2017). E73, 1087-1091 [https://doi.org/10.1107/S2056989017009422]

Two closely related 2-(benzofuran-2-yl)-2-oxoethyl benzoates: structural differences and C—H...O hydrogen-bonded supramolecular assemblies

Li Yee Then, C. S. Chidan Kumar, Huey Chong Kwong, Yip-Foo Win, Siau Hui Mah, Ching Kheng Quah, S. Naveen and Ismail Warad

Computing details

For both compounds, data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINTE* (Bruker, 2009); data reduction: *SAINTE* (Bruker, 2009); program(s) used to solve structure: *SHELXT2013* (Sheldrick, 2015a); program(s) used to refine structure: *SHELXL2013* (Sheldrick, 2015b). Molecular graphics: *SHELXL2013* (Sheldrick, 2015b) and *Mercury* (Macrae *et al.*, 2006) for (I); *SHELXL2013* (Sheldrick, 2015b) for (II). For both compounds, software used to prepare material for publication: *SHELXL2013* (Sheldrick, 2015b) and *PLATON* (Spek, 2009).

(I) 2-(1*H*-1-Benzofuran-2-yl)-2-oxoethyl 2-nitrobenzoate

Crystal data

$C_{17}H_{11}NO_6$	$D_x = 1.477 \text{ Mg m}^{-3}$
$M_r = 325.27$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Orthorhombic, $Pna2_1$	Cell parameters from 3687 reflections
$a = 9.3022 (10) \text{ \AA}$	$\theta = 2.3\text{--}25.3^\circ$
$b = 28.482 (3) \text{ \AA}$	$\mu = 0.11 \text{ mm}^{-1}$
$c = 5.5208 (6) \text{ \AA}$	$T = 100 \text{ K}$
$V = 1462.7 (3) \text{ \AA}^3$	Block, colourless
$Z = 4$	$0.27 \times 0.16 \times 0.13 \text{ mm}$
$F(000) = 672$	

Data collection

Bruker APEXII DUO CCD area-detector diffractometer	15875 measured reflections
Radiation source: fine-focus sealed tube	3358 independent reflections
Graphite monochromator	2915 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.037$
Absorption correction: multi-scan (SADABS; Bruker, 2009)	$\theta_{\text{max}} = 27.6^\circ$, $\theta_{\text{min}} = 1.4^\circ$
$T_{\text{min}} = 0.933$, $T_{\text{max}} = 0.985$	$h = -12 \rightarrow 12$
	$k = -36 \rightarrow 37$
	$l = -7 \rightarrow 7$

Refinement

Refinement on F^2	217 parameters
Least-squares matrix: full	1 restraint
$R[F^2 > 2\sigma(F^2)] = 0.038$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.085$	H-atom parameters constrained
$S = 1.08$	
3358 reflections	

$$w = 1/[\sigma^2(F_o^2) + (0.0345P)^2 + 0.279P],$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$

$$\Delta\rho_{\max} = 0.18 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.17 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.5075 (2)	0.27391 (8)	0.1493 (4)	0.0290 (5)
O1	0.72961 (19)	0.53112 (6)	0.6220 (3)	0.0309 (4)
O2	0.51143 (18)	0.39614 (6)	0.2506 (3)	0.0289 (4)
O3	0.5376 (2)	0.45933 (6)	0.6044 (4)	0.0328 (4)
O4	0.6811 (2)	0.36342 (6)	0.4887 (4)	0.0350 (5)
O5	0.5946 (2)	0.30243 (6)	0.0721 (4)	0.0343 (4)
O6	0.4877 (2)	0.23524 (7)	0.0592 (4)	0.0460 (6)
C1	0.8415 (3)	0.56187 (8)	0.5803 (5)	0.0270 (6)
C2	0.8830 (3)	0.59794 (10)	0.7294 (6)	0.0362 (6)
H2A	0.8343	0.6045	0.8768	0.043*
C3	0.9982 (3)	0.62384 (10)	0.6543 (6)	0.0412 (7)
H3A	1.0313	0.6488	0.7539	0.049*
C4	1.0686 (3)	0.61491 (10)	0.4372 (6)	0.0413 (8)
H4A	1.1475	0.6341	0.3910	0.050*
C5	1.0259 (3)	0.57859 (11)	0.2866 (6)	0.0380 (7)
H5A	1.0743	0.5725	0.1385	0.046*
C6	0.9074 (3)	0.55093 (9)	0.3613 (5)	0.0278 (6)
C7	0.8303 (3)	0.51167 (9)	0.2669 (5)	0.0284 (6)
H7A	0.8489	0.4958	0.1187	0.034*
C8	0.7272 (3)	0.50142 (9)	0.4259 (5)	0.0312 (6)
C9	0.6161 (3)	0.46488 (9)	0.4317 (5)	0.0286 (6)
C10	0.6080 (3)	0.43440 (9)	0.2082 (5)	0.0299 (6)
H10A	0.5737	0.4533	0.0691	0.036*
H10B	0.7047	0.4221	0.1685	0.036*
C11	0.5621 (3)	0.36359 (9)	0.4045 (5)	0.0272 (6)
C12	0.4463 (3)	0.32984 (8)	0.4742 (5)	0.0250 (5)
C13	0.4219 (3)	0.28707 (9)	0.3614 (4)	0.0247 (5)
C14	0.3186 (3)	0.25567 (9)	0.4415 (5)	0.0307 (6)
H14A	0.3052	0.2264	0.3622	0.037*
C15	0.2354 (3)	0.26790 (10)	0.6397 (5)	0.0345 (6)
H15A	0.1640	0.2469	0.6979	0.041*
C16	0.2562 (3)	0.31066 (10)	0.7531 (5)	0.0348 (6)
H16A	0.1978	0.3191	0.8874	0.042*
C17	0.3614 (3)	0.34129 (10)	0.6725 (5)	0.0319 (6)
H17A	0.3755	0.3704	0.7535	0.038*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0311 (11)	0.0268 (11)	0.0292 (12)	0.0017 (9)	0.0055 (10)	-0.0006 (10)
O1	0.0304 (10)	0.0307 (9)	0.0316 (9)	0.0006 (8)	0.0030 (8)	0.0025 (8)
O2	0.0337 (10)	0.0216 (9)	0.0314 (9)	0.0024 (7)	-0.0008 (9)	0.0025 (8)
O3	0.0371 (10)	0.0267 (9)	0.0346 (10)	0.0017 (8)	0.0038 (9)	-0.0001 (8)
O4	0.0278 (10)	0.0305 (10)	0.0467 (12)	0.0005 (8)	-0.0025 (9)	0.0064 (9)
O5	0.0374 (10)	0.0291 (9)	0.0364 (11)	0.0010 (8)	0.0133 (9)	0.0021 (9)
O6	0.0516 (13)	0.0338 (11)	0.0527 (13)	-0.0047 (10)	0.0184 (11)	-0.0161 (10)
C1	0.0267 (13)	0.0262 (12)	0.0280 (13)	0.0043 (10)	-0.0017 (11)	0.0067 (11)
C2	0.0419 (16)	0.0360 (16)	0.0307 (14)	0.0065 (13)	-0.0075 (13)	-0.0020 (12)
C3	0.0426 (17)	0.0341 (15)	0.0470 (18)	0.0033 (13)	-0.0196 (15)	0.0017 (14)
C4	0.0293 (15)	0.0395 (17)	0.055 (2)	-0.0040 (13)	-0.0109 (15)	0.0193 (15)
C5	0.0326 (15)	0.0502 (18)	0.0312 (15)	0.0123 (13)	0.0035 (13)	0.0142 (13)
C6	0.0281 (13)	0.0299 (14)	0.0256 (13)	0.0072 (11)	-0.0027 (11)	0.0043 (11)
C7	0.0316 (14)	0.0247 (13)	0.0289 (13)	0.0081 (11)	-0.0035 (12)	-0.0022 (11)
C8	0.0349 (14)	0.0230 (12)	0.0357 (15)	0.0058 (11)	-0.0062 (13)	-0.0002 (11)
C9	0.0293 (13)	0.0222 (12)	0.0342 (14)	0.0065 (11)	-0.0007 (12)	0.0042 (11)
C10	0.0343 (14)	0.0208 (12)	0.0345 (15)	0.0017 (11)	0.0044 (12)	0.0032 (11)
C11	0.0299 (14)	0.0217 (12)	0.0299 (14)	0.0059 (11)	0.0028 (12)	-0.0010 (11)
C12	0.0234 (12)	0.0244 (12)	0.0273 (13)	0.0056 (10)	0.0004 (11)	0.0021 (11)
C13	0.0233 (12)	0.0281 (13)	0.0227 (12)	0.0052 (10)	0.0021 (10)	0.0008 (11)
C14	0.0301 (13)	0.0308 (14)	0.0313 (13)	-0.0033 (11)	-0.0001 (12)	-0.0007 (12)
C15	0.0275 (13)	0.0436 (16)	0.0324 (14)	-0.0054 (12)	0.0042 (12)	0.0049 (13)
C16	0.0289 (14)	0.0471 (17)	0.0286 (13)	0.0011 (13)	0.0069 (12)	-0.0014 (14)
C17	0.0320 (14)	0.0343 (15)	0.0295 (14)	0.0050 (12)	0.0012 (12)	-0.0051 (12)

Geometric parameters (Å, °)

N1—O6	1.223 (3)	C6—C7	1.427 (4)
N1—O5	1.224 (3)	C7—C8	1.333 (4)
N1—C13	1.465 (3)	C7—H7A	0.9500
O1—C8	1.374 (3)	C8—C9	1.467 (4)
O1—C1	1.380 (3)	C9—C10	1.511 (4)
O2—C11	1.343 (3)	C10—H10A	0.9900
O2—C10	1.431 (3)	C10—H10B	0.9900
O3—C9	1.211 (3)	C11—C12	1.494 (4)
O4—C11	1.201 (3)	C12—C13	1.387 (3)
C1—C2	1.372 (4)	C12—C17	1.389 (4)
C1—C6	1.391 (4)	C13—C14	1.385 (4)
C2—C3	1.365 (4)	C14—C15	1.385 (4)
C2—H2A	0.9500	C14—H14A	0.9500
C3—C4	1.389 (5)	C15—C16	1.383 (4)
C3—H3A	0.9500	C15—H15A	0.9500
C4—C5	1.385 (4)	C16—C17	1.384 (4)
C4—H4A	0.9500	C16—H16A	0.9500
C5—C6	1.416 (4)	C17—H17A	0.9500

C5—H5A	0.9500		
O6—N1—O5	123.8 (2)	O3—C9—C10	122.6 (2)
O6—N1—C13	118.3 (2)	C8—C9—C10	115.1 (2)
O5—N1—C13	117.9 (2)	O2—C10—C9	109.6 (2)
C8—O1—C1	105.8 (2)	O2—C10—H10A	109.8
C11—O2—C10	114.1 (2)	C9—C10—H10A	109.8
C2—C1—O1	126.0 (3)	O2—C10—H10B	109.8
C2—C1—C6	124.4 (3)	C9—C10—H10B	109.8
O1—C1—C6	109.6 (2)	H10A—C10—H10B	108.2
C3—C2—C1	116.3 (3)	O4—C11—O2	124.9 (2)
C3—C2—H2A	121.8	O4—C11—C12	124.2 (2)
C1—C2—H2A	121.8	O2—C11—C12	110.7 (2)
C2—C3—C4	122.2 (3)	C13—C12—C17	117.8 (2)
C2—C3—H3A	118.9	C13—C12—C11	124.6 (2)
C4—C3—H3A	118.9	C17—C12—C11	117.5 (2)
C5—C4—C3	121.3 (3)	C14—C13—C12	122.5 (2)
C5—C4—H4A	119.4	C14—C13—N1	117.8 (2)
C3—C4—H4A	119.4	C12—C13—N1	119.7 (2)
C4—C5—C6	117.6 (3)	C13—C14—C15	118.5 (3)
C4—C5—H5A	121.2	C13—C14—H14A	120.7
C6—C5—H5A	121.2	C15—C14—H14A	120.7
C1—C6—C5	118.1 (3)	C16—C15—C14	120.1 (3)
C1—C6—C7	105.7 (2)	C16—C15—H15A	120.0
C5—C6—C7	136.1 (3)	C14—C15—H15A	120.0
C8—C7—C6	107.0 (2)	C15—C16—C17	120.6 (3)
C8—C7—H7A	126.5	C15—C16—H16A	119.7
C6—C7—H7A	126.5	C17—C16—H16A	119.7
C7—C8—O1	111.9 (2)	C16—C17—C12	120.5 (3)
C7—C8—C9	132.6 (3)	C16—C17—H17A	119.8
O1—C8—C9	115.5 (2)	C12—C17—H17A	119.8
O3—C9—C8	122.3 (3)		
C8—O1—C1—C2	179.6 (2)	O3—C9—C10—O2	-7.9 (3)
C8—O1—C1—C6	-0.3 (3)	C8—C9—C10—O2	171.4 (2)
O1—C1—C2—C3	179.2 (2)	C10—O2—C11—O4	-5.4 (4)
C6—C1—C2—C3	-0.8 (4)	C10—O2—C11—C12	169.6 (2)
C1—C2—C3—C4	1.1 (4)	O4—C11—C12—C13	-90.2 (4)
C2—C3—C4—C5	-0.8 (4)	O2—C11—C12—C13	94.7 (3)
C3—C4—C5—C6	0.3 (4)	O4—C11—C12—C17	86.9 (3)
C2—C1—C6—C5	0.3 (4)	O2—C11—C12—C17	-88.2 (3)
O1—C1—C6—C5	-179.7 (2)	C17—C12—C13—C14	-1.0 (4)
C2—C1—C6—C7	-179.7 (2)	C11—C12—C13—C14	176.1 (2)
O1—C1—C6—C7	0.3 (3)	C17—C12—C13—N1	179.1 (2)
C4—C5—C6—C1	0.0 (4)	C11—C12—C13—N1	-3.8 (4)
C4—C5—C6—C7	-180.0 (3)	O6—N1—C13—C14	-2.8 (3)
C1—C6—C7—C8	-0.1 (3)	O5—N1—C13—C14	177.0 (2)
C5—C6—C7—C8	179.9 (3)	O6—N1—C13—C12	177.1 (2)

C6—C7—C8—O1	−0.1 (3)	O5—N1—C13—C12	−3.1 (3)
C6—C7—C8—C9	−179.1 (3)	C12—C13—C14—C15	1.0 (4)
C1—O1—C8—C7	0.3 (3)	N1—C13—C14—C15	−179.1 (2)
C1—O1—C8—C9	179.4 (2)	C13—C14—C15—C16	0.0 (4)
C7—C8—C9—O3	174.5 (3)	C14—C15—C16—C17	−1.0 (4)
O1—C8—C9—O3	−4.5 (3)	C15—C16—C17—C12	1.0 (4)
C7—C8—C9—C10	−4.9 (4)	C13—C12—C17—C16	0.1 (4)
O1—C8—C9—C10	176.2 (2)	C11—C12—C17—C16	−177.3 (2)
C11—O2—C10—C9	−71.4 (3)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C10—H10 <i>A</i> ...O3 ⁱ	0.99	2.59	3.471 (4)	148
C15—H15 <i>A</i> ...O5 ⁱⁱ	0.95	2.58	3.380 (3)	142

Symmetry codes: (i) *x*, *y*, *z*−1; (ii) *x*−1/2, −*y*+1/2, *z*+1.(II) 2-(1*H*-1-Benzofuran-2-yl)-2-oxoethyl 2-aminobenzoate

Crystal data

C₁₇H₁₃NO₄*M_r* = 295.28Triclinic, *P*1̄*a* = 5.1839 (12) Å*b* = 10.853 (3) Å*c* = 12.269 (3) Å

α = 93.562 (3)°

β = 91.167 (3)°

γ = 98.714 (3)°

V = 680.6 (3) Å³*Z* = 2*F*(000) = 308*D_x* = 1.441 Mg m^{−3}Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 5796 reflections

θ = 2.4–27.9°

μ = 0.10 mm^{−1}*T* = 297 K

Block, orange

0.40 × 0.32 × 0.21 mm

Data collection

Bruker APEXII DUO CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan

(SADABS; Bruker, 2009)

T_{min} = 0.871, *T_{max}* = 0.978

17052 measured reflections

3105 independent reflections

2214 reflections with *I* > 2σ(*I*)*R_{int}* = 0.037θ_{max} = 27.5°, θ_{min} = 1.7°*h* = −6→6*k* = −14→14*l* = −15→15

Refinement

Refinement on *F*²

Least-squares matrix: full

R[*F*² > 2σ(*F*²)] = 0.046*wR*(*F*²) = 0.123*S* = 1.08

3105 reflections

207 parameters

0 restraints

Hydrogen site location: mixed

H atoms treated by a mixture of independent
and constrained refinement*w* = 1/[σ²(*F_o*²) + (0.0462*P*)² + 0.1899*P*]where *P* = (*F_o*² + 2*F_c*²)/3(Δ/σ)_{max} < 0.001Δρ_{max} = 0.18 e Å^{−3}Δρ_{min} = −0.18 e Å^{−3}

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.6563 (4)	0.67887 (18)	0.63733 (16)	0.0583 (4)
H1A	0.611 (4)	0.617 (2)	0.5840 (19)	0.069 (7)*
H1B	0.797 (5)	0.682 (2)	0.6702 (19)	0.074 (7)*
O1	0.2314 (2)	0.47254 (11)	0.10178 (9)	0.0478 (3)
O2	0.0514 (2)	0.72867 (11)	0.41702 (9)	0.0500 (3)
O3	0.3365 (3)	0.67869 (12)	0.24247 (11)	0.0587 (4)
O4	0.2949 (3)	0.59366 (11)	0.47797 (11)	0.0579 (4)
C1	0.1297 (3)	0.35882 (16)	0.05120 (14)	0.0448 (4)
C2	0.2360 (4)	0.3001 (2)	-0.03521 (16)	0.0581 (5)
H2A	0.3890	0.3356	-0.0667	0.070*
C3	0.1032 (5)	0.1866 (2)	-0.07204 (17)	0.0677 (6)
H3A	0.1669	0.1434	-0.1309	0.081*
C4	-0.1240 (5)	0.1334 (2)	-0.02449 (18)	0.0672 (6)
H4A	-0.2072	0.0551	-0.0514	0.081*
C5	-0.2283 (4)	0.19380 (18)	0.06135 (17)	0.0585 (5)
H5A	-0.3814	0.1581	0.0926	0.070*
C6	-0.0976 (3)	0.31045 (16)	0.10033 (14)	0.0452 (4)
C7	-0.1376 (3)	0.40069 (16)	0.18454 (14)	0.0453 (4)
H7A	-0.2762	0.3957	0.2318	0.054*
C8	0.0625 (3)	0.49452 (16)	0.18298 (13)	0.0429 (4)
C9	0.1355 (3)	0.60838 (16)	0.25240 (14)	0.0439 (4)
C10	-0.0582 (3)	0.63287 (18)	0.33799 (14)	0.0507 (4)
H10A	-0.2107	0.6571	0.3031	0.061*
H10B	-0.1141	0.5567	0.3741	0.061*
C11	0.2338 (3)	0.69697 (16)	0.48521 (13)	0.0431 (4)
C12	0.3415 (3)	0.79867 (15)	0.56403 (13)	0.0407 (4)
C13	0.5539 (3)	0.78708 (16)	0.63365 (13)	0.0441 (4)
C14	0.6573 (4)	0.89064 (19)	0.70265 (15)	0.0567 (5)
H14A	0.7996	0.8855	0.7486	0.068*
C15	0.5549 (4)	0.9990 (2)	0.70424 (16)	0.0611 (5)
H15A	0.6289	1.0667	0.7506	0.073*
C16	0.3431 (4)	1.00947 (18)	0.63798 (16)	0.0603 (5)
H16A	0.2715	1.0831	0.6403	0.072*
C17	0.2401 (4)	0.91065 (17)	0.56910 (15)	0.0515 (4)
H17A	0.0976	0.9179	0.5240	0.062*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0538 (10)	0.0643 (11)	0.0604 (11)	0.0199 (9)	-0.0078 (8)	0.0097 (9)
O1	0.0432 (7)	0.0505 (7)	0.0489 (7)	0.0035 (5)	0.0037 (5)	0.0054 (5)
O2	0.0511 (7)	0.0514 (7)	0.0492 (7)	0.0158 (6)	-0.0066 (6)	-0.0011 (6)
O3	0.0505 (8)	0.0579 (8)	0.0635 (8)	-0.0041 (6)	0.0051 (6)	-0.0002 (6)
O4	0.0647 (8)	0.0426 (7)	0.0680 (8)	0.0158 (6)	-0.0080 (7)	0.0023 (6)
C1	0.0456 (9)	0.0457 (10)	0.0441 (9)	0.0097 (8)	-0.0051 (7)	0.0072 (8)
C2	0.0588 (12)	0.0676 (13)	0.0508 (11)	0.0174 (10)	0.0053 (9)	0.0058 (9)
C3	0.0846 (16)	0.0697 (14)	0.0532 (12)	0.0292 (12)	-0.0042 (11)	-0.0027 (10)
C4	0.0846 (16)	0.0519 (12)	0.0630 (13)	0.0095 (11)	-0.0195 (12)	-0.0043 (10)
C5	0.0565 (11)	0.0540 (11)	0.0622 (12)	-0.0018 (9)	-0.0073 (9)	0.0083 (10)
C6	0.0457 (9)	0.0462 (10)	0.0442 (9)	0.0075 (8)	-0.0060 (7)	0.0081 (8)
C7	0.0399 (9)	0.0523 (10)	0.0438 (9)	0.0047 (8)	0.0017 (7)	0.0087 (8)
C8	0.0408 (9)	0.0489 (10)	0.0409 (9)	0.0105 (7)	-0.0014 (7)	0.0093 (7)
C9	0.0405 (9)	0.0458 (10)	0.0458 (9)	0.0065 (8)	-0.0051 (7)	0.0080 (7)
C10	0.0432 (10)	0.0576 (11)	0.0508 (10)	0.0083 (8)	-0.0033 (8)	-0.0008 (8)
C11	0.0422 (9)	0.0447 (10)	0.0444 (9)	0.0098 (7)	0.0042 (7)	0.0093 (7)
C12	0.0416 (9)	0.0414 (9)	0.0403 (8)	0.0078 (7)	0.0058 (7)	0.0064 (7)
C13	0.0417 (9)	0.0511 (10)	0.0404 (9)	0.0064 (8)	0.0069 (7)	0.0104 (7)
C14	0.0538 (11)	0.0684 (13)	0.0461 (10)	0.0039 (10)	-0.0026 (8)	0.0032 (9)
C15	0.0724 (13)	0.0573 (12)	0.0488 (11)	-0.0023 (10)	0.0052 (9)	-0.0049 (9)
C16	0.0776 (14)	0.0471 (11)	0.0579 (11)	0.0156 (10)	0.0074 (10)	-0.0006 (9)
C17	0.0565 (11)	0.0482 (10)	0.0520 (10)	0.0152 (9)	0.0008 (8)	0.0034 (8)

Geometric parameters (Å, °)

N1—C13	1.363 (2)	C6—C7	1.419 (2)
N1—H1A	0.91 (2)	C7—C8	1.340 (2)
N1—H1B	0.82 (2)	C7—H7A	0.9300
O1—C1	1.371 (2)	C8—C9	1.453 (2)
O1—C8	1.373 (2)	C9—C10	1.507 (3)
O2—C11	1.3469 (19)	C10—H10A	0.9700
O2—C10	1.420 (2)	C10—H10B	0.9700
O3—C9	1.208 (2)	C11—C12	1.457 (2)
O4—C11	1.209 (2)	C12—C17	1.394 (2)
C1—C2	1.371 (3)	C12—C13	1.405 (2)
C1—C6	1.382 (2)	C13—C14	1.396 (3)
C2—C3	1.363 (3)	C14—C15	1.361 (3)
C2—H2A	0.9300	C14—H14A	0.9300
C3—C4	1.387 (3)	C15—C16	1.376 (3)
C3—H3A	0.9300	C15—H15A	0.9300
C4—C5	1.370 (3)	C16—C17	1.358 (3)
C4—H4A	0.9300	C16—H16A	0.9300
C5—C6	1.393 (3)	C17—H17A	0.9300
C5—H5A	0.9300		

C13—N1—H1A	119.3 (14)	O3—C9—C10	122.21 (16)
C13—N1—H1B	117.8 (16)	C8—C9—C10	115.14 (15)
H1A—N1—H1B	118 (2)	O2—C10—C9	111.46 (14)
C1—O1—C8	105.72 (13)	O2—C10—H10A	109.3
C11—O2—C10	115.04 (13)	C9—C10—H10A	109.3
C2—C1—O1	125.63 (17)	O2—C10—H10B	109.3
C2—C1—C6	124.29 (18)	C9—C10—H10B	109.3
O1—C1—C6	110.08 (15)	H10A—C10—H10B	108.0
C3—C2—C1	115.8 (2)	O4—C11—O2	121.09 (16)
C3—C2—H2A	122.1	O4—C11—C12	126.02 (15)
C1—C2—H2A	122.1	O2—C11—C12	112.89 (14)
C2—C3—C4	122.1 (2)	C17—C12—C13	118.93 (16)
C2—C3—H3A	119.0	C17—C12—C11	120.32 (15)
C4—C3—H3A	119.0	C13—C12—C11	120.72 (15)
C5—C4—C3	121.3 (2)	N1—C13—C14	119.72 (17)
C5—C4—H4A	119.3	N1—C13—C12	122.51 (17)
C3—C4—H4A	119.3	C14—C13—C12	117.76 (16)
C4—C5—C6	117.9 (2)	C15—C14—C13	121.56 (18)
C4—C5—H5A	121.1	C15—C14—H14A	119.2
C6—C5—H5A	121.1	C13—C14—H14A	119.2
C1—C6—C5	118.65 (17)	C14—C15—C16	120.74 (19)
C1—C6—C7	105.84 (15)	C14—C15—H15A	119.6
C5—C6—C7	135.50 (18)	C16—C15—H15A	119.6
C8—C7—C6	106.96 (16)	C17—C16—C15	119.00 (18)
C8—C7—H7A	126.5	C17—C16—H16A	120.5
C6—C7—H7A	126.5	C15—C16—H16A	120.5
C7—C8—O1	111.38 (15)	C16—C17—C12	121.99 (18)
C7—C8—C9	132.36 (17)	C16—C17—H17A	119.0
O1—C8—C9	116.19 (14)	C12—C17—H17A	119.0
O3—C9—C8	122.65 (17)		
C8—O1—C1—C2	179.91 (16)	O1—C8—C9—C10	177.70 (13)
C8—O1—C1—C6	-0.32 (16)	C11—O2—C10—C9	-70.85 (18)
O1—C1—C2—C3	179.33 (15)	O3—C9—C10—O2	-13.3 (2)
C6—C1—C2—C3	-0.4 (3)	C8—C9—C10—O2	166.80 (13)
C1—C2—C3—C4	-0.5 (3)	C10—O2—C11—O4	-0.3 (2)
C2—C3—C4—C5	1.0 (3)	C10—O2—C11—C12	179.52 (14)
C3—C4—C5—C6	-0.5 (3)	O4—C11—C12—C17	-175.17 (17)
C2—C1—C6—C5	0.8 (3)	O2—C11—C12—C17	5.1 (2)
O1—C1—C6—C5	-178.95 (14)	O4—C11—C12—C13	6.8 (3)
C2—C1—C6—C7	-179.49 (16)	O2—C11—C12—C13	-172.97 (14)
O1—C1—C6—C7	0.74 (17)	C17—C12—C13—N1	176.84 (17)
C4—C5—C6—C1	-0.3 (2)	C11—C12—C13—N1	-5.1 (2)
C4—C5—C6—C7	-179.90 (18)	C17—C12—C13—C14	-1.8 (2)
C1—C6—C7—C8	-0.87 (18)	C11—C12—C13—C14	176.29 (15)
C5—C6—C7—C8	178.74 (18)	N1—C13—C14—C15	-177.67 (18)
C6—C7—C8—O1	0.71 (18)	C12—C13—C14—C15	1.0 (3)
C6—C7—C8—C9	-176.21 (16)	C13—C14—C15—C16	0.6 (3)

C1—O1—C8—C7	-0.26 (17)	C14—C15—C16—C17	-1.3 (3)
C1—O1—C8—C9	177.21 (13)	C15—C16—C17—C12	0.4 (3)
C7—C8—C9—O3	174.59 (17)	C13—C12—C17—C16	1.1 (3)
O1—C8—C9—O3	-2.2 (2)	C11—C12—C17—C16	-176.94 (17)
C7—C8—C9—C10	-5.5 (3)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1A \cdots O4	0.91 (2)	2.05 (2)	2.700 (3)	127.7 (18)
N1—H1A \cdots O4 ⁱ	0.91 (2)	2.49 (2)	3.246 (2)	141.4 (18)
C10—H10A \cdots O3 ⁱⁱ	0.97	2.50	3.444 (2)	165
C17—H17A \cdots O2	0.93	2.35	2.687 (2)	101

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $x-1, y, z$.