

Received 2 May 2022

Accepted 5 June 2022

Edited by G. Díaz de Delgado, Universidad de Los Andes, Venezuela

Keywords: crystal structure; hydrogen bonding; piperazine; biological activity.**CCDC references:** 2177037; 2177036; 2177035; 2177034**Supporting information:** this article has supporting information at journals.iucr.org/e

Crystal-structure studies of 4-phenylpiperazin-1-ium 4-ethoxybenzoate monohydrate, 4-phenylpiperazin-1-ium 4-methoxybenzoate monohydrate, 4-phenylpiperazin-1-ium 4-methylbenzoate monohydrate and 4-phenylpiperazin-1-ium trifluoroacetate 0.12-hydrate

Ninganayaka Mahesha,^a Haruvegowda Kiran Kumar,^a Mehmet Akkurt,^b Hemmige S. Yathirajan,^a Sabine Foro,^c Mohammed S. M. Abdelbaky^d and Santiago Garcia-Granda^{d*}


^aDepartment of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore-570 006, India, ^bDepartment of Physics, Faculty of Sciences, Erciyes University, 38039 Kayseri, Turkey, ^cInstitute of Materials Science, Darmstadt University of Technology, Alarich-Weiss-Strasse 2, D-64287 Darmstadt, Germany, and ^dDepartment of Physical and Analytical Chemistry, Faculty of Chemistry, Oviedo University-CINN, Oviedo 33006, Spain. *Correspondence e-mail: sgg@uniovi.es

In this study, four new piperazinium salts, namely, 4-phenylpiperazin-1-ium 4-ethoxybenzoate monohydrate, $C_9H_9O_3 \cdot C_{10}H_{15}N_2 \cdot H_2O$ (**I**); 4-phenylpiperazin-1-ium 4-methoxybenzoate monohydrate, $C_{10}H_{15}N_2 \cdot C_8H_7O_3 \cdot H_2O$ (**II**); 4-phenylpiperazin-1-ium 4-methylbenzoate monohydrate, $C_{10}H_{15}N_2 \cdot C_8H_7O_2 \cdot H_2O$ (**III**); and 4-phenylpiperazin-1-ium trifluoroacetate 0.12 hydrate, $C_{10}H_{15}N_2 \cdot C_2F_3O_2 \cdot 0.12H_2O$ (**IV**), have been synthesized. The single-crystal structures of these compounds reveal that all of them crystallize in the triclinic $P\bar{1}$ space group and the crystal packing of (**I**)–(**III**) is built up of ribbons formed by a combination of hydrogen bonds of type $N-H \cdots O$, $O-H \cdots O$ and other weak interactions of type $C-H \cdots O$ and $C-H \cdots \pi$, leading to a three-dimensional network. In the crystal of (**IV**), the cations and the anions are connected by $C-H \cdots O$, $N-H \cdots O$ and $C-H \cdots F$ hydrogen bonds and by $C-H \cdots \pi$ interactions, forming sheets which in turn interact to maintain the crystal structure by linking through the oxygen atoms of water molecules and van der Waals interactions, giving the whole structure.

1. Chemical context

Piperazines are among the most important building blocks in today's drug discovery efforts and are found in biologically active compounds across a number of different therapeutic areas (Brockunier *et al.*, 2004; Bogatcheva *et al.*, 2006). For a review on the current pharmacological and toxicological information for piperazine derivative, see Elliott (2011). Various pharmacological properties of phenylpiperazines and their derivatives have been discussed by several authors (Cohen *et al.*, 1982; Conrado *et al.*, 2010; Neves *et al.*, 2003; Hanano *et al.*, 2000). The design and synthesis of phenylpiperazine derivatives as potent anticancer agents for prostate cancer have been described (Demirci *et al.*, 2019). Many pharmaceutical compounds are derived from 1-phenylpiperazine, *viz.*, oxyperline, trazodone, nefazodone, *etc.*


The crystal structures of 2-(4-methyl-2-phenylpiperazin-4-ium-1-yl)pyridine-3-carboxylate dehydrate (Li *et al.*, 2008),



OPEN ACCESS

Published under a CC BY 4.0 licence

1-chloro-2-(4-phenylpiperazin-1-yl)-ethanone (Xu & Jing, 2009), 4-phenylpiperazin-1-ium dihydrogen phosphate (Essid *et al.*, 2010) and 1-phenylpiperazine-1,4-diium bis(hydrogen sulfate) (Marouani *et al.*, 2010) have been reported, as have those of 4-phenylpiperazin-1-ium 6-chloro-5-ethyl-2,4-dioxopyrimidin-1-ide and 4-phenylpiperazin-1-ium 6-chloro-5-isopropyl-2,4-dioxopyrimidin-1-ide (Al-Alshaikh *et al.*, 2015). We have reported the crystal structures of some salts of 4-methoxyphenylpiperazine (Kiran Kumar *et al.*, 2019a), six 1-aryol-4-(4-methoxyphenyl)piperazines (Kiran Kumar *et al.*, 2019b), 2-methoxyphenylpiperazine (Harish Chinthali *et al.*, 2020) and the recreational drug *N*-(4-methoxyphenyl)piperazine (MeOPP) and three of its salts (Kiran Kumar *et al.*, 2020a).



In view of the importance of piperazines in general and the use of 1-phenylpiperazine in particular, the present paper reports the crystal structure studies of some salts of 1-phenylpiperazine with organic acids *viz.*, 4-phenylpiperazin-1-ium 4-ethoxybenzoate monohydrate, $C_9H_9O_3 \cdot C_{10}H_{15}N_2 \cdot H_2O$ (**I**); 4-phenylpiperazin-1-ium 4-methoxybenzoate monohydrate, $C_{10}H_{15}N_2 \cdot C_8H_7O_3 \cdot H_2O$ (**II**); 4-phenylpiperazin-1-ium 4-methylbenzoate monohydrate, $C_{10}H_{15}N_2 \cdot C_8H_7O_2 \cdot H_2O$ (**III**);




Figure 1
The independent components of compound (**I**) showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level.




Figure 2
The independent components of compound (**II**) showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level.

and 4-phenylpiperazin-1-ium trifluoroacetate 0.12 hydrate $C_{10}H_{15}N_2 \cdot C_2F_3O_2 \cdot 0.12H_2O$ (**IV**).

2. Structural commentary

The asymmetric unit of the compound (**I**), (Fig. 1), consists of a 4-phenylpiperazin-1-ium cation, a 4-ethoxybenzoate anion and one water molecule. The aromatic ring of the cation is essentially planar while the protonated piperazine ring adopts a chair conformation, with puckering parameters (Cremer & Pople, 1975) $Q_T = 0.553$ (2) Å, $\theta = 175.0$ (2)° and $\varphi = 15$ (3)°. In compound (**II**) the asymmetric unit (Fig. 2) comprises a 4-phenylpiperazin-1-ium cation, a 4-methoxybenzoate anion and one water molecule. The aromatic ring of the cation is essentially planar while the protonated piperazine ring adopts a chair conformation, with puckering parameters $Q_T = 0.5614$ (18) Å, $\theta = 175.89$ (17)° and $\varphi = 346$ (3)°. Compound (**III**) presents an asymmetric unit (Fig. 3) composed of a 4-phenylpiperazin-1-ium cation, a 4-methylbenzoate anion and one water molecule. The aromatic ring of the cation is essentially planar but the protonated piperazine ring adopts a distorted chair conformation, with puckering parameters $Q_T = 0.5486$ (19) Å, $\theta = 9.38$ (19)° and $\varphi = 167.9$ (13)°. On the other hand, the asymmetric unit of (**IV**) (Fig. 4) contains two 4-phenylpiperazin-1-ium cations (*A1* with *N1*, *A2* with *N3*) and two trifluoroacetate anions (*B1* with *F1*, *B2* with *F4*) and a 0.12 occupancy water molecule. The aromatic rings of the cations (*A1*, *A2*) are essentially planar while the protonated piperazine rings adopt a chair conformation for cation *A1*, with puckering parameters (Cremer & Pople, 1975) $Q_T = 0.552$ (4) Å, $\theta = 0.0$ (4)° and $\varphi = 207$ (14)°, and a distorted




Figure 3
The independent components of compound (**III**) showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level.




Figure 4
The independent components of compound (IV) showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level. (Atom splitting is omitted for clarity.)

chair conformation for the cation A2, with puckering parameters $Q_T = 0.559$ (5) Å, $\theta = 6.6$ (4)° and $\varphi = 168$ (4)°.

3. Supramolecular features

In the crystal structure of (I), the cation pairs are connected across two water molecules by C—H···O and N—H···O

Table 1
Hydrogen-bond geometry (Å, °) for (I).

Cg1 and Cg3 are the centroids of the C12–C17 and C1–C6 benzene rings, respectively.

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N2—HN1···OW1 ⁱ	0.89 (2)	1.94 (2)	2.817 (3)	167 (2)
N2—HN2···O1	0.93 (2)	1.80 (2)	2.724 (3)	174 (2)
OW1—HW1···O2	0.88 (3)	1.75 (3)	2.630 (3)	178 (4)
OW1—HW2···O1 ⁱⁱ	0.91 (3)	1.89 (3)	2.789 (3)	167 (3)
C9—H9A···OW1	0.97	2.52	3.308 (3)	138
C1—H1···Cg1 ⁱⁱⁱ	0.93	2.91	3.607 (3)	133
C5—H5···Cg1 ^{iv}	0.93	2.79	3.570 (3)	142
C18—H18B···Cg3 ^v	0.97	2.88	3.737 (4)	148

Symmetry codes: (i) $-x + 1, -y + 2, -z + 1$; (ii) $x + 1, y, z$; (iii) $-x, -y + 2, -z + 1$; (iv) $-x + 1, -y + 1, -z + 1$; (v) $-x, -y + 1, -z + 1$.

hydrogen bonds, forming an $R_2^4(10)$ ring motif in which the anions and cations are linked through the water molecules by O—H···O and N—H···O hydrogen bonds, forming ribbons along the *a*-axis direction (Table 1, Fig. 5*a*). In addition, a set of C—H··· π interactions, through the benzene rings of the anions and the cations, connect the molecules together in ribbons along the *a*-axis direction (Table 1, Fig. 5*b*). The C—H···O, N—H···O, O—H···O hydrogen bonds and C—H··· π interactions together form a three-dimensional network, contributing to the stabilization of the crystal structure.

In the crystal structure of (II), the cations, the anions and the water molecules are connected by C—H···O, N—H···O and O—H···O hydrogen bonds, forming ribbons along the *a*-axis direction (Table 2, Fig. 6*a*). Furthermore, the cations interact *via* C—H··· π interactions through the benzene ring of the anion, forming ribbons along the *b*-axis direction (Table 2, Fig. 6*b*). The C—H···O, N—H···O, O—H···O hydrogen bonds and C—H··· π interactions together form a




Figure 5
Parts of the crystal structure of compound (I) showing (a) the formation of a cyclic hydrogen-bonded $R_2^4(10)$ aggregate and (b) a general view of C—H··· π interactions parallel to [100]. Hydrogen bonds and C—H··· π interactions are drawn as dashed lines.

Table 2
Hydrogen-bond geometry (Å, °) for (II).

Cg3 is the centroid of the C12–C17 benzene ring.

D–H...A	D–H	H...A	D...A	D–H...A
N2–HN1...OW1 ⁱ	0.93 (2)	1.91 (2)	2.815 (2)	166 (2)
OW1–HW1...O1 ⁱⁱⁱ	0.84 (2)	1.80 (2)	2.633 (2)	175 (2)
N2–HN2...O2	0.93 (2)	1.81 (2)	2.7350 (19)	176 (2)
OW1–HW2...O2 ⁱⁱⁱ	0.85 (2)	1.96 (2)	2.7876 (19)	168 (2)
C8–H8B...OW1 ⁱⁱ	0.97	2.53	3.331 (2)	140
C1–H1...Cg3 ⁱⁱⁱ	0.93	2.76	3.549 (2)	144
C5–H5...Cg3 ^{iv}	0.93	2.86	3.625 (2)	140

Symmetry codes: (i) $x, y - 1, z$; (ii) $-x, -y + 1, -z + 1$; (iii) $-x + 1, -y + 1, -z + 1$; (iv) $-x + 1, -y, -z + 1$.

Table 3
Hydrogen-bond geometry (Å, °) for (III).

D–H...A	D–H	H...A	D...A	D–H...A
OW1–HW1...O2 ⁱ	0.89 (3)	1.90 (3)	2.782 (2)	171 (4)
OW1–HW2...O1	0.84 (2)	1.92 (3)	2.751 (2)	172 (3)
N2–HN1...O1 ⁱⁱ	0.90 (2)	1.94 (2)	2.819 (2)	164 (2)
N2–HN2...O2	0.92 (2)	1.80 (2)	2.7207 (19)	176 (2)
C8–H8A...OW1	0.97	2.33	3.116 (3)	138

Symmetry codes: (i) $x + 1, y, z$; (ii) $-x + 1, -y + 1, -z + 1$.

three-dimensional network, contributing to the stabilization of the crystal structure.

In the crystal structure of (III), the cations, the anions and the water molecules are connected by C–H...O, N–H...O and O–H...O hydrogen bonds, forming ribbons along the *a*-axis direction (Table 3, Fig. 7). There are no C–H... π interactions or π - π stacking interactions. The crystal structure is stabilized by C–H...O, N–H...O, O–H...O hydrogen bonds and van der Waals interactions between the ribbons, which run along the *a*-axis direction.

In the crystal structure of (IV), the cations and the anions are connected by C–H...O, N–H...O and C–H...F hydrogen bonds (Table 4, Fig. 8a) and C–H... π interactions, generating sheets parallel to the (100) plane (Table 4, Fig. 8).




Figure 6
Parts of the crystal structure of compound (II) showing (a) the formation of hydrogen-bonded ribbons parallel to [010] and (b) a general view of the C–H... π interactions parallel to [010]. Hydrogen bonds and C–H... π interactions are drawn as dashed lines.

Table 4
Hydrogen-bond geometry (Å, °) for (IV).

Cg2 is the centroid of the C1–C6 phenyl ring.

D–H...A	D–H	H...A	D...A	D–H...A
N2–H21...O1 ⁱ	0.88	1.91	2.790 (4)	174
N2–H22...O3	0.87	2.04	2.860 (4)	157
N2–H22...O4	0.87	2.47	3.164 (5)	137
N4–H41...O4 ⁱⁱ	0.86	1.95	2.759 (6)	156
N4–H42...O2 ⁱⁱⁱ	0.89	1.90	2.758 (4)	164
C18–H18A...F5 ^{iv}	0.97	2.53	3.273 (18)	134
C18–H18B...Ow1	0.97	2.08	2.929 (15)	145
C19–H19B...O3 ^{iv}	0.97	2.59	3.420 (5)	144
C20–H20A...F5 ^{iv}	0.97	2.64	3.468 (8)	144
C16–H16...Cg2 ^v	0.93	2.99	3.745 (4)	140

Symmetry codes: (i) $x, y + 1, z$; (ii) $x + 1, y, z$; (iii) $-x + 1, -y + 1, -z$; (iv) $-x + 1, -y + 2, -z$; (v) $-x + 1, -y + 1, -z + 1$.

These sheets further interact to maintain the crystal structure by linking through the oxygen atoms of water molecules and by van der Waals interactions. As shown in Table 4, the main interactions in the structure of (IV) involve the oxygen atoms




Figure 7
Part of the crystal structure of compound (III) showing the formation of a hydrogen-bonded chain of rings parallel to [001]. Hydrogen bonds are drawn as dashed lines.




Figure 8

Parts of the crystal structure of compound (IV) showing (a) a general view of the C—H...O, N—H...O and C—H...F hydrogen bonds and C—H... π interactions and (b) the molecular packing of (IV) down the *a*-axis. Hydrogen bonds and C—H... π interactions are drawn as dashed lines.

of carboxylate groups, while the 0.12 fraction of the water molecule contributes with one interaction of the type C—H...O and it is weak in comparison to the other oxygen-based ones.

4. Database survey

A search of the Cambridge Structural Database (Version 2020.3, last update February 2022; Groom *et al.*, 2016) for an unsubstituted 4-phenylpiperazin-1-ium cation and *para*-substituted benzoate anion involved in the reported salts (I)–(III) gave no hits. However, searching for a branched phenyl piperazinium cation and *para*-substituted benzoate anion gave comparable hits, namely; 4-(4-methoxyphenyl)piperazin-1-ium 4-fluorobenzoate monohydrate, 4-(4-methoxyphenyl)piperazin-1-ium 4-chlorobenzoate monohydrate, 4-(4-methoxyphenyl)piperazin-1-ium 4-bromobenzoate monohydrate (FOVPOY, FOVPUE, FOVQAL; Kiran Kumar *et al.*, 2019a), 4-(4-methoxyphenyl)piperazin-1-ium 4-iodobenzoate monohydrate (KUJPUD; Kiran Kumar *et al.*, 2020b). They exhibit a methoxy group as a substituent in the 4-phenylpiperazin-1-ium cation while the reported compounds (I)–(IV) have no substituent. They also crystallize as monohydrates, and their crystal structures are based on differently sized chains of rings formed *via* a combination of hydrogen bonds of the type N—H...O and O—H...O and other weak interactions of types C—H...O and C—H... π to form sheets. In 4-(4-methoxyphenyl)piperazin-1-ium 4-aminobenzoate monohydrate (IHIMEU; Kiran Kumar *et al.*, 2020a) the presence of the amino substituent in the anion, which acts as both a donor and as an

acceptor of hydrogen bonds, makes the supramolecular assembly of this compound more complex than those reported here. A search for 4-phenylpiperazin-1-ium and acetate derivatives involved in the reported compound (IV) gave no hits.

5. Synthesis and crystallization

For the synthesis of salts (I)–(IV), a solution of commercially available (from Sigma–Aldrich) 1-phenylpiperazine (100 mg, 0.62 mol) in methanol (10 ml) was mixed with equimolar solutions of the appropriate organic acids in methanol (10 ml) *viz.*, 4-ethoxybenzoic acid (103 mg, 0.62 mol) for (I), 4-methoxybenzoic acid (94 mg, 0.62 mol) for (II), 4-methylbenzoic acid (84 mg, 0.62 mol) for (III) and trifluoroacetic acid (71 mg, 0.62 mol) for (IV). The corresponding solutions were stirred for 15 min at room temperature and allowed to stand at the same temperature. X-ray quality crystals were formed on slow evaporation in a week for all compounds, where ethanol:ethylacetate (1:1) was used for crystallization. The corresponding melting points were 353–355 K for (I), 368–370 K for (II), 338–340 K for (III) and 385–387 K for (IV).

6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 5. All H atoms bonded to C atoms were fixed geometrically and treated as riding with C—H = 0.93 Å (aromatic), C—H = 0.96 Å (methyl) or 0.97 Å (methylene), with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{C})$. For the H

Table 5
Experimental details.

	(I)	(II)	(III)	(IV)
Crystal data				
Chemical formula	C ₁₀ H ₁₅ N ₂ ⁺ ·C ₉ H ₉ O ₃ ⁻ ·H ₂ O	C ₁₀ H ₁₅ N ₂ ⁺ ·C ₈ H ₇ O ₃ ⁻ ·H ₂ O	C ₁₀ H ₁₅ N ₂ ⁺ ·C ₈ H ₇ O ₂ ⁻ ·H ₂ O	C ₁₀ H ₁₅ N ₂ ⁺ ·C ₂ F ₃ O ₂ ⁻ ·0.123H ₂ O
<i>M_r</i>	346.42	332.39	316.39	278.47
Crystal system, space group	Triclinic, <i>P</i> $\bar{1}$	Triclinic, <i>P</i> $\bar{1}$	Triclinic, <i>P</i> $\bar{1}$	Triclinic, <i>P</i> $\bar{1}$
Temperature (K)	293	293	293	293
<i>a</i> , <i>b</i> , <i>c</i> (Å)	6.1635 (5), 7.5946 (6), 20.458 (2)	6.2039 (4), 7.5565 (7), 18.614 (1)	6.1175 (5), 7.6225 (7), 18.452 (1)	9.6544 (6), 9.9029 (6), 15.2090 (9)
α , β , γ (°)	79.545 (7), 86.521 (7), 83.791 (7)	81.799 (7), 87.020 (7), 84.852 (7)	97.421 (9), 90.403 (8), 92.405 (8)	79.621 (6), 86.579 (6), 70.603 (6)
<i>V</i> (Å ³)	935.38 (14)	859.53 (11)	852.40 (12)	1349.10 (15)
<i>Z</i>	2	2	2	4
Radiation type	Mo <i>K</i> α	Mo <i>K</i> α	Mo <i>K</i> α	Mo <i>K</i> α
μ (mm ⁻¹)	0.09	0.09	0.08	0.12
Crystal size (mm)	0.48 × 0.42 × 0.1	0.48 × 0.48 × 0.32	0.5 × 0.4 × 0.08	0.48 × 0.48 × 0.36
Data collection				
Diffractometer	Oxford Diffraction Xcalibur	Oxford Diffraction Xcalibur	Oxford Diffraction Xcalibur	Oxford Diffraction Xcalibur
Absorption correction	Multi-scan (<i>CrysAlis RED</i> ; Oxford Diffraction, 2009)	Multi-scan (<i>CrysAlis RED</i> ; Oxford Diffraction, 2009)	Multi-scan (<i>CrysAlis RED</i> ; Oxford Diffraction, 2009)	Multi-scan (<i>CrysAlis RED</i> ; Oxford Diffraction, 2009)
<i>T</i> _{min} , <i>T</i> _{max}	0.623, 1.000	0.520, 1.000	0.837, 1.000	0.724, 1.000
No. of measured, independent and observed [<i>I</i> > 2 σ (<i>I</i>)] reflections	5989, 3429, 2159	5360, 3142, 2322	5354, 3126, 2248	9220, 4940, 2777
<i>R</i> _{int}	0.022	0.016	0.013	0.014
(<i>sin</i> θ / λ) _{max} (Å ⁻¹)	0.602	0.602	0.602	0.602
Refinement				
<i>R</i> [<i>F</i> ² > 2 σ (<i>F</i> ²)], <i>wR</i> (<i>F</i> ²), <i>S</i>	0.054, 0.124, 1.08	0.045, 0.125, 1.06	0.046, 0.128, 1.03	0.070, 0.235, 1.07
No. of reflections	3424	3139	3118	4927
No. of parameters	244	230	226	375
No. of restraints	2	4	4	4
H-atom treatment	H atoms treated by a mixture of independent and constrained refinement	H atoms treated by a mixture of independent and constrained refinement	H atoms treated by a mixture of independent and constrained refinement	H-atom parameters constrained
$\Delta\rho_{\max}$, $\Delta\rho_{\min}$ (e Å ⁻³)	0.15, -0.15	0.2, -0.17	0.16, -0.16	0.42, -0.28

Computer programs: *CrysAlis CCD* (Oxford Diffraction, 2009), *CrysAlis RED* (Oxford Diffraction, 2009), *SHELXT* (Sheldrick, 2015a), *SHELXL2014* (Sheldrick, 2015b), *Mercury* (Macrae et al., 2020), *PLATON* (Spek, 2020) and *publCIF* (Westrip, 2010).

atoms bonded to the N and O atoms, the atomic coordinates were refined with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$ and $1.5U_{\text{eq}}(\text{O})$, [for (I), N2–HN2 = 0.931 (19), N2–HN1 = 0.888 (17) Å and OW1–HW2 = 0.91 (3), OW1–HW1 = 0.88 (3) Å; for (II), N2–HN1 = 0.927 (16), N2–HN2 = 0.931 (18) Å and OW–HW1 = 0.840 (19), OW1–HW2 = 0.85 (2) Å; for (III), N2–HN1 = 0.900 (16), N2–HN2 = 0.918 (17) Å and OW1–HW1 = 0.89 (3), OW1–HW2 = 0.84 (2) Å and for (IV), N2–H22 = 0.87 (2) and N2–H21 = 0.88 (3) Å]. In (IV), the atoms of the CF₃ groups of two trifluoroacetate anions (B1, B2) are disordered over two sets of sites with site occupancies of 0.737 (3) and 0.263 (3). The corresponding bond distances in the disordered groups were restrained to be equal. The U^{ij} components of these atoms were restrained to be equal and were restrained to approximate isotropic behaviour. The OW1 water molecule was refined with a resulting occupation factor of 0.245 (10) and the H atoms of the water molecule were placed geometrically.

Acknowledgements

NM is grateful to the University of Mysore for research facilities. HSY thanks the UGC for a BSR Faculty fellowship

for three years. SGG gratefully acknowledges financial support from the Spanish Ministerio de Ciencia e Innovación (PID2020–113558RB-C41) and Gobierno del Principado de Asturias (AYUD/2021/50997).

Funding information

Funding for this research was provided by: Ministerio de Ciencia e Innovación (grant No. PID2020-113558RB-C41 to S. Garcia-Granda, M. S. M. Abdelbaky); Gobierno del Principado de Asturias (grant No. AYUD/2021/50997 to S. Garcia-Granda, M. S. M. Abdelbaky).

References

- Al-Alshaiikh, M. A., El-Emam, A. A., Al-Deeb, O. A., Abdelbaky, M. S. M. & Garcia-Granda, S. (2015). *Acta Cryst.* E71, 956–959.
- Bogatcheva, E., Hanrahan, C., Nikonenko, B., Samala, R., Chen, P., Gearhart, J., Barbosa, F., Einck, L., Nacy, C. A. & Protopopova, M. (2006). *J. Med. Chem.* 49, 3045–3048.
- Brockunier, L. L., He, J., Colwell, L. F. Jr, Habulihaz, B., He, H., Leiting, B., Lyons, K. A., Marsilio, F., Patel, R. A., Teffera, Y., Wu, J. K., Thornberry, N. A., Weber, A. E. & Parmee, E. R. (2004). *Bioorg. Med. Chem. Lett.* 14, 4763–4766.
- Cohen, M. R., Hinsch, E., Palkoski, Z., Vergona, R., Urbano, S. & Sztokalo, J. (1982). *J. Pharmacol. Exp. Ther.* 223, 110–119.

- Conrado, D. J., Verli, H., Neves, G., Fraga, C. A., Barreiro, E. J., Rates, S. M. & Dalla-Costa, T. (2010). *J. Pharm. Pharmacol.* **60**, 699–707.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Demirci, S., Hayal, T. B., Kıratlı, B., Şişli, H. B., Demirci, S., Şahin, F. & Doğan, A. (2019). *Chem. Biol. Drug Des.* **94**, 1584–1595.
- Elliott, S. (2011). *Drug Test Anal.* **3**, 430–438.
- Essid, M., Marouani, H., Rzaigui, M. & Al-Deyab, S. S. (2010). *Acta Cryst.* **E66**, o2244–o2245.
- Groom, C. R., Bruno, I. J., Lightfoot, M. P. & Ward, S. C. (2016). *Acta Cryst.* **B72**, 171–179.
- Hanano, T., Adachi, K., Aoki, Y., Morimoto, H., Naka, Y., Hisadome, M., Fukuda, T. & Sumichika, H. (2000). *Bioorg. Med. Chem. Lett.* **10**, 875–879.
- Harish Chinthala, C., Kavitha, C. N., Yathirajan, H. S., Foro, S., Rathore, R. S. & Glidewell, C. (2020). *Acta Cryst.* **E76**, 1779–1793.
- Kiran Kumar, H., Yathirajan, H. S., Foro, S. & Glidewell, C. (2019a). *Acta Cryst.* **E75**, 1494–1506.
- Kiran Kumar, H., Yathirajan, H. S., Harish Chinthala, C., Foro, S. & Glidewell, C. (2020a). *Acta Cryst.* **E76**, 488–495.
- Kiran Kumar, H., Yathirajan, H. S., Harish Chinthala, C., Foro, S. & Glidewell, C. (2020b). CSD Communication (refcode KUJPUJ). CCDC, Cambridge, England.
- Kiran Kumar, H., Yathirajan, H. S., Sagar, B. K., Foro, S. & Glidewell, C. (2019b). *Acta Cryst.* **E75**, 1253–1260.
- Li, A.-J., Zhang, X.-H., Sun, W.-Q., Zhou, X.-Q. & Liu, D.-Z. (2008). *Acta Cryst.* **E64**, o1234.
- Macrae, C. F., Sovago, I., Cottrell, S. J., Galek, P. T. A., McCabe, P., Pidcock, E., Platings, M., Shields, G. P., Stevens, J. S., Towler, M. & Wood, P. A. (2020). *J. Appl. Cryst.* **53**, 226–235.
- Marouani, H., Rzaigui, M. & Al-Deyab, S. S. (2010). *Acta Cryst.* **E66**, o2613.
- Neves, G., Fenner, R., Heckler, A. P., Viana, A. F., Tasso, L., Menegatti, R., Fraga, C. A. M., Barreiro, E. J., Dalla-Costa, T. & Rates, S. M. K. (2003). *Braz. J. Med. Biol. Res.* **36**, 625–629.
- Oxford Diffraction (2009). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Abingdon, England.
- Sheldrick, G. M. (2015a). *Acta Cryst.* **A71**, 3–8.
- Sheldrick, G. M. (2015b). *Acta Cryst.* **C71**, 3–8.
- Spek, A. L. (2020). *Acta Cryst.* **E76**, 1–11.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Xu, Y.-J. & Jing, F. (2009). *Acta Cryst.* **E65**, o1211.

supporting information

Acta Cryst. (2022). E78, 709-715 [https://doi.org/10.1107/S2056989022006004]

Crystal-structure studies of 4-phenylpiperazin-1-ium 4-ethoxybenzoate monohydrate, 4-phenylpiperazin-1-ium 4-methoxybenzoate monohydrate, 4-phenylpiperazin-1-ium 4-methylbenzoate monohydrate and 4-phenylpiperazin-1-ium trifluoroacetate 0.12-hydrate

Ninganayaka Mahesha, Haruvegowda Kiran Kumar, Mehmet Akkurt, Hemmige S. Yathirajan, Sabine Foro, Mohammed S. M. Abdelbaky and Santiago Garcia-Granda

Computing details

For all structures, data collection: *CrysAlis CCD* (Oxford Diffraction, 2009); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2009); data reduction: *CrysAlis RED* (Oxford Diffraction, 2009); program(s) used to solve structure: SHELXT (Sheldrick, 2015a); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015b); molecular graphics: *Mercury* (Macrae *et al.*, 2020); software used to prepare material for publication: *SHELXL2014* (Sheldrick, 2015b), *PLATON* (Spek, 2020) and *publCIF* (Westrip, 2010).

4-Phenylpiperazin-1-ium 4-ethoxybenzoate monohydrate (I)

Crystal data

$C_{10}H_{15}N_2^+ \cdot C_9H_9O_3^- \cdot H_2O$

$M_r = 346.42$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 6.1635$ (5) Å

$b = 7.5946$ (6) Å

$c = 20.458$ (2) Å

$\alpha = 79.545$ (7)°

$\beta = 86.521$ (7)°

$\gamma = 83.791$ (7)°

$V = 935.38$ (14) Å³

$Z = 2$

$F(000) = 372$

$D_x = 1.23$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2352 reflections

$\theta = 3.0$ – 27.8 °

$\mu = 0.09$ mm⁻¹

$T = 293$ K

Plate, colourless

$0.48 \times 0.42 \times 0.1$ mm

Data collection

Oxford Diffraction Xcalibur
diffractometer

ω scans

Absorption correction: multi-scan
(*CrysAlis RED*; Oxford Diffraction, 2009)

$T_{\min} = 0.623$, $T_{\max} = 1.000$

5989 measured reflections

3429 independent reflections

2159 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.022$

$\theta_{\text{max}} = 25.4$ °, $\theta_{\text{min}} = 3.0$ °

$h = -7 \rightarrow 7$

$k = -9 \rightarrow 9$

$l = -24 \rightarrow 14$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.054$

$wR(F^2) = 0.124$

$S = 1.08$

3424 reflections

244 parameters

2 restraints

0 constraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: structure-
invariant direct methods

Hydrogen site location: mixed

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0392P)^2 + 0.2859P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.15 \text{ e } \text{Å}^{-3}$

$\Delta\rho_{\min} = -0.15 \text{ e } \text{Å}^{-3}$

Extinction correction: SHELXL2018/3
(Sheldrick 2015b),

$F_c^* = kFc[1 + 0.001x\text{Fc}^2\lambda^3/\sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0058 (17)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.1990 (4)	0.7508 (4)	0.79338 (12)	0.0626 (7)
H1	0.069252	0.809989	0.776125	0.075*
C2	0.2437 (5)	0.7539 (4)	0.85869 (13)	0.0782 (9)
H2	0.142853	0.814295	0.884641	0.094*
C3	0.4331 (5)	0.6699 (4)	0.88559 (14)	0.0759 (9)
H3	0.462163	0.672073	0.929504	0.091*
C4	0.5786 (5)	0.5827 (4)	0.84658 (14)	0.0753 (8)
H4	0.709095	0.525999	0.864083	0.09*
C5	0.5363 (4)	0.5771 (3)	0.78183 (12)	0.0579 (7)
H5	0.638106	0.515837	0.756445	0.069*
C6	0.3432 (3)	0.6615 (3)	0.75355 (10)	0.0407 (5)
C7	0.1046 (3)	0.7635 (3)	0.66062 (11)	0.0484 (6)
H7A	0.130663	0.888733	0.656771	0.058*
H7B	-0.01969	0.742138	0.691398	0.058*
C8	0.0502 (3)	0.7292 (3)	0.59343 (11)	0.0496 (6)
H8A	0.005085	0.609001	0.59798	0.059*
H8B	-0.07032	0.814458	0.576145	0.059*
C9	0.4290 (3)	0.6216 (3)	0.57366 (10)	0.0480 (6)
H9A	0.555062	0.634835	0.543053	0.058*
H9B	0.392361	0.498713	0.578625	0.058*
C10	0.4839 (3)	0.6594 (3)	0.64012 (10)	0.0447 (6)
H10A	0.602615	0.572985	0.657958	0.054*
H10B	0.5333	0.778458	0.634186	0.054*
C11	0.2356 (4)	0.7988 (3)	0.37750 (13)	0.0504 (6)
C12	0.1740 (3)	0.8093 (3)	0.30715 (11)	0.0409 (5)
C13	-0.0108 (3)	0.7381 (3)	0.29259 (11)	0.0462 (6)

H13	-0.09987	0.685018	0.327326	0.055*
C14	-0.0669 (4)	0.7436 (3)	0.22762 (11)	0.0496 (6)
H14	-0.191724	0.694349	0.218988	0.06*
C15	0.0633 (4)	0.8223 (3)	0.17610 (11)	0.0480 (6)
C16	0.2462 (4)	0.8988 (3)	0.18948 (12)	0.0547 (6)
H16	0.332264	0.955074	0.154649	0.066*
C17	0.3005 (4)	0.8916 (3)	0.25403 (12)	0.0521 (6)
H17	0.424178	0.942678	0.262421	0.063*
C18	-0.1614 (5)	0.7562 (5)	0.09410 (14)	0.0931 (10)
H18A	-0.293805	0.818359	0.110377	0.112*
H18B	-0.155291	0.63023	0.114607	0.112*
C19	-0.1605 (8)	0.7747 (7)	0.02032 (17)	0.178 (2)
H19A	-0.195823	0.898767	0.00101	0.268*
H19B	-0.266871	0.703796	0.008494	0.268*
H19C	-0.018288	0.73364	0.003915	0.268*
N1	0.2969 (3)	0.6493 (2)	0.68767 (8)	0.0382 (4)
N2	0.2420 (3)	0.7474 (3)	0.54621 (10)	0.0452 (5)
O1	0.1211 (3)	0.7133 (2)	0.42359 (8)	0.0622 (5)
O2	0.3970 (3)	0.8727 (3)	0.38724 (10)	0.0885 (7)
O3	0.0253 (3)	0.8330 (2)	0.11014 (8)	0.0681 (5)
OW1	0.7230 (4)	0.8819 (2)	0.46338 (9)	0.0591 (5)
HN1	0.275 (4)	0.860 (2)	0.5421 (11)	0.056 (7)*
HN2	0.208 (4)	0.729 (3)	0.5044 (9)	0.067 (8)*
HW1	0.612 (5)	0.879 (4)	0.4385 (15)	0.102 (11)*
HW2	0.842 (5)	0.825 (4)	0.4445 (15)	0.102 (12)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0719 (17)	0.0673 (18)	0.0465 (15)	0.0151 (14)	-0.0066 (13)	-0.0156 (13)
C2	0.103 (2)	0.079 (2)	0.0522 (17)	0.0150 (18)	-0.0020 (16)	-0.0248 (15)
C3	0.106 (2)	0.076 (2)	0.0510 (17)	-0.0055 (18)	-0.0223 (17)	-0.0207 (15)
C4	0.0735 (18)	0.090 (2)	0.0636 (18)	0.0027 (16)	-0.0267 (15)	-0.0145 (17)
C5	0.0561 (15)	0.0696 (18)	0.0486 (15)	0.0031 (13)	-0.0116 (12)	-0.0147 (13)
C6	0.0467 (12)	0.0353 (13)	0.0414 (13)	-0.0092 (10)	-0.0045 (10)	-0.0067 (10)
C7	0.0399 (12)	0.0595 (16)	0.0441 (13)	0.0007 (11)	-0.0025 (10)	-0.0078 (11)
C8	0.0407 (12)	0.0622 (16)	0.0445 (14)	-0.0045 (11)	-0.0053 (10)	-0.0053 (11)
C9	0.0480 (13)	0.0512 (15)	0.0423 (13)	-0.0011 (11)	0.0026 (10)	-0.0057 (11)
C10	0.0396 (12)	0.0496 (14)	0.0435 (13)	-0.0024 (10)	-0.0018 (10)	-0.0062 (11)
C11	0.0524 (14)	0.0427 (14)	0.0593 (17)	0.0050 (12)	-0.0162 (13)	-0.0190 (12)
C12	0.0398 (12)	0.0373 (13)	0.0475 (13)	-0.0007 (10)	-0.0072 (10)	-0.0121 (10)
C13	0.0494 (13)	0.0462 (14)	0.0432 (13)	-0.0111 (11)	-0.0024 (11)	-0.0048 (11)
C14	0.0502 (13)	0.0519 (15)	0.0490 (14)	-0.0163 (11)	-0.0091 (11)	-0.0063 (12)
C15	0.0556 (14)	0.0469 (15)	0.0412 (14)	-0.0042 (11)	-0.0052 (11)	-0.0065 (11)
C16	0.0534 (14)	0.0566 (16)	0.0533 (16)	-0.0136 (12)	0.0088 (12)	-0.0063 (12)
C17	0.0409 (12)	0.0533 (16)	0.0659 (17)	-0.0098 (11)	-0.0026 (12)	-0.0174 (13)
C18	0.111 (2)	0.121 (3)	0.0572 (18)	-0.046 (2)	-0.0218 (17)	-0.0148 (18)
C19	0.242 (6)	0.251 (6)	0.063 (2)	-0.137 (5)	-0.039 (3)	-0.007 (3)

N1	0.0374 (9)	0.0419 (11)	0.0350 (10)	-0.0028 (8)	-0.0021 (8)	-0.0062 (8)
N2	0.0548 (12)	0.0428 (13)	0.0390 (11)	-0.0088 (10)	-0.0079 (9)	-0.0058 (9)
O1	0.0706 (11)	0.0703 (12)	0.0465 (10)	-0.0048 (10)	-0.0127 (9)	-0.0108 (9)
O2	0.0835 (13)	0.1109 (17)	0.0825 (14)	-0.0368 (12)	-0.0321 (11)	-0.0215 (12)
O3	0.0816 (12)	0.0792 (13)	0.0450 (10)	-0.0182 (10)	-0.0052 (9)	-0.0080 (9)
OW1	0.0593 (11)	0.0598 (12)	0.0604 (12)	-0.0122 (10)	-0.0135 (10)	-0.0092 (9)

Geometric parameters (Å, °)

C1—C6	1.377 (3)	C11—O2	1.237 (3)
C1—C2	1.385 (3)	C11—O1	1.267 (3)
C1—H1	0.93	C11—C12	1.497 (3)
C2—C3	1.365 (4)	C12—C13	1.383 (3)
C2—H2	0.93	C12—C17	1.391 (3)
C3—C4	1.362 (4)	C13—C14	1.386 (3)
C3—H3	0.93	C13—H13	0.93
C4—C5	1.375 (3)	C14—C15	1.373 (3)
C4—H4	0.93	C14—H14	0.93
C5—C6	1.395 (3)	C15—O3	1.370 (3)
C5—H5	0.93	C15—C16	1.386 (3)
C6—N1	1.416 (3)	C16—C17	1.372 (3)
C7—N1	1.465 (2)	C16—H16	0.93
C7—C8	1.508 (3)	C17—H17	0.93
C7—H7A	0.97	C18—O3	1.425 (3)
C7—H7B	0.97	C18—C19	1.490 (4)
C8—N2	1.482 (3)	C18—H18A	0.97
C8—H8A	0.97	C18—H18B	0.97
C8—H8B	0.97	C19—H19A	0.96
C9—N2	1.484 (3)	C19—H19B	0.96
C9—C10	1.504 (3)	C19—H19C	0.96
C9—H9A	0.97	N2—HN1	0.887 (16)
C9—H9B	0.97	N2—HN2	0.930 (16)
C10—N1	1.462 (3)	OW1—HW1	0.88 (3)
C10—H10A	0.97	OW1—HW2	0.91 (3)
C10—H10B	0.97		
C6—C1—C2	121.2 (2)	O2—C11—C12	118.0 (2)
C6—C1—H1	119.4	O1—C11—C12	118.2 (2)
C2—C1—H1	119.4	C13—C12—C17	117.6 (2)
C3—C2—C1	121.1 (3)	C13—C12—C11	121.3 (2)
C3—C2—H2	119.4	C17—C12—C11	121.1 (2)
C1—C2—H2	119.4	C12—C13—C14	121.8 (2)
C4—C3—C2	118.4 (3)	C12—C13—H13	119.1
C4—C3—H3	120.8	C14—C13—H13	119.1
C2—C3—H3	120.8	C15—C14—C13	119.4 (2)
C3—C4—C5	121.3 (3)	C15—C14—H14	120.3
C3—C4—H4	119.4	C13—C14—H14	120.3
C5—C4—H4	119.4	O3—C15—C14	124.4 (2)

C4—C5—C6	121.1 (2)	O3—C15—C16	115.8 (2)
C4—C5—H5	119.4	C14—C15—C16	119.8 (2)
C6—C5—H5	119.4	C17—C16—C15	120.1 (2)
C1—C6—C5	116.9 (2)	C17—C16—H16	119.9
C1—C6—N1	122.23 (19)	C15—C16—H16	119.9
C5—C6—N1	120.8 (2)	C16—C17—C12	121.2 (2)
N1—C7—C8	112.74 (18)	C16—C17—H17	119.4
N1—C7—H7A	109	C12—C17—H17	119.4
C8—C7—H7A	109	O3—C18—C19	107.7 (3)
N1—C7—H7B	109	O3—C18—H18A	110.2
C8—C7—H7B	109	C19—C18—H18A	110.2
H7A—C7—H7B	107.8	O3—C18—H18B	110.2
N2—C8—C7	110.70 (17)	C19—C18—H18B	110.2
N2—C8—H8A	109.5	H18A—C18—H18B	108.5
C7—C8—H8A	109.5	C18—C19—H19A	109.5
N2—C8—H8B	109.5	C18—C19—H19B	109.5
C7—C8—H8B	109.5	H19A—C19—H19B	109.5
H8A—C8—H8B	108.1	C18—C19—H19C	109.5
N2—C9—C10	110.47 (18)	H19A—C19—H19C	109.5
N2—C9—H9A	109.6	H19B—C19—H19C	109.5
C10—C9—H9A	109.6	C6—N1—C10	115.21 (16)
N2—C9—H9B	109.6	C6—N1—C7	115.50 (17)
C10—C9—H9B	109.6	C10—N1—C7	111.72 (16)
H9A—C9—H9B	108.1	C8—N2—C9	109.60 (17)
N1—C10—C9	112.16 (17)	C8—N2—HN1	106.9 (15)
N1—C10—H10A	109.2	C9—N2—HN1	109.4 (15)
C9—C10—H10A	109.2	C8—N2—HN2	110.8 (15)
N1—C10—H10B	109.2	C9—N2—HN2	112.2 (15)
C9—C10—H10B	109.2	HN1—N2—HN2	108 (2)
H10A—C10—H10B	107.9	C15—O3—C18	117.7 (2)
O2—C11—O1	123.7 (2)	HW1—OW1—HW2	107 (3)
C6—C1—C2—C3	0.6 (4)	O3—C15—C16—C17	-178.7 (2)
C1—C2—C3—C4	0.2 (5)	C14—C15—C16—C17	1.8 (4)
C2—C3—C4—C5	-0.7 (5)	C15—C16—C17—C12	-0.4 (4)
C3—C4—C5—C6	0.5 (4)	C13—C12—C17—C16	-1.2 (3)
C2—C1—C6—C5	-0.8 (4)	C11—C12—C17—C16	178.9 (2)
C2—C1—C6—N1	177.0 (2)	C1—C6—N1—C10	143.0 (2)
C4—C5—C6—C1	0.3 (4)	C5—C6—N1—C10	-39.3 (3)
C4—C5—C6—N1	-177.6 (2)	C1—C6—N1—C7	10.3 (3)
N1—C7—C8—N2	-54.6 (3)	C5—C6—N1—C7	-171.9 (2)
N2—C9—C10—N1	56.7 (2)	C9—C10—N1—C6	172.59 (17)
O2—C11—C12—C13	-176.6 (2)	C9—C10—N1—C7	-53.0 (2)
O1—C11—C12—C13	4.1 (3)	C8—C7—N1—C6	-173.71 (18)
O2—C11—C12—C17	3.3 (3)	C8—C7—N1—C10	52.0 (2)
O1—C11—C12—C17	-176.0 (2)	C7—C8—N2—C9	57.2 (2)
C17—C12—C13—C14	1.6 (3)	C10—C9—N2—C8	-58.4 (2)
C11—C12—C13—C14	-178.6 (2)	C14—C15—O3—C18	-0.4 (4)

C12—C13—C14—C15	−0.3 (3)	C16—C15—O3—C18	−179.9 (2)
C13—C14—C15—O3	179.1 (2)	C19—C18—O3—C15	−176.7 (3)
C13—C14—C15—C16	−1.4 (3)		

Hydrogen-bond geometry (Å, °)

*Cg*1 and *Cg*3 are the centroids of the C12–C17 and C1–C6 benzene rings, respectively.

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N2—HN1...OW1 ⁱ	0.89 (2)	1.94 (2)	2.817 (3)	167 (2)
N2—HN2...O1	0.93 (2)	1.80 (2)	2.724 (3)	174 (2)
OW1—HW1...O2	0.88 (3)	1.75 (3)	2.630 (3)	178 (4)
OW1—HW2...O1 ⁱⁱ	0.91 (3)	1.89 (3)	2.789 (3)	167 (3)
C9—H9A...OW1	0.97	2.52	3.308 (3)	138
C1—H1... <i>Cg</i> 1 ⁱⁱⁱ	0.93	2.91	3.607 (3)	133
C5—H5... <i>Cg</i> 1 ^{iv}	0.93	2.79	3.570 (3)	142
C18—H18B... <i>Cg</i> 3 ^v	0.97	2.88	3.737 (4)	148

Symmetry codes: (i) $-x+1, -y+2, -z+1$; (ii) $x+1, y, z$; (iii) $-x, -y+2, -z+1$; (iv) $-x+1, -y+1, -z+1$; (v) $-x, -y+1, -z+1$.

4-Phenylpiperazin-1-ium 4-methoxybenzoate monohydrate (II)*Crystal data*

$C_{10}H_{15}N_2^+ \cdot C_8H_7O_3^- \cdot H_2O$

$M_r = 332.39$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 6.2039$ (4) Å

$b = 7.5565$ (7) Å

$c = 18.614$ (1) Å

$\alpha = 81.799$ (7)°

$\beta = 87.020$ (7)°

$\gamma = 84.852$ (7)°

$V = 859.53$ (11) Å³

$Z = 2$

$F(000) = 356$

$D_x = 1.284$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2855 reflections

$\theta = 3.1$ – 27.8 °

$\mu = 0.09$ mm⁻¹

$T = 293$ K

Prism, colourless

$0.48 \times 0.48 \times 0.32$ mm

Data collection

Oxford Diffraction Xcalibur
diffractometer

ω scans

Absorption correction: multi-scan

(CrysAlis RED; Oxford Diffraction, 2009)

$T_{\min} = 0.520$, $T_{\max} = 1.000$

5360 measured reflections

3142 independent reflections

2322 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.016$

$\theta_{\max} = 25.3$ °, $\theta_{\min} = 3.1$ °

$h = -7 \rightarrow 5$

$k = -9 \rightarrow 9$

$l = -22 \rightarrow 22$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.045$

$wR(F^2) = 0.125$

$S = 1.06$

3139 reflections

230 parameters

4 restraints

0 constraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: structure-
invariant direct methods

Hydrogen site location: mixed

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0613P)^2 + 0.1503P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$$\Delta\rho_{\max} = 0.2 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.16 \text{ e } \text{\AA}^{-3}$$

Extinction correction: SHELXL2018/3
(Sheldrick 2015b),
 $F_c^* = kFc[1 + 0.001x\text{Fc}^2\lambda^3/\sin(2\theta)]^{-1/4}$
Extinction coefficient: 0.032 (4)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	-0.0254 (3)	0.4086 (3)	0.19209 (10)	0.0516 (5)
H1	-0.120858	0.476382	0.219629	0.062*
C2	-0.0695 (3)	0.3996 (3)	0.12102 (11)	0.0625 (6)
H2	-0.194898	0.46027	0.101533	0.075*
C3	0.0686 (3)	0.3024 (3)	0.07854 (10)	0.0597 (6)
H3	0.039957	0.298402	0.030279	0.072*
C4	0.2497 (3)	0.2117 (3)	0.10901 (10)	0.0593 (6)
H4	0.343702	0.143909	0.08102	0.071*
C5	0.2964 (3)	0.2181 (3)	0.18024 (9)	0.0512 (5)
H5	0.420732	0.154912	0.199433	0.061*
C6	0.1594 (2)	0.3180 (2)	0.22353 (8)	0.0361 (4)
C7	0.0222 (3)	0.3258 (2)	0.34797 (8)	0.0419 (4)
H7A	-0.026461	0.205915	0.354319	0.05*
H7B	-0.096084	0.40906	0.32889	0.05*
C8	0.0795 (3)	0.3708 (2)	0.42029 (9)	0.0456 (4)
H8A	0.116889	0.494094	0.414858	0.055*
H8B	-0.044531	0.359357	0.454053	0.055*
C9	0.4547 (3)	0.2652 (3)	0.39777 (9)	0.0485 (5)
H9A	0.574598	0.182967	0.416336	0.058*
H9B	0.499661	0.386146	0.392734	0.058*
C10	0.3979 (3)	0.2229 (3)	0.32457 (9)	0.0445 (4)
H10A	0.520581	0.241283	0.29068	0.053*
H10B	0.36928	0.097655	0.328948	0.053*
C11	0.2655 (3)	0.2146 (2)	0.63263 (10)	0.0474 (5)
C12	0.3208 (3)	0.2073 (2)	0.71052 (9)	0.0389 (4)
C13	0.1897 (3)	0.1251 (2)	0.76647 (10)	0.0479 (5)
H13	0.067829	0.072733	0.755273	0.057*
C14	0.2386 (3)	0.1207 (3)	0.83782 (10)	0.0496 (5)
H14	0.149875	0.065178	0.874461	0.06*
C15	0.4186 (3)	0.1982 (2)	0.85567 (9)	0.0442 (4)
C16	0.5528 (3)	0.2784 (2)	0.80102 (9)	0.0453 (4)
H16	0.675183	0.329713	0.812349	0.054*
C17	0.5021 (3)	0.2808 (2)	0.72949 (9)	0.0424 (4)
H17	0.592994	0.333766	0.6929	0.051*
C18	0.6395 (4)	0.2628 (4)	0.94845 (12)	0.0858 (8)

H13A	0.643815	0.248303	1.000449	0.129*
H13B	0.76663	0.201465	0.928951	0.129*
H13C	0.63455	0.388125	0.929616	0.129*
N1	0.2079 (2)	0.33487 (18)	0.29565 (7)	0.0359 (3)
N2	0.2652 (2)	0.2489 (2)	0.44963 (8)	0.0434 (4)
O1	0.1058 (3)	0.1395 (3)	0.61995 (9)	0.0873 (6)
O2	0.3847 (2)	0.29717 (19)	0.58441 (7)	0.0598 (4)
O3	0.4517 (2)	0.1895 (2)	0.92816 (7)	0.0620 (4)
OW1	0.2271 (2)	0.8777 (2)	0.46022 (8)	0.0579 (4)
HN1	0.230 (3)	0.131 (2)	0.4565 (11)	0.07*
HW1	0.120 (3)	0.865 (3)	0.4356 (11)	0.07*
HN2	0.303 (3)	0.270 (3)	0.4956 (9)	0.07*
HW2	0.336 (3)	0.822 (3)	0.4413 (12)	0.07*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0534 (11)	0.0579 (12)	0.0441 (10)	0.0127 (9)	-0.0106 (8)	-0.0169 (9)
C2	0.0654 (13)	0.0739 (14)	0.0491 (11)	0.0143 (11)	-0.0228 (10)	-0.0166 (10)
C3	0.0820 (14)	0.0654 (13)	0.0347 (10)	-0.0036 (11)	-0.0147 (10)	-0.0145 (9)
C4	0.0717 (13)	0.0660 (13)	0.0411 (10)	0.0070 (11)	0.0022 (9)	-0.0202 (9)
C5	0.0535 (11)	0.0597 (12)	0.0399 (10)	0.0105 (9)	-0.0032 (8)	-0.0137 (8)
C6	0.0405 (9)	0.0344 (9)	0.0346 (8)	-0.0042 (7)	-0.0035 (7)	-0.0071 (7)
C7	0.0388 (9)	0.0515 (11)	0.0352 (9)	0.0009 (8)	-0.0006 (7)	-0.0083 (7)
C8	0.0518 (10)	0.0495 (11)	0.0354 (9)	0.0016 (8)	0.0004 (8)	-0.0104 (8)
C9	0.0425 (10)	0.0619 (12)	0.0419 (10)	-0.0015 (8)	-0.0105 (8)	-0.0090 (8)
C10	0.0385 (9)	0.0569 (11)	0.0381 (9)	0.0049 (8)	-0.0053 (7)	-0.0113 (8)
C11	0.0484 (11)	0.0464 (11)	0.0508 (11)	0.0051 (8)	-0.0161 (9)	-0.0190 (9)
C12	0.0387 (9)	0.0356 (9)	0.0443 (9)	0.0023 (7)	-0.0086 (7)	-0.0131 (7)
C13	0.0366 (9)	0.0485 (11)	0.0617 (12)	-0.0058 (8)	-0.0057 (8)	-0.0157 (9)
C14	0.0437 (10)	0.0551 (12)	0.0500 (11)	-0.0060 (8)	0.0047 (8)	-0.0079 (9)
C15	0.0450 (10)	0.0485 (10)	0.0398 (9)	0.0020 (8)	-0.0042 (8)	-0.0110 (8)
C16	0.0461 (10)	0.0501 (11)	0.0430 (10)	-0.0103 (8)	-0.0101 (8)	-0.0112 (8)
C17	0.0447 (10)	0.0437 (10)	0.0400 (9)	-0.0072 (8)	-0.0036 (7)	-0.0075 (7)
C18	0.0785 (16)	0.136 (2)	0.0509 (13)	-0.0228 (16)	-0.0171 (11)	-0.0274 (14)
N1	0.0347 (7)	0.0416 (8)	0.0320 (7)	0.0006 (6)	-0.0034 (5)	-0.0091 (6)
N2	0.0553 (9)	0.0450 (8)	0.0319 (7)	-0.0061 (7)	-0.0089 (6)	-0.0082 (6)
O1	0.0815 (11)	0.1184 (15)	0.0727 (11)	-0.0352 (10)	-0.0327 (9)	-0.0230 (10)
O2	0.0721 (9)	0.0700 (10)	0.0402 (7)	-0.0038 (7)	-0.0139 (7)	-0.0144 (7)
O3	0.0641 (9)	0.0866 (11)	0.0365 (7)	-0.0067 (8)	-0.0047 (6)	-0.0117 (7)
OW1	0.0595 (9)	0.0631 (9)	0.0544 (8)	-0.0097 (7)	-0.0177 (7)	-0.0108 (7)

Geometric parameters (Å, °)

C1—C2	1.377 (3)	C10—H10A	0.97
C1—C6	1.393 (2)	C10—H10B	0.97
C1—H1	0.93	C11—O1	1.234 (2)
C2—C3	1.371 (3)	C11—O2	1.263 (2)

C2—H2	0.93	C11—C12	1.499 (2)
C3—C4	1.368 (3)	C12—C17	1.381 (2)
C3—H3	0.93	C12—C13	1.396 (3)
C4—C5	1.380 (3)	C13—C14	1.373 (3)
C4—H4	0.93	C13—H13	0.93
C5—C6	1.388 (2)	C14—C15	1.383 (3)
C5—H5	0.93	C14—H14	0.93
C6—N1	1.4165 (19)	C15—O3	1.367 (2)
C7—N1	1.469 (2)	C15—C16	1.386 (2)
C7—C8	1.502 (2)	C16—C17	1.381 (2)
C7—H7A	0.97	C16—H16	0.93
C7—H7B	0.97	C17—H17	0.93
C8—N2	1.485 (2)	C18—O3	1.424 (3)
C8—H8A	0.97	C18—H13A	0.96
C8—H8B	0.97	C18—H13B	0.96
C9—N2	1.484 (2)	C18—H13C	0.96
C9—C10	1.509 (2)	N2—HN1	0.924 (15)
C9—H9A	0.97	N2—HN2	0.938 (16)
C9—H9B	0.97	OW1—HW1	0.847 (16)
C10—N1	1.467 (2)	OW1—HW2	0.850 (16)
C2—C1—C6	121.26 (16)	C9—C10—H10B	109.1
C2—C1—H1	119.4	H10A—C10—H10B	107.9
C6—C1—H1	119.4	O1—C11—O2	124.33 (18)
C3—C2—C1	121.01 (18)	O1—C11—C12	117.62 (19)
C3—C2—H2	119.5	O2—C11—C12	118.06 (16)
C1—C2—H2	119.5	C17—C12—C13	117.75 (16)
C4—C3—C2	118.25 (17)	C17—C12—C11	121.49 (16)
C4—C3—H3	120.9	C13—C12—C11	120.77 (16)
C2—C3—H3	120.9	C14—C13—C12	120.75 (16)
C3—C4—C5	121.67 (17)	C14—C13—H13	119.6
C3—C4—H4	119.2	C12—C13—H13	119.6
C5—C4—H4	119.2	C13—C14—C15	120.56 (17)
C4—C5—C6	120.65 (17)	C13—C14—H14	119.7
C4—C5—H5	119.7	C15—C14—H14	119.7
C6—C5—H5	119.7	O3—C15—C14	116.20 (16)
C5—C6—C1	117.15 (15)	O3—C15—C16	124.10 (16)
C5—C6—N1	122.11 (14)	C14—C15—C16	119.69 (16)
C1—C6—N1	120.68 (14)	C17—C16—C15	119.06 (16)
N1—C7—C8	111.59 (14)	C17—C16—H16	120.5
N1—C7—H7A	109.3	C15—C16—H16	120.5
C8—C7—H7A	109.3	C16—C17—C12	122.16 (16)
N1—C7—H7B	109.3	C16—C17—H17	118.9
C8—C7—H7B	109.3	C12—C17—H17	118.9
H7A—C7—H7B	108	O3—C18—H13A	109.5
N2—C8—C7	110.40 (13)	O3—C18—H13B	109.5
N2—C8—H8A	109.6	H13A—C18—H13B	109.5
C7—C8—H8A	109.6	O3—C18—H13C	109.5

N2—C8—H8B	109.6	H13A—C18—H13C	109.5
C7—C8—H8B	109.6	H13B—C18—H13C	109.5
H8A—C8—H8B	108.1	C6—N1—C10	115.62 (12)
N2—C9—C10	110.34 (14)	C6—N1—C7	114.90 (12)
N2—C9—H9A	109.6	C10—N1—C7	111.53 (12)
C10—C9—H9A	109.6	C9—N2—C8	109.67 (13)
N2—C9—H9B	109.6	C9—N2—HN1	108.7 (13)
C10—C9—H9B	109.6	C8—N2—HN1	110.6 (13)
H9A—C9—H9B	108.1	C9—N2—HN2	110.0 (13)
N1—C10—C9	112.30 (13)	C8—N2—HN2	113.2 (13)
N1—C10—H10A	109.1	HN1—N2—HN2	104.5 (18)
C9—C10—H10A	109.1	C15—O3—C18	117.68 (16)
N1—C10—H10B	109.1	HW1—OW1—HW2	106 (2)
C6—C1—C2—C3	-0.7 (3)	C13—C14—C15—C16	-1.1 (3)
C1—C2—C3—C4	1.2 (3)	O3—C15—C16—C17	-179.35 (16)
C2—C3—C4—C5	-0.9 (3)	C14—C15—C16—C17	0.7 (3)
C3—C4—C5—C6	0.0 (3)	C15—C16—C17—C12	0.5 (3)
C4—C5—C6—C1	0.6 (3)	C13—C12—C17—C16	-1.4 (2)
C4—C5—C6—N1	-176.76 (18)	C11—C12—C17—C16	178.68 (15)
C2—C1—C6—C5	-0.2 (3)	C5—C6—N1—C10	-7.1 (2)
C2—C1—C6—N1	177.14 (18)	C1—C6—N1—C10	175.65 (16)
N1—C7—C8—N2	-57.13 (19)	C5—C6—N1—C7	-139.28 (17)
N2—C9—C10—N1	55.4 (2)	C1—C6—N1—C7	43.5 (2)
O1—C11—C12—C17	177.14 (17)	C9—C10—N1—C6	172.93 (14)
O2—C11—C12—C17	-3.0 (2)	C9—C10—N1—C7	-53.34 (19)
O1—C11—C12—C13	-2.8 (2)	C8—C7—N1—C6	-171.86 (13)
O2—C11—C12—C13	177.07 (16)	C8—C7—N1—C10	54.06 (18)
C17—C12—C13—C14	1.0 (2)	C10—C9—N2—C8	-57.61 (19)
C11—C12—C13—C14	-179.05 (15)	C7—C8—N2—C9	58.76 (19)
C12—C13—C14—C15	0.2 (3)	C14—C15—O3—C18	178.00 (19)
C13—C14—C15—O3	178.99 (15)	C16—C15—O3—C18	-1.9 (3)

Hydrogen-bond geometry (Å, °)

*Cg*3 is the centroid of the C12—C17 benzene ring.

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N2—HN1...OW1 ⁱ	0.93 (2)	1.91 (2)	2.815 (2)	166 (2)
OW1—HW1...O1 ⁱⁱ	0.84 (2)	1.80 (2)	2.633 (2)	175 (2)
N2—HN2...O2	0.93 (2)	1.81 (2)	2.7350 (19)	176 (2)
OW1—HW2...O2 ⁱⁱⁱ	0.85 (2)	1.96 (2)	2.7876 (19)	168 (2)
C8—H8B...OW1 ⁱⁱ	0.97	2.53	3.331 (2)	140
C1—H1... <i>Cg</i> 3 ⁱⁱ	0.93	2.76	3.549 (2)	144
C5—H5... <i>Cg</i> 3 ^{iv}	0.93	2.86	3.625 (2)	140

Symmetry codes: (i) *x*, *y*-1, *z*; (ii) -*x*, -*y*+1, -*z*+1; (iii) -*x*+1, -*y*+1, -*z*+1; (iv) -*x*+1, -*y*, -*z*+1.

4-Phenylpiperazin-1-ium 4-methylbenzoate monohydrate (III)

Crystal data

 $C_{10}H_{15}N_2^+ \cdot C_8H_7O_2^- \cdot H_2O$ $M_r = 316.39$ Triclinic, $P\bar{1}$

Hall symbol: -P 1

 $a = 6.1175$ (5) Å $b = 7.6225$ (7) Å $c = 18.452$ (1) Å $\alpha = 97.421$ (9)° $\beta = 90.403$ (8)° $\gamma = 92.405$ (8)° $V = 852.40$ (12) Å³ $Z = 2$ $F(000) = 340$ $D_x = 1.233$ Mg m⁻³Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2877 reflections

 $\theta = 3.0$ – 27.8 ° $\mu = 0.08$ mm⁻¹ $T = 293$ K

Plate, colourless

 $0.5 \times 0.4 \times 0.08$ mm

Data collection

Oxford Diffraction Xcalibur
diffractometer ω scans

Absorption correction: multi-scan

(CrysAlis RED; Oxford Diffraction, 2009)

 $T_{\min} = 0.837$, $T_{\max} = 1.000$

5354 measured reflections

3126 independent reflections

2248 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.013$ $\theta_{\max} = 25.4$ °, $\theta_{\min} = 3.1$ ° $h = -7 \rightarrow 7$ $k = -7 \rightarrow 9$ $l = -22 \rightarrow 22$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.046$ $wR(F^2) = 0.128$ $S = 1.03$

3118 reflections

226 parameters

4 restraints

0 constraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: structure-
invariant direct methods

Hydrogen site location: mixed

H atoms treated by a mixture of independent
and constrained refinement $w = 1/[\sigma^2(F_o^2) + (0.0582P)^2 + 0.2086P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.16$ e Å⁻³ $\Delta\rho_{\min} = -0.16$ e Å⁻³

Extinction correction: SHELXL2018/3

(Sheldrick 2015b),

 $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.011 (3)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.5691 (4)	0.8192 (4)	0.17484 (12)	0.0759 (7)
H1	0.680156	0.871649	0.206032	0.091*
C2	0.5967 (5)	0.8079 (4)	0.10029 (13)	0.0979 (9)
H2	0.725499	0.854112	0.082348	0.117*
C3	0.4416 (5)	0.7314 (4)	0.05273 (13)	0.0932 (8)
H3	0.462879	0.722711	0.002561	0.112*

C4	0.2536 (5)	0.6676 (4)	0.08016 (13)	0.0966 (9)
H4	0.144014	0.6157	0.048248	0.116*
C5	0.2219 (4)	0.6782 (3)	0.15497 (11)	0.0776 (7)
H5	0.091389	0.633226	0.172178	0.093*
C6	0.3793 (3)	0.7539 (2)	0.20421 (9)	0.0472 (4)
C7	0.5393 (3)	0.8071 (3)	0.32686 (9)	0.0497 (4)
H7A	0.626347	0.904686	0.31147	0.06*
H7B	0.626834	0.703106	0.320724	0.06*
C8	0.4853 (3)	0.8506 (3)	0.40599 (10)	0.0534 (5)
H8A	0.619196	0.865746	0.435007	0.064*
H8B	0.410139	0.960943	0.413335	0.064*
C9	0.1390 (3)	0.6889 (3)	0.38632 (10)	0.0554 (5)
H9A	0.062182	0.798326	0.394853	0.067*
H9B	0.04531	0.594989	0.401599	0.067*
C10	0.1858 (3)	0.6476 (3)	0.30616 (9)	0.0502 (5)
H10A	0.241603	0.529612	0.296889	0.06*
H10B	0.050244	0.647749	0.278608	0.06*
C11	0.3129 (3)	0.7088 (2)	0.62385 (9)	0.0445 (4)
C12	0.2060 (3)	0.7367 (2)	0.69737 (9)	0.0413 (4)
C13	0.2989 (3)	0.6749 (3)	0.75734 (10)	0.0535 (5)
H13	0.428845	0.615675	0.752062	0.064*
C14	0.2008 (4)	0.7003 (3)	0.82503 (11)	0.0656 (6)
H14	0.265661	0.657329	0.864556	0.079*
C15	0.0081 (4)	0.7885 (3)	0.83492 (11)	0.0611 (5)
C16	-0.0837 (3)	0.8490 (3)	0.77515 (11)	0.0583 (5)
H16	-0.214079	0.907628	0.780509	0.07*
C17	0.0129 (3)	0.8250 (2)	0.70719 (10)	0.0484 (4)
H17	-0.052163	0.868519	0.667839	0.058*
C18	-0.0996 (5)	0.8177 (4)	0.90877 (13)	0.0968 (9)
H18A	-0.171921	0.927853	0.913714	0.145*
H18B	0.009588	0.821099	0.946613	0.145*
H18C	-0.204829	0.722525	0.912877	0.145*
N1	0.3441 (2)	0.77371 (19)	0.28025 (7)	0.0434 (4)
N2	0.3444 (2)	0.7069 (2)	0.43010 (8)	0.0487 (4)
O1	0.4844 (2)	0.62506 (19)	0.61816 (7)	0.0632 (4)
O2	0.2244 (2)	0.77468 (19)	0.57233 (7)	0.0601 (4)
OW1	0.7937 (3)	0.7241 (4)	0.52207 (11)	0.1232 (9)
HW1	0.926 (4)	0.748 (5)	0.542 (2)	0.177 (17)*
HW2	0.702 (4)	0.704 (4)	0.5539 (13)	0.119 (11)*
HN1	0.415 (3)	0.605 (2)	0.4225 (11)	0.061 (6)*
HN2	0.310 (3)	0.731 (3)	0.4786 (9)	0.065 (6)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0620 (13)	0.114 (2)	0.0494 (12)	-0.0113 (13)	0.0106 (10)	0.0080 (12)
C2	0.0883 (18)	0.148 (3)	0.0556 (15)	-0.0138 (18)	0.0233 (13)	0.0120 (15)
C3	0.123 (2)	0.109 (2)	0.0456 (13)	-0.0055 (18)	0.0201 (15)	0.0043 (13)

C4	0.130 (2)	0.106 (2)	0.0480 (13)	-0.0340 (18)	-0.0161 (14)	0.0036 (13)
C5	0.0895 (16)	0.0936 (17)	0.0466 (12)	-0.0337 (13)	-0.0042 (11)	0.0105 (11)
C6	0.0561 (11)	0.0456 (10)	0.0397 (9)	0.0045 (8)	0.0038 (8)	0.0042 (7)
C7	0.0473 (10)	0.0550 (11)	0.0456 (10)	-0.0123 (8)	0.0008 (8)	0.0070 (8)
C8	0.0581 (11)	0.0569 (12)	0.0440 (10)	-0.0065 (9)	-0.0031 (8)	0.0045 (8)
C9	0.0439 (10)	0.0734 (13)	0.0494 (11)	-0.0025 (9)	0.0087 (8)	0.0105 (9)
C10	0.0383 (9)	0.0668 (12)	0.0446 (10)	-0.0056 (8)	0.0005 (7)	0.0064 (8)
C11	0.0429 (10)	0.0454 (10)	0.0432 (10)	-0.0062 (8)	0.0042 (7)	0.0012 (7)
C12	0.0413 (9)	0.0376 (9)	0.0439 (9)	-0.0047 (7)	0.0046 (7)	0.0025 (7)
C13	0.0525 (11)	0.0601 (12)	0.0479 (11)	0.0081 (9)	0.0012 (8)	0.0054 (9)
C14	0.0798 (15)	0.0739 (14)	0.0439 (11)	0.0046 (11)	-0.0013 (10)	0.0111 (9)
C15	0.0742 (14)	0.0597 (12)	0.0479 (11)	-0.0009 (10)	0.0167 (10)	0.0012 (9)
C16	0.0577 (12)	0.0554 (12)	0.0614 (12)	0.0098 (9)	0.0173 (9)	0.0025 (9)
C17	0.0525 (10)	0.0458 (10)	0.0476 (10)	0.0047 (8)	0.0056 (8)	0.0072 (8)
C18	0.125 (2)	0.108 (2)	0.0567 (14)	0.0110 (17)	0.0378 (15)	0.0048 (13)
N1	0.0436 (8)	0.0487 (8)	0.0379 (8)	-0.0025 (6)	0.0016 (6)	0.0068 (6)
N2	0.0519 (9)	0.0577 (10)	0.0371 (8)	0.0057 (8)	0.0054 (7)	0.0068 (7)
O1	0.0546 (8)	0.0720 (9)	0.0633 (9)	0.0149 (7)	0.0170 (6)	0.0047 (7)
O2	0.0557 (8)	0.0833 (10)	0.0426 (7)	0.0032 (7)	0.0096 (6)	0.0120 (7)
OW1	0.0607 (12)	0.250 (3)	0.0691 (12)	0.0102 (15)	0.0088 (10)	0.0577 (15)

Geometric parameters (Å, °)

C1—C2	1.379 (3)	C10—H10A	0.97
C1—C6	1.385 (3)	C10—H10B	0.97
C1—H1	0.93	C11—O1	1.249 (2)
C2—C3	1.349 (4)	C11—O2	1.260 (2)
C2—H2	0.93	C11—C12	1.504 (2)
C3—C4	1.356 (4)	C12—C13	1.385 (2)
C3—H3	0.93	C12—C17	1.386 (2)
C4—C5	1.388 (3)	C13—C14	1.384 (3)
C4—H4	0.93	C13—H13	0.93
C5—C6	1.375 (3)	C14—C15	1.383 (3)
C5—H5	0.93	C14—H14	0.93
C6—N1	1.411 (2)	C15—C16	1.374 (3)
C7—N1	1.461 (2)	C15—C18	1.512 (3)
C7—C8	1.497 (2)	C16—C17	1.384 (2)
C7—H7A	0.97	C16—H16	0.93
C7—H7B	0.97	C17—H17	0.93
C8—N2	1.481 (2)	C18—H18A	0.96
C8—H8A	0.97	C18—H18B	0.96
C8—H8B	0.97	C18—H18C	0.96
C9—N2	1.481 (2)	N2—HN1	0.900 (15)
C9—C10	1.504 (2)	N2—HN2	0.918 (15)
C9—H9A	0.97	OW1—HW1	0.886 (19)
C9—H9B	0.97	OW1—HW2	0.839 (18)
C10—N1	1.462 (2)		

C2—C1—C6	121.3 (2)	C9—C10—H10B	109
C2—C1—H1	119.3	H10A—C10—H10B	107.8
C6—C1—H1	119.3	O1—C11—O2	124.44 (16)
C3—C2—C1	121.6 (2)	O1—C11—C12	118.17 (16)
C3—C2—H2	119.2	O2—C11—C12	117.37 (15)
C1—C2—H2	119.2	C13—C12—C17	118.11 (16)
C2—C3—C4	118.1 (2)	C13—C12—C11	120.69 (15)
C2—C3—H3	120.9	C17—C12—C11	121.19 (15)
C4—C3—H3	120.9	C14—C13—C12	120.81 (17)
C3—C4—C5	121.2 (2)	C14—C13—H13	119.6
C3—C4—H4	119.4	C12—C13—H13	119.6
C5—C4—H4	119.4	C15—C14—C13	121.14 (19)
C6—C5—C4	121.4 (2)	C15—C14—H14	119.4
C6—C5—H5	119.3	C13—C14—H14	119.4
C4—C5—H5	119.3	C16—C15—C14	117.78 (17)
C5—C6—C1	116.25 (18)	C16—C15—C18	120.7 (2)
C5—C6—N1	121.84 (18)	C14—C15—C18	121.5 (2)
C1—C6—N1	121.81 (17)	C15—C16—C17	121.68 (18)
N1—C7—C8	112.51 (14)	C15—C16—H16	119.2
N1—C7—H7A	109.1	C17—C16—H16	119.2
C8—C7—H7A	109.1	C16—C17—C12	120.47 (17)
N1—C7—H7B	109.1	C16—C17—H17	119.8
C8—C7—H7B	109.1	C12—C17—H17	119.8
H7A—C7—H7B	107.8	C15—C18—H18A	109.5
N2—C8—C7	110.19 (15)	C15—C18—H18B	109.5
N2—C8—H8A	109.6	H18A—C18—H18B	109.5
C7—C8—H8A	109.6	C15—C18—H18C	109.5
N2—C8—H8B	109.6	H18A—C18—H18C	109.5
C7—C8—H8B	109.6	H18B—C18—H18C	109.5
H8A—C8—H8B	108.1	C6—N1—C7	116.19 (14)
N2—C9—C10	110.84 (14)	C6—N1—C10	116.09 (14)
N2—C9—H9A	109.5	C7—N1—C10	113.11 (13)
C10—C9—H9A	109.5	C8—N2—C9	108.70 (15)
N2—C9—H9B	109.5	C8—N2—HN1	108.9 (13)
C10—C9—H9B	109.5	C9—N2—HN1	109.5 (13)
H9A—C9—H9B	108.1	C8—N2—HN2	111.2 (13)
N1—C10—C9	112.96 (15)	C9—N2—HN2	108.7 (13)
N1—C10—H10A	109	HN1—N2—HN2	109.8 (18)
C9—C10—H10A	109	HW1—OW1—HW2	111 (3)
N1—C10—H10B	109		
C6—C1—C2—C3	-0.7 (5)	C13—C14—C15—C16	0.4 (3)
C1—C2—C3—C4	1.1 (5)	C13—C14—C15—C18	-179.7 (2)
C2—C3—C4—C5	-0.8 (5)	C14—C15—C16—C17	-0.6 (3)
C3—C4—C5—C6	0.1 (4)	C18—C15—C16—C17	179.5 (2)
C4—C5—C6—C1	0.4 (4)	C15—C16—C17—C12	0.7 (3)
C4—C5—C6—N1	176.6 (2)	C13—C12—C17—C16	-0.5 (3)
C2—C1—C6—C5	-0.1 (4)	C11—C12—C17—C16	179.76 (17)

C2—C1—C6—N1	-176.3 (2)	C5—C6—N1—C7	162.90 (19)
N1—C7—C8—N2	56.4 (2)	C1—C6—N1—C7	-21.0 (3)
N2—C9—C10—N1	-53.3 (2)	C5—C6—N1—C10	26.2 (3)
O1—C11—C12—C13	1.6 (2)	C1—C6—N1—C10	-157.72 (19)
O2—C11—C12—C13	-177.16 (17)	C8—C7—N1—C6	172.21 (15)
O1—C11—C12—C17	-178.61 (16)	C8—C7—N1—C10	-49.8 (2)
O2—C11—C12—C17	2.6 (2)	C9—C10—N1—C6	-173.76 (14)
C17—C12—C13—C14	0.3 (3)	C9—C10—N1—C7	48.2 (2)
C11—C12—C13—C14	-179.94 (18)	C7—C8—N2—C9	-60.59 (19)
C12—C13—C14—C15	-0.3 (3)	C10—C9—N2—C8	59.1 (2)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$OW1-HW1\cdots O2^i$	0.89 (3)	1.90 (3)	2.782 (2)	171 (4)
$OW1-HW2\cdots O1$	0.84 (2)	1.92 (3)	2.751 (2)	172 (3)
$N2-HN1\cdots O1^{ii}$	0.90 (2)	1.94 (2)	2.819 (2)	164 (2)
$N2-HN2\cdots O2$	0.92 (2)	1.80 (2)	2.7207 (19)	176 (2)
$C8-H8A\cdots OW1$	0.97	2.33	3.116 (3)	138

Symmetry codes: (i) $x+1, y, z$; (ii) $-x+1, -y+1, -z+1$.

4-Phenylpiperazin-1-ium trifluoroacetate 0.123-hydrate (IV)

Crystal data

 $C_{10}H_{15}N_2^+ \cdot C_2F_3O_2^- \cdot 0.123H_2O$ $M_r = 278.47$ Triclinic, $P\bar{1}$

Hall symbol: -P 1

 $a = 9.6544$ (6) \AA $b = 9.9029$ (6) \AA $c = 15.2090$ (9) \AA $\alpha = 79.621$ (6) $^\circ$ $\beta = 86.579$ (6) $^\circ$ $\gamma = 70.603$ (6) $^\circ$ $V = 1349.10$ (15) \AA^3 $Z = 4$ $F(000) = 580.9$ $D_x = 1.371$ Mg m^{-3} Mo $K\alpha$ radiation, $\lambda = 0.71073$ \AA

Cell parameters from 3827 reflections

 $\theta = 2.6\text{--}27.7^\circ$ $\mu = 0.12$ mm^{-1} $T = 293$ K

Prism, colourless

 $0.48 \times 0.48 \times 0.36$ mm

Data collection

Oxford Diffraction Xcalibur
diffractometer ω scans

Absorption correction: multi-scan

(CrysAlis RED; Oxford Diffraction, 2009)

 $T_{\min} = 0.724$, $T_{\max} = 1.000$

9220 measured reflections

4940 independent reflections

2777 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.014$ $\theta_{\max} = 25.3^\circ$, $\theta_{\min} = 2.6^\circ$ $h = -11 \rightarrow 11$ $k = -11 \rightarrow 11$ $l = -18 \rightarrow 18$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.070$ $wR(F^2) = 0.235$ $S = 1.07$

4927 reflections

375 parameters

4 restraints

0 constraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: structure-invariant direct methods
 Hydrogen site location: mixed
 H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.1128P)^2 + 0.3819P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.42 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.28 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C1	0.2559 (3)	0.7294 (3)	0.51366 (18)	0.0565 (7)	
C2	0.1392 (4)	0.7473 (3)	0.5737 (2)	0.0733 (9)	
H2	0.043765	0.775704	0.552306	0.088*	
C3	0.1634 (5)	0.7235 (4)	0.6648 (3)	0.0913 (12)	
H3	0.083976	0.737161	0.703837	0.11*	
C4	0.3019 (6)	0.6803 (4)	0.6979 (2)	0.0945 (12)	
H4	0.317436	0.664341	0.759246	0.113*	
C5	0.4172 (5)	0.6606 (4)	0.6405 (3)	0.0939 (11)	
H5	0.512209	0.630205	0.66299	0.113*	
C6	0.3958 (4)	0.6850 (4)	0.5497 (2)	0.0746 (9)	
H6	0.476639	0.671553	0.511724	0.089*	
C7	0.3582 (4)	0.7522 (4)	0.3646 (2)	0.0748 (9)	
H7A	0.439678	0.665346	0.385364	0.09*	
H7B	0.386552	0.835603	0.369838	0.09*	
C8	0.3289 (5)	0.7565 (4)	0.2686 (2)	0.0924 (11)	
H8A	0.41407	0.762586	0.232966	0.111*	
H8B	0.311597	0.667892	0.261853	0.111*	
C9	0.0709 (5)	0.8815 (5)	0.2926 (3)	0.1078 (14)	
H9A	0.044836	0.796648	0.287542	0.129*	
H9B	-0.01211	0.967225	0.272441	0.129*	
C10	0.1038 (4)	0.8774 (5)	0.3876 (2)	0.0933 (12)	
H10A	0.121556	0.966325	0.393067	0.112*	
H10B	0.019188	0.872467	0.424128	0.112*	
C11	0.7162 (3)	0.7837 (3)	0.34093 (18)	0.0568 (7)	
C12	0.6756 (3)	0.9091 (3)	0.3787 (2)	0.0682 (8)	
H12	0.637995	0.999179	0.342289	0.082*	
C13	0.6902 (4)	0.9023 (4)	0.4692 (2)	0.0777 (9)	
H13	0.660653	0.987687	0.492938	0.093*	
C14	0.7475 (4)	0.7719 (5)	0.5247 (2)	0.0786 (10)	
H14	0.758672	0.767891	0.585529	0.094*	
C15	0.7878 (4)	0.6476 (4)	0.4884 (2)	0.0856 (10)	
H15	0.826468	0.558096	0.525259	0.103*	
C16	0.7723 (4)	0.6527 (4)	0.3984 (2)	0.0760 (9)	
H16	0.799969	0.566471	0.375625	0.091*	

C17	0.8021 (5)	0.6701 (4)	0.2099 (2)	0.0988 (13)	
H17A	0.807268	0.577457	0.246337	0.119*	
H17B	0.898326	0.680875	0.21027	0.119*	
C18	0.7624 (6)	0.6706 (5)	0.1152 (3)	0.1280 (18)	
H18A	0.837937	0.594874	0.090434	0.154*	
H18B	0.670336	0.65126	0.114843	0.154*	
C19	0.6330 (5)	0.9270 (5)	0.0978 (2)	0.0940 (11)	
H19A	0.540215	0.909315	0.096567	0.113*	
H19B	0.622668	1.021169	0.061705	0.113*	
C20	0.6691 (4)	0.9279 (4)	0.1919 (2)	0.0773 (9)	
H20A	0.756179	0.95605	0.191813	0.093*	
H20B	0.58882	0.999996	0.216184	0.093*	
C21	0.2670 (4)	0.2041 (4)	0.1833 (2)	0.0776 (9)	
C22	0.2420 (8)	0.3636 (6)	0.1772 (3)	0.1240 (19)	0.736 (3)
C22'	0.2420 (8)	0.3636 (6)	0.1772 (3)	0.1240 (19)	0.264 (3)
C23	0.1345 (7)	0.8145 (4)	0.0390 (2)	0.0940 (13)	
C24	0.1356 (7)	0.7597 (5)	-0.0474 (3)	0.1064 (14)	0.736 (3)
C24'	0.1356 (7)	0.7597 (5)	-0.0474 (3)	0.1064 (14)	0.264 (3)
N1	0.2321 (3)	0.7528 (2)	0.42089 (15)	0.0596 (6)	
N2	0.1985 (4)	0.8842 (3)	0.23622 (18)	0.0888 (9)	
N3	0.6944 (3)	0.7878 (3)	0.24907 (15)	0.0687 (7)	
N4	0.7485 (4)	0.8143 (4)	0.05995 (19)	0.1087 (12)	
O1	0.2367 (4)	0.1448 (3)	0.25394 (17)	0.1168 (10)	
O2	0.3178 (4)	0.1492 (4)	0.11854 (18)	0.1323 (12)	
O3	0.2462 (4)	0.8383 (4)	0.05509 (19)	0.1252 (11)	
O4	0.0222 (4)	0.8316 (3)	0.08423 (17)	0.1113 (10)	
F1	0.3669 (7)	0.3949 (7)	0.1721 (5)	0.202 (3)	0.736 (3)
F2	0.1775 (9)	0.4242 (5)	0.2477 (4)	0.184 (3)	0.736 (3)
F3	0.1706 (8)	0.4421 (5)	0.1039 (4)	0.179 (3)	0.736 (3)
F1'	0.302 (2)	0.407 (2)	0.0998 (15)	0.202 (3)	0.264 (3)
F2'	0.282 (3)	0.3950 (17)	0.2311 (15)	0.184 (3)	0.264 (3)
F3'	0.102 (2)	0.4303 (16)	0.1642 (11)	0.179 (3)	0.264 (3)
F4	0.2609 (7)	0.6549 (8)	-0.0615 (4)	0.175 (2)	0.736 (3)
F5	0.1187 (10)	0.8524 (4)	-0.1159 (2)	0.181 (3)	0.736 (3)
F6	0.0480 (7)	0.6817 (8)	-0.0460 (4)	0.153 (2)	0.736 (3)
F4'	0.238 (2)	0.789 (2)	-0.1015 (13)	0.175 (2)	0.264 (3)
F5'	0.157 (4)	0.6370 (19)	-0.0412 (9)	0.181 (3)	0.264 (3)
F6'	-0.0072 (19)	0.8256 (19)	-0.0913 (10)	0.153 (2)	0.264 (3)
OW1	0.5122 (12)	0.6320 (14)	0.0361 (10)	0.150 (8)	0.245 (10)
H21	0.207318	0.969376	0.238227	0.18*	
H22	0.187596	0.886717	0.179696	0.18*	
H41	0.826598	0.838953	0.053835	0.18*	
H42	0.722096	0.810784	0.005806	0.18*	
HW1A	0.572085	0.618747	-0.007306	0.225*	0.245 (10)
HW1B	0.441675	0.707638	0.015874	0.225*	0.245 (10)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.065 (2)	0.0490 (15)	0.0531 (16)	-0.0184 (13)	0.0022 (14)	-0.0040 (12)
C2	0.075 (2)	0.069 (2)	0.068 (2)	-0.0195 (16)	0.0111 (17)	-0.0051 (15)
C3	0.116 (3)	0.074 (2)	0.075 (2)	-0.025 (2)	0.033 (2)	-0.0120 (18)
C4	0.141 (4)	0.079 (2)	0.056 (2)	-0.028 (2)	-0.001 (2)	-0.0089 (17)
C5	0.107 (3)	0.098 (3)	0.072 (2)	-0.025 (2)	-0.026 (2)	-0.0087 (19)
C6	0.064 (2)	0.089 (2)	0.066 (2)	-0.0204 (17)	-0.0069 (16)	-0.0074 (16)
C7	0.073 (2)	0.086 (2)	0.0618 (18)	-0.0228 (17)	0.0105 (16)	-0.0132 (16)
C8	0.122 (3)	0.098 (3)	0.062 (2)	-0.039 (2)	0.010 (2)	-0.0215 (19)
C9	0.099 (3)	0.139 (4)	0.079 (3)	-0.044 (3)	-0.030 (2)	0.017 (2)
C10	0.065 (2)	0.128 (3)	0.063 (2)	-0.010 (2)	-0.0084 (17)	0.0067 (19)
C11	0.0597 (18)	0.0649 (18)	0.0512 (15)	-0.0254 (14)	0.0087 (13)	-0.0168 (13)
C12	0.086 (2)	0.0636 (18)	0.0605 (17)	-0.0280 (16)	0.0031 (15)	-0.0183 (14)
C13	0.089 (2)	0.093 (2)	0.066 (2)	-0.0390 (19)	0.0122 (18)	-0.0374 (19)
C14	0.076 (2)	0.119 (3)	0.0513 (17)	-0.044 (2)	0.0031 (16)	-0.019 (2)
C15	0.092 (3)	0.093 (3)	0.063 (2)	-0.024 (2)	-0.0075 (18)	-0.0006 (18)
C16	0.091 (2)	0.065 (2)	0.066 (2)	-0.0172 (16)	0.0037 (17)	-0.0140 (15)
C17	0.167 (4)	0.071 (2)	0.064 (2)	-0.042 (2)	0.028 (2)	-0.0283 (17)
C18	0.221 (6)	0.115 (4)	0.074 (3)	-0.078 (4)	0.039 (3)	-0.050 (2)
C19	0.118 (3)	0.118 (3)	0.0538 (19)	-0.051 (2)	0.0026 (19)	-0.0131 (19)
C20	0.094 (3)	0.076 (2)	0.0602 (18)	-0.0257 (18)	-0.0026 (17)	-0.0121 (16)
C21	0.102 (3)	0.086 (2)	0.0458 (17)	-0.0309 (19)	-0.0063 (17)	-0.0135 (16)
C22	0.194 (6)	0.122 (4)	0.068 (3)	-0.074 (4)	-0.044 (3)	0.011 (3)
C22'	0.194 (6)	0.122 (4)	0.068 (3)	-0.074 (4)	-0.044 (3)	0.011 (3)
C23	0.159 (4)	0.066 (2)	0.052 (2)	-0.032 (2)	0.011 (3)	-0.0113 (16)
C24	0.172 (5)	0.084 (3)	0.070 (2)	-0.047 (3)	0.028 (3)	-0.028 (2)
C24'	0.172 (5)	0.084 (3)	0.070 (2)	-0.047 (3)	0.028 (3)	-0.028 (2)
N1	0.0591 (15)	0.0674 (15)	0.0519 (13)	-0.0234 (12)	-0.0030 (11)	-0.0028 (11)
N2	0.136 (3)	0.082 (2)	0.0530 (15)	-0.0447 (19)	-0.0139 (17)	-0.0015 (13)
N3	0.094 (2)	0.0709 (16)	0.0501 (13)	-0.0357 (14)	0.0087 (12)	-0.0184 (12)
N4	0.162 (3)	0.127 (3)	0.0538 (16)	-0.066 (2)	0.0207 (19)	-0.0303 (18)
O1	0.205 (3)	0.0908 (18)	0.0635 (16)	-0.0618 (19)	0.0128 (17)	-0.0133 (13)
O2	0.163 (3)	0.164 (3)	0.0675 (16)	-0.038 (2)	0.0098 (17)	-0.0474 (18)
O3	0.161 (3)	0.157 (3)	0.0719 (18)	-0.062 (3)	0.0133 (19)	-0.0395 (18)
O4	0.154 (3)	0.121 (2)	0.0633 (15)	-0.0477 (19)	0.0290 (17)	-0.0313 (14)
F1	0.249 (6)	0.248 (6)	0.198 (6)	-0.196 (5)	-0.015 (4)	-0.040 (4)
F2	0.289 (8)	0.078 (3)	0.159 (4)	-0.016 (4)	0.037 (5)	-0.053 (2)
F3	0.278 (7)	0.117 (3)	0.142 (4)	-0.095 (3)	-0.104 (5)	0.068 (3)
F1'	0.249 (6)	0.248 (6)	0.198 (6)	-0.196 (5)	-0.015 (4)	-0.040 (4)
F2'	0.289 (8)	0.078 (3)	0.159 (4)	-0.016 (4)	0.037 (5)	-0.053 (2)
F3'	0.278 (7)	0.117 (3)	0.142 (4)	-0.095 (3)	-0.104 (5)	0.068 (3)
F4	0.227 (5)	0.149 (4)	0.137 (4)	-0.024 (4)	0.046 (3)	-0.082 (4)
F5	0.411 (11)	0.084 (2)	0.0504 (17)	-0.087 (4)	-0.014 (3)	0.0007 (16)
F6	0.226 (5)	0.152 (4)	0.118 (3)	-0.092 (4)	0.009 (3)	-0.061 (3)
F4'	0.227 (5)	0.149 (4)	0.137 (4)	-0.024 (4)	0.046 (3)	-0.082 (4)
F5'	0.411 (11)	0.084 (2)	0.0504 (17)	-0.087 (4)	-0.014 (3)	0.0007 (16)

F6'	0.226 (5)	0.152 (4)	0.118 (3)	-0.092 (4)	0.009 (3)	-0.061 (3)
OW1	0.082 (10)	0.146 (13)	0.219 (18)	-0.019 (8)	-0.031 (9)	-0.048 (11)

Geometric parameters (Å, °)

C1—C6	1.388 (4)	C17—C18	1.512 (5)
C1—C2	1.392 (4)	C17—H17A	0.97
C1—N1	1.408 (3)	C17—H17B	0.97
C2—C3	1.384 (5)	C18—N4	1.485 (6)
C2—H2	0.93	C18—H18A	0.97
C3—C4	1.359 (6)	C18—H18B	0.97
C3—H3	0.93	C19—N4	1.465 (5)
C4—C5	1.357 (5)	C19—C20	1.495 (4)
C4—H4	0.93	C19—H19A	0.97
C5—C6	1.375 (5)	C19—H19B	0.97
C5—H5	0.93	C20—N3	1.451 (4)
C6—H6	0.93	C20—H20A	0.97
C7—N1	1.445 (4)	C20—H20B	0.97
C7—C8	1.494 (5)	C21—O1	1.197 (4)
C7—H7A	0.97	C21—O2	1.211 (4)
C7—H7B	0.97	C21—C22'	1.502 (7)
C8—N2	1.489 (5)	C21—C22	1.502 (7)
C8—H8A	0.97	C22—F3	1.321 (6)
C8—H8B	0.97	C22—F1	1.337 (7)
C9—N2	1.464 (5)	C22—F2	1.343 (7)
C9—C10	1.488 (5)	C22'—F2'	1.06 (2)
C9—H9A	0.97	C22'—F3'	1.306 (19)
C9—H9B	0.97	C22'—F1'	1.346 (19)
C10—N1	1.466 (4)	C23—O3	1.224 (5)
C10—H10A	0.97	C23—O4	1.228 (5)
C10—H10B	0.97	C23—C24'	1.506 (6)
C11—C16	1.386 (4)	C23—C24	1.506 (6)
C11—C12	1.390 (4)	C24—F5	1.238 (5)
C11—N3	1.417 (3)	C24—F6	1.319 (7)
C12—C13	1.379 (4)	C24—F4	1.342 (7)
C12—H12	0.93	C24'—F5'	1.151 (17)
C13—C14	1.369 (5)	C24'—F4'	1.32 (2)
C13—H13	0.93	C24'—F6'	1.459 (17)
C14—C15	1.368 (5)	N2—H21	0.8818
C14—H14	0.93	N2—H22	0.8672
C15—C16	1.377 (5)	N4—H41	0.8621
C15—H15	0.93	N4—H42	0.8861
C16—H16	0.93	OW1—HW1A	0.8501
C17—N3	1.471 (4)	OW1—HW1B	0.8501
C6—C1—C2	116.9 (3)	N4—C18—H18B	109.8
C6—C1—N1	122.0 (3)	C17—C18—H18B	109.8
C2—C1—N1	121.1 (3)	H18A—C18—H18B	108.2

C3—C2—C1	120.8 (3)	N4—C19—C20	110.7 (3)
C3—C2—H2	119.6	N4—C19—H19A	109.5
C1—C2—H2	119.6	C20—C19—H19A	109.5
C4—C3—C2	120.8 (3)	N4—C19—H19B	109.5
C4—C3—H3	119.6	C20—C19—H19B	109.5
C2—C3—H3	119.6	H19A—C19—H19B	108.1
C5—C4—C3	119.2 (4)	N3—C20—C19	112.8 (3)
C5—C4—H4	120.4	N3—C20—H20A	109
C3—C4—H4	120.4	C19—C20—H20A	109
C4—C5—C6	121.0 (4)	N3—C20—H20B	109
C4—C5—H5	119.5	C19—C20—H20B	109
C6—C5—H5	119.5	H20A—C20—H20B	107.8
C5—C6—C1	121.2 (3)	O1—C21—O2	127.2 (4)
C5—C6—H6	119.4	O1—C21—C22'	115.1 (4)
C1—C6—H6	119.4	O2—C21—C22'	117.7 (4)
N1—C7—C8	112.5 (3)	O1—C21—C22	115.1 (4)
N1—C7—H7A	109.1	O2—C21—C22	117.7 (4)
C8—C7—H7A	109.1	F3—C22—F1	103.5 (5)
N1—C7—H7B	109.1	F3—C22—F2	108.8 (6)
C8—C7—H7B	109.1	F1—C22—F2	100.9 (6)
H7A—C7—H7B	107.8	F3—C22—C21	112.7 (4)
N2—C8—C7	110.4 (3)	F1—C22—C21	112.9 (6)
N2—C8—H8A	109.6	F2—C22—C21	116.7 (4)
C7—C8—H8A	109.6	F2'—C22'—F3'	112.1 (15)
N2—C8—H8B	109.6	F2'—C22'—F1'	110.0 (16)
C7—C8—H8B	109.6	F3'—C22'—F1'	103.4 (13)
H8A—C8—H8B	108.1	F2'—C22'—C21	116.5 (11)
N2—C9—C10	110.4 (3)	F3'—C22'—C21	106.9 (7)
N2—C9—H9A	109.6	F1'—C22'—C21	107.0 (10)
C10—C9—H9A	109.6	O3—C23—O4	127.7 (4)
N2—C9—H9B	109.6	O3—C23—C24'	115.5 (5)
C10—C9—H9B	109.6	O4—C23—C24'	116.8 (5)
H9A—C9—H9B	108.1	O3—C23—C24	115.5 (5)
N1—C10—C9	112.1 (3)	O4—C23—C24	116.8 (5)
N1—C10—H10A	109.2	F5—C24—F6	111.9 (6)
C9—C10—H10A	109.2	F5—C24—F4	104.5 (6)
N1—C10—H10B	109.2	F6—C24—F4	96.7 (5)
C9—C10—H10B	109.2	F5—C24—C23	115.4 (4)
H10A—C10—H10B	107.9	F6—C24—C23	113.1 (4)
C16—C11—C12	116.8 (3)	F4—C24—C23	113.6 (5)
C16—C11—N3	121.0 (2)	F5'—C24'—F4'	106.2 (16)
C12—C11—N3	122.0 (3)	F5'—C24'—F6'	103.6 (18)
C13—C12—C11	121.1 (3)	F4'—C24'—F6'	109.6 (11)
C13—C12—H12	119.4	F5'—C24'—C23	116.0 (8)
C11—C12—H12	119.4	F4'—C24'—C23	110.3 (9)
C14—C13—C12	121.1 (3)	F6'—C24'—C23	110.7 (6)
C14—C13—H13	119.4	C1—N1—C7	116.1 (2)
C12—C13—H13	119.4	C1—N1—C10	115.0 (2)

C15—C14—C13	118.3 (3)	C7—N1—C10	110.5 (2)
C15—C14—H14	120.8	C9—N2—C8	110.4 (3)
C13—C14—H14	120.8	C9—N2—H21	104.1
C14—C15—C16	121.2 (3)	C8—N2—H21	115
C14—C15—H15	119.4	C9—N2—H22	115.8
C16—C15—H15	119.4	C8—N2—H22	108.2
C15—C16—C11	121.4 (3)	H21—N2—H22	103.4
C15—C16—H16	119.3	C11—N3—C20	115.9 (2)
C11—C16—H16	119.3	C11—N3—C17	114.7 (3)
N3—C17—C18	111.9 (4)	C20—N3—C17	111.7 (2)
N3—C17—H17A	109.2	C19—N4—C18	109.0 (3)
C18—C17—H17A	109.2	C19—N4—H41	108.2
N3—C17—H17B	109.2	C18—N4—H41	116.5
C18—C17—H17B	109.2	C19—N4—H42	109.5
H17A—C17—H17B	107.9	C18—N4—H42	106.4
N4—C18—C17	109.5 (3)	H41—N4—H42	107.1
N4—C18—H18A	109.8	HW1A—OW1—HW1B	104.5
C17—C18—H18A	109.8		
C6—C1—C2—C3	0.8 (4)	O3—C23—C24—F5	-75.6 (8)
N1—C1—C2—C3	179.7 (3)	O4—C23—C24—F5	103.6 (7)
C1—C2—C3—C4	-0.8 (5)	O3—C23—C24—F6	153.8 (6)
C2—C3—C4—C5	0.1 (6)	O4—C23—C24—F6	-26.9 (7)
C3—C4—C5—C6	0.6 (6)	O3—C23—C24—F4	44.9 (7)
C4—C5—C6—C1	-0.5 (6)	O4—C23—C24—F4	-135.8 (6)
C2—C1—C6—C5	-0.2 (5)	O3—C23—C24'—F5'	102 (2)
N1—C1—C6—C5	-179.0 (3)	O4—C23—C24'—F5'	-79 (2)
N1—C7—C8—N2	55.4 (4)	O3—C23—C24'—F4'	-18.9 (12)
N2—C9—C10—N1	-57.1 (5)	O4—C23—C24'—F4'	160.3 (11)
C16—C11—C12—C13	-0.3 (5)	O3—C23—C24'—F6'	-140.4 (8)
N3—C11—C12—C13	176.4 (3)	O4—C23—C24'—F6'	38.8 (9)
C11—C12—C13—C14	1.1 (5)	C6—C1—N1—C7	-10.1 (4)
C12—C13—C14—C15	-1.1 (5)	C2—C1—N1—C7	171.1 (3)
C13—C14—C15—C16	0.3 (5)	C6—C1—N1—C10	-141.4 (3)
C14—C15—C16—C11	0.5 (6)	C2—C1—N1—C10	39.8 (4)
C12—C11—C16—C15	-0.5 (5)	C8—C7—N1—C1	171.8 (3)
N3—C11—C16—C15	-177.2 (3)	C8—C7—N1—C10	-54.8 (4)
N3—C17—C18—N4	56.6 (5)	C9—C10—N1—C1	-170.6 (3)
N4—C19—C20—N3	-55.9 (4)	C9—C10—N1—C7	55.5 (4)
O1—C21—C22—F3	-132.4 (6)	C10—C9—N2—C8	56.8 (4)
O2—C21—C22—F3	49.2 (8)	C7—C8—N2—C9	-55.8 (4)
O1—C21—C22—F1	110.7 (6)	C16—C11—N3—C20	-165.9 (3)
O2—C21—C22—F1	-67.7 (6)	C12—C11—N3—C20	17.6 (4)
O1—C21—C22—F2	-5.5 (8)	C16—C11—N3—C17	-33.3 (4)
O2—C21—C22—F2	176.1 (6)	C12—C11—N3—C17	150.2 (3)
O1—C21—C22'—F2'	47.1 (18)	C19—C20—N3—C11	-174.6 (3)
O2—C21—C22'—F2'	-131.3 (17)	C19—C20—N3—C17	51.5 (4)
O1—C21—C22'—F3'	-79.2 (9)	C18—C17—N3—C11	173.3 (3)

O2—C21—C22'—F3'	102.4 (10)	C18—C17—N3—C20	-52.1 (4)
O1—C21—C22'—F1'	170.6 (10)	C20—C19—N4—C18	59.6 (4)
O2—C21—C22'—F1'	-7.8 (12)	C17—C18—N4—C19	-60.0 (5)

Hydrogen-bond geometry (Å, °)

Cg2 is the centroid of the C1–C6 phenyl ring.

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N2—H21...O1 ⁱ	0.88	1.91	2.790 (4)	174
N2—H22...O3	0.87	2.04	2.860 (4)	157
N2—H22...O4	0.87	2.47	3.164 (5)	137
N4—H41...O4 ⁱⁱ	0.86	1.95	2.759 (6)	156
N4—H42...O2 ⁱⁱⁱ	0.89	1.90	2.758 (4)	164
C18—H18 <i>A</i> ...F5 ⁱⁱⁱ	0.97	2.53	3.273 (18)	134
C18—H18 <i>B</i> ...Ow1	0.97	2.08	2.929 (15)	145
C19—H19 <i>B</i> ...O3 ^{iv}	0.97	2.59	3.420 (5)	144
C20—H20 <i>A</i> ...F5 ^{iv}	0.97	2.64	3.468 (8)	144
C16—H16...Cg2 ^v	0.93	2.99	3.745 (4)	140

Symmetry codes: (i) $x, y+1, z$; (ii) $x+1, y, z$; (iii) $-x+1, -y+1, -z$; (iv) $-x+1, -y+2, -z$; (v) $-x+1, -y+1, -z+1$.