



Online carbohydrate 3D structure validation with the *Privateer* web app

Jordan S. Dialpuri, Haroldas Bagdonas, Lucy C. Schofield, Phuong Thao Pham, Lou Holland, Paul S. Bond, Filomeno Sánchez Rodríguez, Stuart J. McNicholas and Jon Agirre*

York Structural Biology Laboratory, Department of Chemistry, University of York, York YO10 3BG, United Kingdom.

*Correspondence e-mail: jon.agirre@york.ac.uk

Received 28 November 2023

Accepted 9 January 2024

Edited by M. J. van Raaij, Centro Nacional de Biotecnología – CSIC, Spain

Keywords: *Privateer*; validation; polysaccharides; carbohydrates; N-glycosylation; N-glycans; web apps.


Supporting information: this article has supporting information at journals.iucr.org/f

Owing to the difficulties associated with working with carbohydrates, validating glycan 3D structures prior to deposition into the Protein Data Bank has become a staple of the structure-solution pipeline. The *Privateer* software provides integrative methods for the validation, analysis, refinement and graphical representation of 3D atomic structures of glycans, both as ligands and as protein modifiers. While *Privateer* is free software, it requires users to install any of the structural biology software suites that support it or to build it from source code. Here, the *Privateer* web app is presented, which is always up to date and available to be used online (<https://privateer.york.ac.uk>) without installation. This self-updating tool, which runs locally on the user's machine, will allow structural biologists to simply and quickly analyse carbohydrate ligands and protein glycosylation from a web browser whilst retaining all confidential information on their devices.

1. Introduction

Modelling protein glycosylation is an often-overlooked aspect in the field of structural biology. The underappreciation of oligosaccharide modelling can partially be explained by the challenges encountered during the model-building stage. These challenges are primarily caused by the very nature of the carbohydrates: branched, flexible and prone to register microheterogeneity. This results in a tangible impact on protein structures: the median resolution for glycoproteins (2.4 Å) is lower than that of nonglycosylated proteins (2.0 Å) when X-ray crystallography electron-density maps are considered (van Beusekom *et al.*, 2018). As a result, many glycoprotein models deposited in the wwPDB contain flaws ranging from minor irregularities to gross modelling errors (Agirre, Davies *et al.*, 2015; van Beusekom *et al.*, 2018; Crispin *et al.*, 2007; Lütteke *et al.*, 2004; Bagdonas *et al.*, 2020). These difficulties extend to ligand polysaccharide structures as well, with additional complications due to the non-negligible likelihood of finding both α and β anomeric ring forms (Agirre, 2017), and potentially even linear forms, at the reducing end.

To counteract these issues, carbohydrate-validation software packages have been developed and are currently actively being developed to automate the detection and rectification of modelling errors. Some notable examples are legacy tools such as *pdb-care* (PDB CARbohydrate RESidue check) that are used for glycosidic bond and nomenclature validation; the tool utilizes the *pdb2linucs* algorithm, which generates LINUCS notations for modelled oligosaccharides in the model file, which are later compared with HET records in the PDB file (Lütteke *et al.*, 2004). *CARP* (CARbohydrate Ramachandran Plot) is a tool that can be used to evaluate glycosidic linkage



torsions and compare them against data deposited in GlyTorsionDB or GlycoMapsDB (Lütkeke *et al.*, 2005). These tools, which are no longer available, were originally developed and made available as web servers and, while this made them easy to access and run, it detached them from widely used structure-solution software suites such as *Phenix* (Liebschner *et al.*, 2019) and *CCP4* (Agirre *et al.*, 2023). Being remote services, they also relied on data uploads, prompting confidentiality concerns for many industrial users.

The *Privateer* software (Agirre, Iglesias-Fernández *et al.*, 2015) was originally introduced to detect and address issues affecting the ring conformation of pyranosides (Agirre, Davies *et al.*, 2015) using a combination of metrics such as Cremer–Pople puckering parameters (Cremer & Pople, 1975), the real-space correlation coefficient (Atanasova *et al.*, 2022), average *B* factors and nomenclature checks. It was subsequently extended to implement features of the aforementioned legacy tools in a modern setting (Agirre, Iglesias-Fernández *et al.*, 2015; Agirre, 2017). *Privateer* is now capable of performing the automated validation of carbohydrate-ring conformation, monosaccharide nomenclature, glycosidic linkage stereochemistry and torsional conformation (Dialpuri *et al.*, 2023). Another core aspect of *Privateer* is the capability to compute omit $mF_o - DF_c$ maps that are used to calculate a correlation coefficient between carbohydrate residues and input experimental density. *Privateer* can also be used in the context of structure refinement, as it is able to generate chemical dictionaries with unimodal torsion restraints which aid in keeping pyranosides in their lowest energy conformations during refinement; this feature has been used to aid in the low-resolution refinement of large glycoprotein structures (Gristick *et al.*, 2017) and prompted the creation of updated dictionaries for pyranose sugars (Atanasova *et al.*, 2022). Finally, *Privateer* can be used to qualitatively assess the compositional properties of modelled oligosaccharides in the input model file by querying the glycan composition in the GlyTouCan repository and the GlyConnect database (Bagdonas *et al.*, 2020). The most relevant validation information is visually summarized in vector diagrams that are compliant with the Symbol Nomenclature for Glycans (SNFG), where potential modelling errors are highlighted to the user (Haltiwanger, 2016).

The core functionality of *Privateer* is written in the C++ language, and all of the functionality that allows the user to validate atomic models can be accessed by simply running its executable, or binary (the `privateer` command). One of the downsides of running the binary is that as a user it can be difficult or even impossible to harness many of the analytical capabilities of *Privateer*. To compensate for this, *Privateer* also has a rich set of Python bindings that allow other programs to utilize the validation tools contained within its shared library (`libprivateer.so`). Both of these methods require a user to download and compile the source code, or use a software suite such as *CCP4* (Agirre *et al.*, 2023) or *CCP-EM* (Burnley *et al.*, 2017), which not every glyco biologist will have available.

One of the most accessible methods of distributing code is on the web, where users can freely interact with any func-

tionality without the steep learning curve associated with downloading and compiling code (to access the most up-to-date version of the software) or using a software suite. Here, we present a new way to use *Privateer* through a completely client-side web interface running within the user's web browser. By using *Privateer* online, structural biologists can efficiently validate carbohydrates on any modern operating system or device: desktop and laptop computers, and mobile devices.

2. Materials and methods

The source code of *Privateer* uses C++ as the main language for speed-critical operations and Python for higher level functions and scripting; Python functions and types stem from C++ functions wrapped using PyBind11. The primary repository in which this source is kept is hosted at GitHub (<https://github.com/glycojones/privateer>); due to the marked differences between a typical binary distribution of *Privateer* and the web app presented here, a decision was made to create a separate branch ('webapp') for the web app. Crucially, the *Privateer* web app is automatically compiled, built, packaged and deployed on the web server (<https://privateer.york.ac.uk>) using GitHub Actions, which are triggered the moment new features are ready to be released. Therefore, the *Privateer* web app is always up to date with the latest developments and fixes.

The connection between the *Privateer* source code in C++ and the web interface is made possible using the compilation tool *Emscripten*. This tool is able to compile C++ code into WebAssembly, an assembly language that is compatible with most modern web browsers. Using *Emscripten*, parts of *Privateer* are compiled into WebAssembly and are made accessible using a set of JavaScript bindings in a method that is similar to that of the Python bindings already present in *Privateer*. In addition to this, to allow for more functionality, the dependencies of *Privateer* must also be compiled into WebAssembly libraries and bundled in.

The interface of the website is created using *React* (<https://react.dev/>), which allows a flexible and dynamically loadable site to be built and deployed statically. *React* was chosen to enhance compatibility with the 3D visualization software used in this project, *Moorhen* (<https://moorhen.org>). *Moorhen* is web-based molecular-graphics software based on the *Coot* desktop application (Emsley *et al.*, 2010); *Moorhen* and *Coot* rely on the functionality encapsulated in the *Coot* libraries, and therefore are expected to produce similar results. Like the *Privateer* web app, *Moorhen* is compiled into WebAssembly and runs locally on the user's browser. The *Moorhen* panel displayed on the report page presents only a relevant subset of the *Moorhen* interface, including controls for changing the map level, showing symmetry mates and activating Glyco-blocks (McNicholas & Agirre, 2017), a 3D extension of the Standard Symbol Nomenclature for Glycans, commonly known as SNFG (Neelamegham *et al.*, 2019). A brief overview of the controls can be found by pressing 'h' on the *Moorhen* panel. Finally, while the *Moorhen* panel has all of the functionality of the full web application, users are invited to run a

standalone *Moorhen* session to work on models comfortably. This may be performed easily either by accessing the *Moorhen* website (<https://moorhen.org>) or by running it from *CCP4 Cloud* (Krissinel *et al.*, 2022).

3. Results and discussion

The *Privateer* web app has two main data-entry points: the user either chooses a local file or specifies a PDB code to be fetched from the Protein Data Bank (wwPDB Consortium, 2019), as shown in Fig. 1 (also please refer to our Supplementary Video, which demonstrates the case study presented here). This allows the web app to function as both a validation tool during structure determination and a resource for the analysis of deposited structures. When using the web app as a validation tool, a user can choose a single coordinate file in PDB or mmCIF format for geometric validation and, in addition, an accompanying reflection file for further density-fitness analysis as measured by RSCC, the real-space correlation coefficient (Atanasova *et al.*, 2022). Once the correct files have been selected, *Privateer* is used to evaluate known carbohydrates found in the coordinate file.

The web app displays its results in a table with the different glycans. Each table entry contains information for every carbohydrate found in the provided structure, including the chain ID, GlyTouCan (Fujita *et al.*, 2021) identifier and a diagram in the Standard Symbol Nomenclature For Glycans (Haltiwanger, 2016). This information and the SNFG diagrams allow the rapid identification of a particular oligosaccharide group. Clicking on an entry displays more information about the glycan, as shown in Fig. 2.

4. Data security

Perhaps one of the most common reasons not to use the available web servers for carbohydrate structure validation is the potential lack of data security. Confidential structures are sent to a third-party server to be validated, which is likely to be forbidden by many industrial organizations. With a client-side web app such as that presented here, user data are never sent externally; in fact, once the site content is completely loaded, an internet connection is no longer required. By default, however, the site is loaded dynamically to prevent slow initial page loads.

5. A case study: validating, correcting and extending a structure

We will use the high-resolution (1.2 Å) crystal structure of glucose oxidase (PDB entry 3qvp; Kommoju *et al.*, 2011) as an example. This was originally modelled with a five-sugar *N*-glycan chain linked to Asn89, as well as three single *N*-acetylglucosamine pyranosides linked to Asn161, Asn355 and Asn388. The electron-density map is very clear, as expected for such high resolution, and the global quality indicators show that the overall quality of the model is excellent.

The glycan summary of PDB entry 3qvp generated using the 'Fetch from PDB' input box on the *Privateer* web app shows a table with the detected glycans. From this table view, it is simple to identify any glycans that contain any modelling anomaly by looking for orange highlights around a linkage or sugar icon. In the case of PDB entry 3qvp, the single GlcNAc




Figure 1

Input screen of the *Privateer* web app. The user is asked to either supply a structure or a structure code, which is then verified against the Protein Data Bank (wwPDB Consortium, 2019) and downloaded from it.




Figure 2
Detailed view of the glycan chain linked to Asn89 and the *Moorhen* visualization panel. The terminal mannose (red asterisk) can be seen to be in a 1S_5 high-energy conformation as opposed to the expected 4C_1 conformation. Additionally, it has multiple atoms outside the $2mF_o - F_c$ density, presumably due to a clash with the adjacent water molecule (blue asterisk), which lies at the other end of the electron density for the mannose.

glycans are deemed to be modelled within the expected parameters, whereas the $\alpha 1,2$ -linked mannose sugar and glycosidic linkage require further inspection. Clicking on the first table entry reveals more information about this glycan and its potential issues (Fig. 2). The validation data for this glycan attached to Asn89 via an N atom highlights a single

conformational issue with the $\alpha 1,2$ -linked mannose (red asterisk in Fig. 2). Due to a clash with an adjacent water molecule (blue asterisk in Fig. 2) that lies in the continuous electron density of the mannose, this sugar has been distorted into a 1S_5 conformation as opposed to the expected 4C_1 chair conformation. Most likely as a result of this high-energy ring conformation, the $\alpha 1,2$ link (also highlighted in orange) has uncommon torsion angles.

The *Privateer* web app displays these conformational and torsional issues within the SNFG as pop-up messages when the mouse hovers over a sugar or a linkage; however, it is more commonplace for a structural biologist to want to visualize the glycan using desktop model-visualization software. The inbuilt *Moorhen* visualization panel removes the need to open locally installed visualization software and allows trivial inspection of the glycan chain: users need only to click on the sugars for the 3D graphics window to re-centre on them. Using this visualization, it is clear that the $\alpha 1,2$ -linked mannose sugar is indeed in a high-energy conformation and is additionally not modelled within the density. This monosaccharide is a strong candidate for remodelling and subsequent refinement.

To resolve this conformational anomaly, the mannose and neighbouring modelled water molecules were deleted in a *Coot* (Emsley *et al.*, 2010). An $\alpha 1,2$ link was added to a new 4C_1 mannose, which was then refined with *REFMAC5* (Murshudov *et al.*, 2011). Following refinement, the density of




Figure 3
The refined structure of the rebuilt *N*-glycan. $2mF_o - F_c$ density is contoured at 1σ . The individual monosaccharides have been orientated and coloured according to the updated SNFG diagram shown in Fig. 4 (left panel). The annotations show the rebuilt and extended monosaccharides. This figure was generated with *CCP4mg* (McNicholas *et al.*, 2011).




Figure 4 Right: detailed information about the glycan attached to Asn89 in PDB entry 3qyp (Kommoju *et al.*, 2011) after remodelling and refinement. Two additional α -mannose sugars were added to the chain and the conformational issues of the deposited α 1,2-mannose were corrected. Left: linkage torsion angles panel, which highlights the current torsion angles for this linkage (blue crosses) in relation to the previously reported torsional landscape.

two further α -mannose sugars could be identified. One was attached to the rebuilt α 1,2-linked mannose, and another was attached to the β 1–4 mannose, which were then modelled and refined.

This updated structure (Fig. 3) was finally analysed using the web app, with only the original torsional issue remaining. Further inspection of this issue can be performed using the torsion plots that are also available in the detailed glycan view page (Fig. 4). Upon inspection of the MAN-1,2-MAN torsion tab, the highlighted linkage is very close to the expected clusters and hence is little cause for concern. This assertion is validated by inspection of the linkage in the *Moorhen* visualization panel, which shows a good density fit.

6. Conclusions

In conclusion, the *Privateer* web app is an innovative online tool for carbohydrate 3D structure validation. The web app allows fast, local structure validation without the requirement to send any files to an external service. Harnessing the functionality of *Privateer*, users can validate structural composition and conformation, anomericity and linkage-torsion outliers from a web browser.

7. Availability and reproducibility

A video demonstrating the *Privateer* web app is available as supporting information. All source code is publicly available

on GitHub at <https://github.com/glycojones/privateer>. The original, updated structure and map coefficients in MTZ format are available as supporting information. The *Privateer* web app is available at <https://privateer.york.ac.uk> and will remain automatically updated with respect to the source code on GitHub.

Acknowledgements

We are grateful to the University of York IT Services, and Darren Miller in particular, for accommodating our needs and offering timely and excellent technical support. We would like to thank Manal Alzahrani and Elisha Moran (University of York) for testing the website and reporting issues. Jon Agirre is additionally grateful to Luis Fuentes-Montero (Diamond Light Source, UK) for alerting him of the power of Web-Assembly for web-hosted but locally run services. Finally, we would like to pay tribute to Thomas Lütteke, Martin Frank and the late Willy von der Lieth, pioneers of carbohydrate structure validation, whose contributions inspired or directly informed some of the methods that *Privateer* implements.

Funding information

Jordan Dialpuri is funded by the Biotechnology and Biological Sciences Research Council (BBSRC; grant No. BB/T0072221). Haroldas Bagdonas is funded by The Royal Society (grant No. RGF/R1/181006). Lucy Schofield is funded by STFC/CCP4

PhD studentship agreement 4462290 (University of York)/S2 2024 012 (STFC) awarded to Jon Agirre. Phuong Thao Pham is a self-funded PhD student. Paul S. Bond is funded by the Biotechnology and Biological Sciences Research Council (grant No. BB/S005099/1). Filomeno Sanchez is funded by the STFC/CCP4 Collaboration Agreement, Contract ID 1759647 in relation to the CCP4 Web Molecular Graphics project at the University of York (project ID R24844). Stuart McNicholas is funded by the STFC/CCP4 Collaboration Agreement Number CN8680 in relation to the Development of the *CCP4img* Suite and *CCP4i2* Project at the University of York (project ID R22512). Lou Holland is funded by The Royal Society (URF\R\221006). Jon Agirre is a Royal Society University Research Fellow (awards UF160039 and URF\R\221006).

References

- Agirre, J. (2017). *Acta Cryst.* **D73**, 171–186.
- Agirre, J., Atanasova, M., Bagdonas, H., Ballard, C. B., Baslé, A., Beilstein-Edmands, J., Borges, R. J., Brown, D. G., Burgos-Mármol, J. J., Berrisford, J. M., Bond, P. S., Caballero, I., Catapano, L., Chojnowski, G., Cook, A. G., Cowtan, K. D., Croll, T. I., Debreczeni, J. É., Devenish, N. E., Dodson, E. J., Drevon, T. R., Emsley, P., Evans, G., Evans, P. R., Fando, M., Foadi, J., Fuentes-Montero, L., Garman, E. F., Gerstel, M., Gildea, R. J., Hatti, K., Hekkelman, M. L., Heuser, P., Hoh, S. W., Hough, M. A., Jenkins, H. T., Jiménez, E., Joosten, R. P., Keegan, R. M., Keep, N., Krissinel, E. B., Kolenko, P., Kovalevskiy, O., Lamzin, V. S., Lawson, D. M., Lebedev, A. A., Leslie, A. G. W., Lohkamp, B., Long, F., Malý, M., McCoy, A. J., McNicholas, S. J., Medina, A., Millán, C., Murray, J. W., Murshudov, G. N., Nicholls, R. A., Noble, M. E. M., Oeffner, R., Pannu, N. S., Parkhurst, J. M., Pearce, N., Pereira, J., Perrakis, A., Powell, H. R., Read, R. J., Rigden, D. J., Rochira, W., Sammito, M., Sánchez Rodríguez, F., Sheldrick, G. M., Shelley, K. L., Simkovic, F., Simpkin, A. J., Skubak, P., Sobolev, E., Steiner, R. A., Stevenson, K., Tews, I., Thomas, J. M. H., Thorn, A., Valls, J. T., Uski, V., Usón, I., Vagin, A., Velankar, S., Vollmar, M., Walden, H., Waterman, D., Wilson, K. S., Winn, M. D., Winter, G., Wojdyr, M. & Yamashita, K. (2023). *Acta Cryst.* **D79**, 449–461.
- Agirre, J., Davies, G., Wilson, K. & Cowtan, K. (2015). *Nat. Chem. Biol.* **11**, 303.
- Agirre, J., Iglesias-Fernández, J., Rovira, C., Davies, G. J., Wilson, K. S. & Cowtan, K. D. (2015). *Nat. Struct. Mol. Biol.* **22**, 833–834.
- Atanasova, M., Nicholls, R. A., Joosten, R. P. & Agirre, J. (2022). *Acta Cryst.* **D78**, 455–465.
- Bagdonas, H., Ungar, D. & Agirre, J. (2020). *Beilstein J. Org. Chem.* **16**, 2523–2533.
- Beusekom, B. van, Lütteke, T. & Joosten, R. P. (2018). *Acta Cryst.* **F74**, 463–472.
- Burnley, T., Palmer, C. M. & Winn, M. (2017). *Acta Cryst.* **D73**, 469–477.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Crispin, M., Stuart, D. I. & Jones, E. Y. (2007). *Nat. Struct. Mol. Biol.* **14**, 354.
- Dialpuri, J. S., Bagdonas, H., Atanasova, M., Schofield, L. C., Hekkelman, M. L., Joosten, R. P. & Agirre, J. (2023). *Acta Cryst.* **D79**, 462–472.
- Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. (2010). *Acta Cryst.* **D66**, 486–501.
- Fujita, A., Aoki, N. P., Shinmachi, D., Matsubara, M., Tsuchiya, S., Shiota, M., Ono, T., Yamada, I. & Aoki-Kinoshita, K. F. (2021). *Nucleic Acids Res.* **49**, D1529–D1533.
- Gristick, H. B., Wang, H. & Bjorkman, P. J. (2017). *Acta Cryst.* **D73**, 822–828.
- Haltiwanger, R. S. (2016). *Glycobiology*, **26**, 217.
- Kommoju, P., Chen, Z., Bruckner, R. C., Mathews, F. S. & Jorns, M. S. (2011). *Biochemistry*, **50**, 5521–5534.
- Krissinel, E., Lebedev, A. A., Uski, V., Ballard, C. B., Keegan, R. M., Kovalevskiy, O., Nicholls, R. A., Pannu, N. S., Skubák, P., Berrisford, J., Fando, M., Lohkamp, B., Wojdyr, M., Simpkin, A. J., Thomas, J. M. H., Oliver, C., Vornrhein, C., Chojnowski, G., Basle, A., Purkiss, A., Isupov, M. N., McNicholas, S., Lowe, E., Triviño, J., Cowtan, K., Agirre, J., Rigden, D. J., Uson, I., Lamzin, V., Tews, I., Bricogne, G., Leslie, A. G. W. & Brown, D. G. (2022). *Acta Cryst.* **D78**, 1079–1089.
- Liebschner, D., Afonine, P. V., Baker, M. L., Bunkóczi, G., Chen, V. B., Croll, T. I., Hintze, B., Hung, L.-W., Jain, S., McCoy, A. J., Moriarty, N. W., Oeffner, R. D., Poon, B. K., Prisant, M. G., Read, R. J., Richardson, J. S., Richardson, D. C., Sammito, M. D., Sobolev, O. V., Stockwell, D. H., Terwilliger, T. C., Urzhumtsev, A. G., Videau, L. L., Williams, C. J. & Adams, P. D. (2019). *Acta Cryst.* **D75**, 861–877.
- Lütteke, T., Frank, M. & von der Lieth, C.-W. (2004). *Carbohydr. Res.* **339**, 1015–1020.
- Lütteke, T., Frank, M. & von der Lieth, C.-W. (2005). *Nucleic Acids Res.* **33**, D242–D246.
- McNicholas, S. & Agirre, J. (2017). *Acta Cryst.* **D73**, 187–194.
- McNicholas, S., Potterton, E., Wilson, K. S. & Noble, M. E. M. (2011). *Acta Cryst.* **D67**, 386–394.
- Murshudov, G. N., Skubák, P., Lebedev, A. A., Pannu, N. S., Steiner, R. A., Nicholls, R. A., Winn, M. D., Long, F. & Vagin, A. A. (2011). *Acta Cryst.* **D67**, 355–367.
- Neelamegham, S., Aoki-Kinoshita, K., Bolton, E., Frank, M., Lisacek, F., Lütteke, T., O’Boyle, N., Packer, N. H., Stanley, P., Toukach, P., Varki, A., Woods, R. J., Darvill, A., Dell, A., Henrissat, B., Bertozzi, C., Hart, G., Narimatsu, H., Freeze, H., Yamada, I., Paulson, J., Prestegard, J., Marth, J., Vliegthart, J. F. G., Etzler, M., Aebi, M., Kanehisa, M., Taniguchi, N., Edwards, N., Rudd, P., Seeberger, P., Mazumder, R., Ranzinger, R., Cummings, R., Schnaar, R., Perez, S., Kornfeld, S., Kinoshita, T., York, W. & Knirel, Y. (2019). *Glycobiology*, **29**, 620–624.
- wwPDB Consortium (2019). *Nucleic Acids Res.* **47**, D520–D528.