

02-Methods for Structure Determination and Analysis,
Computing and Graphics

49

stronger chemical constraints. One such constraint is the connectivity of the macromolecule. We have developed a rapid algorithm for measuring the connectivity of a map which shows promise in reducing the multiplicity of solutions to the phase problem. We have also developed a refinement method (PRISM) which exploits the connectivity constraint to iteratively improve phases. An initial electron density map is generated with inaccurate phases derived from a partial structure or from isomorphous replacement. A linear connected skeleton is then constructed from the map using a modified version of Greer's algorithm and a new map is created from the skeleton. This "skeletonized" map is Fourier transformed to obtain new phases, which are combined with any starting phase information and the experimental structure factor amplitudes to produce a new map. The procedure is iterated until convergence is reached. The method has been applied to problems with starting phase information from either molecular replacement or isomorphous replacement and appears to be a significant improvement over solvent flattening in both cases.

DS-02.07.04 DIRECT METHODS AND MACROMOLECULAR CRYSTALLOGRAPHY: LIGHTS AND LIMITS. By C. Giacobazzo*, A. Guagliardi, Dipartimento Geomineralogico, Università di Bari, 70124 Bari, Italy; D. Siliqi, Department of Inorganic Chemistry, University of Tirana, Tirana, Albania.

Several papers can be found in literature which describe the application of Direct Methods to macromolecules. Their efficiency is tested both for *ab initio* phasing and for phase refinement and extension. In this paper the role of direct methods in the field of macromolecular crystallography is analyzed. A criterion is formulated which suggests the necessary conditions for the success or the failure of the *ab initio* direct procedure. Most of the experimental protein data do not satisfy such a criterion, therefore their *ab initio* solution is a quite improbable event.

DS-02.07.05 JOINT X-RAY AND NMR REFINEMENT. By B. Shaanan, Department of Biological Chemistry, The Institute of Life Sciences, The Hebrew University of Jerusalem, Israel

PS-02.07.06 AMORE, AN INTEGRATED MOLECULAR REPLACEMENT PROGRAM IN PROTEIN CRYSTALLOGRAPHY: SOME APPLICATIONS TO MULTIBODY SYSTEMS. By J.Navaza¹, Y.Mauguen¹, P.Sauidjian², T. Prangé², P.Alzari³ and G.A.Bentley³.
¹Laboratoire de Physique, Faculté de Pharmacie, 92290 CHATENAY-MALABRY, France. ²Chimie Biomoléculaire, URA 1430 CNRS, UFR Biomédicale, 93012-BOBIGNY CEDEX, France. ³Unité d'Immunologie Structurale, URA 359 CNRS, Institut Pasteur, 25 rue du Docteur Roux, 75724 PARIS CEDEX, France.

A new strategy for Molecular Replacement calculations in protein crystallography has been implemented in the AMoRe package of programs. The algorithms have now been extensively tested in several crystal structures containing multiple copies of the proteins in the asymmetric unit. The examples discussed in the present communication include:

- The complex between Fab F9.13.7 and Guinea-fowl lysozyme with two molecules in the asymmetric unit, using three different search probes (lysozyme, variable and constant regions of the Fab), five out of the six subunits could be sequentially positioned in a single run of AMoRe.
- The trigonal form of Tumor Necrosis Factor, with six copies in the asymmetric unit (a dimer of trimers), using a trimer as the search model.
- A new orthorhombic form of Erabutoxin-b crystallized in presence of KSCN (two copies in the asymmetric unit).
- The complex of a bacterial ribonuclease, barnase, with its specific proteic inhibitor, bastar. There are three copies of the complex in the asymmetric unit. The barnase structure, representing approximately 1/6th of the a.u., was used as the search model.
- An hexagonal form of bastar with four molecules in the a.u., using as the search model the inhibitor subunit, taken out from the above refined complex.

PS-02.07.07 PHASE PERTURBATION AS A MEANS OF REDUCING MODEL BIAS IN MACROMOLECULAR CRYSTALLOGRAPHY. By M.V.Hosur and K.K.Kannan Solid State Physics Division, Bhabha Atomic Research Centre, Trombay, Bombay-400085, INDIA.

The **Molecular Replacement Method** is being increasingly used to solve protein structures by X-ray crystallography. It has also been recognized that the search model used in the above method, introduces a phase bias that complicates interpretation of the calculated electron density maps. A number of attempts have been made to reduce this model bias either by calculating OMIT maps or by using modified amplitudes in Fourier calculations. However, the features of a Fourier map are determined more by the phase rather than the amplitude of the coefficient. We have therefore explored the possibility of altering the phases of the coefficients as a means of reducing model bias in Fourier calculations. A variety of schemes of phase perturbation have been tried, with interesting results. These results and their implications to solving protein structures by the Molecular Replacement Method will be discussed.

PS-02.07.08 ASSESSMENT OF BULK SOLVENT MODELS BY CROSS-VALIDATION. By A.T. Brünger and J.-S. Jiang*, Department of Molecular Biophysics, and Biochemistry, Yale University, U.S.A.

Bulk solvent models can play an important part in the modeling of macromolecular diffraction data. Bulk solvent models are aimed at reducing the residual for the low-resolution reflections. A low R value is not necessarily an indicator for the quality of the model. For example, we have shown earlier (Brünger, A.T., *Nature* **355**,472-474, 1992) that a bulk solvent model consisting of a disordered liquid of point atoms actually worsens the information content of the crystal structure. Cross-validation showed considerable promise to avoid this type of overfitting.