

03-Crystallography of Biological Macromolecules

101

The double-headed Bowman-Birk type trypsin inhibitor(BBI), I (Mr=14,000, 123 amino acid residues) was isolated from wheat germs. To compare the tertiary structures of BBIs in cereal grains with those in leguminous plants (for example, Suzuki et al, J. Biochem.(Tokyo), 1987, **101**, 267-274), the structural study of I-2, the major component of the group I, have been carried out.

I-2 and its 1:2 complex with bovine β -trypsin have been crystallized. For I-2 two morphologically different crystal forms were obtained. Crystal form (1) is tetragonal, P4₁22 or P4₃22, a=55.45(2), c=129.1(2)Å, Vm=3.55Å³/dalton, D₀=1.30gcm⁻³, Z=8 and solvent content(S)=65%. Crystal form (2) is monoclinic, space group C2. The cell parameters show significant variation even for crystals in a same batch. The median parameters are: a=83.9, b=41.5, c=45.7Å, β =95.9°, Vm=2.83Å³/dalton, D₀=1.33gcm⁻³, Z=4, and S=44%. The crystals of trypsin-I-2 (2:1) complex are orthorhombic, P2₁2₁2₁, a=73.49(2), b=120.56(3), c=70.04(2)Å, Vm=2.58Å³/dalton, D₀=1.28gcm⁻³, Z=4 and S=50%. The X-ray intensity data of trypsin-I-2 complex were collected using a Weissenberg camera (Sakabe, J. Appl. Cryst, 1983, **16**, 542-547) at the BL6A2 station, Photon Factory, National Laboratory for High Energy Physics. A total of 26,565 independent reflections between 25 and 2.3Å resolution were measured, giving the merging R of 0.130.

The structure of the trypsin-I-2 complex was determined by the molecular replacement method using 6~3.5Å data and the structure of bovine trypsin. The structure of I-2 was traced on a 3A difference-Fourier map, and the model building was carried out using TURBO-FRODO (Cambillau, 1992). Currently 18 residues were constructed. The R-factor was 0.324 for 10,674 reflections (Fobs>5 σ (F)) between 6 and 3Å resolution. The partial structure of the two reactive site regions of I-2 is essentially similar to that of the 1:1 complex between trypsin and azdzuki inhibitor AB-1 (Sunogae et al., J. Biochem.(Tokyo), 1986, **100**, 1637-1646).

PS-03.07.09 STRUCTURE OF OLIGO-1,6-GLUCOSIDASE REFINED AT 2.0 Å RESOLUTION. By Y. Hata¹⁾, H. Kizaki²⁾, K. Watanabe²⁾, Y. Suzuki²⁾ and Y. Katsube³⁾, ¹⁾Institute for Chemical Research, Kyoto University, ²⁾Department of Agricultural Chemistry, Kyoto Prefectural University, and ³⁾Institute for Protein Research, Osaka University, Japan.

Bacillus oligo-1,6-glucosidases exhibit a strong correlation between their proline content and thermostability. In order to elucidate the contribution of proline to their thermostability as well as the enzyme function, we have analyzed the structure of oligo-1,6-glucosidase (Mr=66010, 558 residues) from a mesophile *B. cereus* by X-ray method. Crystals of the enzyme grew in hanging drops (Watanabe, K., Kitamura, K., Hata, Y., Katsube, Y., & Suzuki, Y. (1991). *FEBS Lett.* 290, 221-223). The structure was solved by MIR method with three derivatives prepared in HgCl₂, UO₂(NO₃)₂ and Sm(NO₃)₃ solutions. The intensity data were collected using synchrotron radiation (λ =1.04 Å) at BL6A₂ station in KEK-PF, Tsukuba. The data were recorded on Fujii image-plates using a Weissenberg camera. The structure model was built with the FRODO on PS390 graphics by interpreting a 3.0 Å resolution electron density map (m)=0.76). The structure refinement was started at this resolution with the program XPLOR, and the resolution has been extended to 2.0 Å. The R-factor for the current model with 221 waters is 0.196 for 43328 reflections (92%) between 8.0 and 2.0 Å resolution. The structure of *B. cereus* oligo-1,6-glucosidase is similar to those of α -amylases and can be subdivided into three domains. The N-terminal domain has a (β/α)-barrel with four additional α -helices. The sub-domain between the third β -strand and α -helix of the N-domain is a long loop containing a small sheet of three β -strands. The C-terminal domain forms an irregular β -barrel of eight strands. The sequence homology and consequent structural similarity between the oligo-1,6-glucosidases of a thermophile and a mesophile suggest that most of the prolines responsible for increase of the thermostability should be positioned on second sites of β -turns.

PS-03.07.10 CRYSTAL STRUCTURE OF ALKALINE SERINE

PROTEASE OF BACTERIAL ORIGIN. By T. Yamane¹, T. Hatanaka¹, T. Kani¹, T. Naruse¹, A. Suzuki¹, T. Ashida^{1*}, T. Kobayashi² and S. Ito², ¹Department of Biotechnology, School of Engineering, Nagoya University, Nagoya 464-01, Japan, ²Tochigi Research Laboratories, Kao Corporation, Tochigi 321-34, Japan.

A new alkaline serine protease(ALPT) was obtained from a *Bacillus* strain. It consists of 269 amino acids and has a molecular weight of 26,715. The sequence of ALPT is 60% homologous with subtilisin Carlsberg(SBC).

The enzyme inhibited with phenylmethylsulfonylfluoride was used for crystallization. Needle crystals (form 1) were grown in 0.05M phosphate buffer (pH 5.5), using 1.4M ammonium sulfate as precipitant. The space group is P2₁2₁2₁ with a=57.79(2), b=75.82(2), c=54.19(1)Å, Z=4 and Vm=2.15Å³/dalton. The cell parameters at first suggested that form 1 might be isomorphous to the crystal of SBC(Petsko & Tsernoglou, J. Mol. Biol., 1976, **106**, 453-456) if the a and b axes were exchanged. The other crystals (form 2) were obtained from 0.05M acetate buffer (pH 5) and belong to space group P2₁2₁2₁ with a=47.3, b=62.5, c=75.6Å, Z=4 and Vm=2.09Å³/dalton, which is isomorphous to the crystals of high-alkaline serine protease PB92 (van der Laan et al, Protein Engineering, 1992, **5**, 405-411).

The X-ray intensity data of form 1 were collected using a Weissenberg camera (Sakabe, J. Appl. Cryst, 1983, **16**, 542-547) at the BL6A2 station at the Photon Factory, National Laboratory for High Energy Physics. The needle (a*) axis was mounted parallel to the rotation axis of the camera. A total of 8,513 independent reflections up to 1.9Å resolution were measured, giving the merging R of 0.119. The structure was solved by the molecular replacement method using 6~3.5Å data and the structure of SBC (PDB entry number 1SBC). The crystallographic R-factor was 0.472 at this stage. The structure was refined by the X-PLOR system (Bruenger, 1990, X-PLOR Manual, Ver 2.1, Yale Univ., New Haven, USA), with the amino acid sequence of SBC. The current R factor is 0.255 for 6404 reflections between 7 and 2.5Å resolution. The structure of ALPT is essentially the same as that of SBC, though the orientation parameters from the rotation function revealed that it was not isomorphous to that of SBC.

PS-03.07.11 STRUCTURE ANALYSIS OF PHOSPHOLIPASE A2

FROM VENOM OF *TRIMERESURUS FLAVOVIRIDIS*. By E. Matsueda¹, A. Suzuki^{1*}, T. Yamane¹, T. Ashida¹, H. Kihara², and M. Ohno³, ¹Department of Biotechnology, School of Engineering, Nagoya University, Nagoya 464-01, Japan, ²Department of Physiology, Faculty of Medicine, Kagoshima University, Kagoshima 890, Japan, ³Department of Chemistry, Faculty of Science, Kyushu University, Higashi-ku, Fukuoka 812, Japan

Phospholipase A₂ (PLA₂) from the venom of *Trimeresurus flavoviridis* (habu snake) is a homo-dimeric enzyme composed of subunits with 122 residues. Crystals belong to space group P2₁ with a=44.1 Å, b=55.7 Å, c=48.8 Å, and β =92.4°. An asymmetric unit contains one dimer.

X-ray diffraction data up to 3.5Å resolution were collected on a four-circle diffractometer with CuK α radiation. High resolution data up to 2.0Å were recorded on imaging plates by using a Weissenberg camera installed at BL-6A2 station of Photon Factory. These two datasets were merged to give a total of 13822 unique reflections. The merging R was 0.054.

The structure analysis was proceeded with the molecular replacement method using the modified structure of PLA₂ from the Western Diamondback rattlesnake as a search model. The search model was constructed by omitting two turn regions and side-chain atoms of residues different from the habu-snake PLA₂ except C β atoms. Self- and cross-rotation search were performed with the Crowther fast rotation function using 6.0 to 3.5Å resolution data. Self-rotation search shows that two subunits in an asymmetric unit are related by a non-crystallographic 2-fold axis. Using this symmetry, the peaks on the cross-rotation function could be paired with one another. The pair containing the highest peak gave true orientations of two subunits in an asymmetric unit. Molecular translation along the a and c axes was determined for each subunit with the Crowther-Blow translation function using 6.0 to 3.5Å resolution data. Relative position along the b axis was determined by the R-search method using 6.0 to 4.0Å data.