

Poster Sessions

¹Brown University, Molecular Biology, Cell Biology & Biochemistry, Box G-E139, Providence, Rhode Island, 02912, USA, ²Brown University, Providence, Rhode Island, 02912, USA, ³Brown University, Providence, Rhode Island, 02912, USA, ⁴Brown University, Providence, Rhode Island, 02912, USA, ⁵Brown University, Box G-E4, Providence, Rhode Island, 02912, USA, E-mail: David_Critton@brown.edu

Cells of the immune system have inherently high levels of tyrosine phosphorylation. In fact, they express more genes encoding protein tyrosine kinases (PTKs) and protein tyrosine phosphatases (PTPs) than nearly any other cell type. One PTP expressed exclusively in cells of the immune system is hematopoietic protein tyrosine phosphatase (HePTP). HePTP functions to negatively regulate the activation of T cells, and it does so by dephosphorylating and inactivating its only known substrates, the mitogen-activated protein kinases (MAPKs) Erk1/2 and p38. The importance of this regulation is highlighted by the fact that dysregulation of HePTP is associated with several diseases of the immune system, including acute myelogenous leukemia and non-Hodgkins lymphoma. In order to understand the molecular basis of the HePTP:MAPK interaction, we have generated a series of HePTP substrate-trapping mutants (STMs) in order to selectively populate an HePTP:MAPK peptide dephosphorylation complex. STMs have low catalytic activities yet retain high affinities for their substrates. We first determined the biochemical and biophysical characteristics of these HePTP STMs to identify those STMs most suitable for structural studies. We then crystallized and determined the structures of several HePTP STMs in both their apo forms and in complex with two distinct substrate peptide mimetics, e.g. peptides that correspond to the singly- and dually-phosphorylated activation loop of the MAPK Erk2. Finally, we describe how this biochemical, biophysical and structural data has allowed us to identify, for the first time, the structural features of HePTP that confer such a high degree of specificity for its MAPK substrates.

Keywords: hematopoietic protein tyrosine phosphatase (HePTP), extracellular signal-regulated kinase (Erk), T cell activation

P04.12.279

Acta Cryst. (2008). A64, C318

Structure of *Escherichia coli* tyrosine kinase Etk reveals novel activation mechanism

Daniel Cho-En Lee, Jimin Zheng, Zongchao Jia

Queen's University, Biochemistry, Rm 634 Botterell Hall, Kingston, Ontario, K7P1S1, Canada, E-mail: dcl.maltz@gmail.com

While protein tyrosine kinases (PTKs) have been extensively characterized in eukaryotes, far less is known about their emerging counterparts in prokaryotes. The inner-membrane Wzc/Etk protein belongs to the bacterial PTK family, which plays a critical role in regulating the polymerization and transportation of virulence-determining capsular polysaccharide (CPS). The kinase utilizes a unique two-step activation mechanism centering on the intraphosphorylation of a tyrosine residue, although the specific detail remains unknown. Herein we report the first crystal structure of a bacterial PTK, the C-terminal kinase domain of *E. coli* tyrosine kinase (Etk) at 2.5Å resolution. The folding of the Etk kinase domain in bacteria differs markedly from that in eukaryotic PTKs. Based on the structure and supporting mass spectrometric evidence of the PTK observed, a unique activation mechanism is consequently proposed that involves the regulation of the phosphorylation of a single tyrosine residue at position 574 and its specific interaction with a previously unidentified key arginine residue at position 614 (R614) to

unblock the active site.

Keywords: protein tyrosine kinases, pathogenic bacterial mechanism, polysaccharides

P04.12.280


Acta Cryst. (2008). A64, C318

Crystallographic analysis of response regulator protein from *Desulfovibrio vulgaris* Hildenborough

Hirofumi Komori, Yoshiki Higuchi

University of Hyogo, 3-2-1 Koto, Kamigori-cho, Aku-gun, Hyogo, 678-1297, Japan, E-mail: komori@sci.u-hyogo.ac.jp

Sulfate-reducing bacteria *Desulfovibrio vulgaris* can generally use hydrogen and organic acids as electron donors for sulfate reduction. Sulfate reduction and hydrogen oxidation are spatially separated in the cytoplasm and periplasm, respectively. It has been proposed that electron transport linking periplasmic hydrogen oxidation to cytoplasmic sulfate reduction is mediated through the high-molecular-mass cytochrome redox protein complex (the Hmc complex). Two genes (*rrf1* and *rrf2*) encoding response regulator proteins with a putative function in the regulation of gene expression are present immediately downstream from the structural genes of the hmc operon. The deletion of the *rrf1*, 2 genes gives rise to increased hmc operon expression. In order to understand the regulation mechanism of Hmc complex expression, we tried to determine the crystal structure of the response regulator proteins (Rrf1 and Rrf2). We cloned the *rrf1* and *rrf2* genes from *Desulfovibrio vulgaris* Hildenborough and successfully overexpressed, purified and crystallized Rrf1 protein. Here we report the crystal structure of Rrf1 at 2.1 Å resolution.



Keywords: signal transduction, transcription regulation, X-ray crystallography of biological macromolecules

P04.12.281

Acta Cryst. (2008). A64, C318-319

Crystal structure of human ERK1 kinase mono-phosphorylated at Tyr204

Takayoshi Kinoshita¹, Ikuyo Yoshida¹, Kouki Okita², Masaki Gouda², Mamoru Matsubara^{2,3}, Koichi Yokota², Hiroshi Ishiguro², Toshiji Tada¹

¹Osaka Prefecture University, Graduate School of Science, 1-1 Gakuencho, Naka-ku, Sakai, Osaka, 599-8531, Japan, ²Carna Biosciences Inc., Kobe, Hyogo, Japan, ³Faculty of Bioenvironment Science, Kyoto Gakuen University, Kyoto, Japan, E-mail: kinotk@b.s.osakafu-u.ac.jp

ERK is a member of MAP kinase family that regulates cell growth and differentiation in response to extracellular stimulation. ERK consists of two major isoforms, ERK1 and ERK2, which have a high degree of amino acid sequence homology and are different from each other in the intravital behavior. In order to investigate *in vivo* function of ERK1, we determined the crystal structure of human ERK1 complexed with 5-iodotubercidin, a potent inhibitor. Purified ERK1 was identified as auto-phosphorylated protein at Tyr204 by Western blot experiments. Crystals of the complex were obtained using a reservoir solution of 30% PEG4000, 0.2 M lithium sulfate,