P24.01.01

Acta Cryst. (2008). A64, C626

The predicted powder diffraction database (P2D2)

Armel Le Bail

Université du Maine, Laboratoire des Oxydes et Fluorures, CNRS UMR 6010, Avenue Olivier Messiaen, Le Mans, Pays de Loire, 72085, France, E-mail:armel.le_bail@univ-lemans.fr

Massive inorganic crystal structure predictions were recently performed. For millions of virtual zeolites or related materials the composition is imposed : SiO2 or AlPO4 etc. For other predicted inorganic compounds the composition was let to be free, but at least chemical elements were selected and some geometrical rules for organizing them were applied (exclusive corner sharing of polyhedra within the GRINSP software [1]). The explosion of the number of predictions justified the creation of new databases. Among them, the PCOD [2] (Predicted Crystallography Open Database) contains the crystal data of predicted titanosilicates, phosphates, vanadates, niobates, fluoroaluminates (etc). These databases open now the possibility for the identification of a newly synthesized compound by the comparison of its experimental powder pattern with predicted ones. The powder patterns calculated from the >100.000 PCOD entries were gathered into the P2D2 (Predicted Powder Diffraction Database) [3-4], allowing for identification by using any classical search-match software. To be successful, identification attempts require mainly accurate predicted cell parameters. Many improvements, by using empirical or ab initio approaches, will be needed in order to restrict the number of structure candidates to those having really a chance to exist (quite a difficult task). Prediction is obviously a large part of our future in crystallography, the P2D2 represents a small new step in that direction.

[1] A. Le Bail, J. Appl. Cryst. 38 (2005) 389-395.

[2] Predicted Crystallography Open Database - http://www. crystallography.net/pcod/

[3] A. Le Bail, Powder Diffraction (2008) in the press.

[4] Predicted Powder Diffraction Database - http://sdpd.univ-lemans. fr/cod/pcod/P2D2/

Keywords: crystal structure prediction, databases, powder diffraction

P24.01.02

Acta Cryst. (2008). A64, C626

Databases for absorption, XAFS and XANES, and future opportunities for research and investigation

Christopher T. Chantler

University of Melbourne, School of Physics, University of Melbourne, School of Physics, Parkville, Victoria, 3010, Australia, E-mail : chantler@unimelb.edu.au

Databases for absorption include the International Union of Crystallography (IUCr) data, particularly of Creagh, Hubbell & Maslen et al.; the XCOM data of Hubbell & Scofield et al.; and the most recent FFAST (Form Factor ans Scattering Tables) of Chantler; together with experimental data using the X-ray Extended Range Technique (XERT) of Chantler. These databases typically have different purposes and uses, so that guidance and explanation is useful for a user to make informed comparisons with his experiments or medical or laboratory applications. Other databases exist, whether particularly for isolated atoms, diffraction, or condensed matter. The needs of X-ray Absorption Fine Structure (XAFS) and X-ray Absorption Near-Edge Structure (XANES) have often been somewhat different, for example needing fine spacing near Keywords: databases, absorption, XAFS

P24.02.03

Acta Cryst. (2008). A64, C626

Modernization of protein crystallography data formats: PDBML as a replacement for PDB?

Howard J Feldman, Paul Labute

Chemical Computing Group, Inc., 1010 Sherbrooke St. W., Suite 910, Montreal, Quebec, H3A 2R7, Canada, E-mail:feldman@chemcomp.com

Recently the Worldwide Protein Data Bank (MSD-EBI in Europe, PDBj in Japan and RCSB in USA) has made available the PDB Exchange Dictionary (http://pdbml.rcsb.org/schema/pdbx.xsd), adapted from the mmCIF dictionary. The data structures provided allow better disambiguation compared to the aging PDB format, for example introducing the concept of entities - unique molecules within the record. Biological assemblies are more easily represented as well. The PDBML format is aligned with modern relational database practices - only store each piece of information once. However the problem remains that most popular software uses the PDB format for both input and output. We have looked at some of the hurdles involved in converting PDB files to PDBML format, and back again, and present a new, scalable database system, Protein SILO (PSILO) which overcomes these. The benefits of using correctly built PDBML files include more accurate interpretation of ligands and small molecules, more precise definitions of experimental conditions, categories for structure validation measures (a protein 'health check'), and far more powerful search capabilities when stored in a relational database.

Keywords: PDBML, structural databases, biomacromolecular structures

P24.02.04

Acta Cryst. (2008). A64, C626-627

The eCrystals Federation: Management and publication of small molecule structure data for all

Simon J Coles¹, Michael B Hursthouse¹, Richard A Stephenson¹, Peter D Cliff², Elizabeth Lyon², Manjula Patel², Jim Downing³, Peter Murray-Rust³

¹University of Southampton, School of Chemistry, School of Chemistry, Southampton, Hampshire, SO17 1BJ, UK, ²UKOLN, University of Bath, UK, ³Unilever Centre for Molecular Informatics, University of Cambridge, UK, E-mail:s.j.coles@soton.ac.uk

Modern data collection methods and publication processes cause a bottleneck in the dissemination of crystal structure data, which hinders the growth of databases. There is also increasing pressure from funders to make results available as widely and rapidly as possible. We have established an institutional data repository that supports, manages and disseminates metadata relating to crystal structure data. The eCrystals Federation project (http://wiki.ecrystals.