

present in microorganisms, plants and animals. From a biotechnological point of view, *Kluyveromyces lactis*  $\beta$ -galactosidase is suitable for many applications due to its neutral optimum pH and for the fact that *K. lactis* is a GRAS organism (Generally Recognized As Safe). Interestingly,  $\beta$ -galactosidases are being used in lactose intolerance treatments and in food industry. Moreover, yeast expressing this enzyme can be used to improve the valorization of the cheese whey, a cheese industry byproduct by coupling the degradation of lactose with ethanol production, biomass production, etc. [1].

On the basis of their sequence,  $\beta$ -galactosidases are classified within families 1, 2, 35 and 42 of glycosyl hydrolases in the CAZy database [2]. Those from eukaryotic organisms are grouped into family 35 with the only exceptions of *K. lactis* and *K. marxianus*  $\beta$ -galactosidases which belong to family 2, together with the prokaryotic  $\beta$ -galactosidases from *Escherichia coli* and *Arthrobacter* sp. Whereas the structures of these last two prokaryotic enzymes have been determined [3], [4], none of the eukaryotic  $\beta$ -galactosidase structures have been reported to date. Although their sequence similarity with the prokaryotic enzymes is significant (48% vs. *E. coli* and 47% vs. *Arthrobacter*) there are many differences, particularly some long insertions and deletions, which play an important role in protein stability and in substrate recognition and specificity.

Gaining insight into the structural features that determine its stability and understanding the specificity determinants and catalytic mechanism should lead to improvements of its biotechnological applications by rational protein engineering. In this study, we describe X-ray crystallographic studies and an analysis of *Kluyveromyces lactis*  $\beta$ -galactosidase structure.

[1] M.I. Gonzalez Siso. *Bioresorcer Technology* **1996**, 57, 1-11 [2] L. Brandi Cantarel, P. M. Coutinho, C. Rancurel, Th. Bernard, V. Lombard, B. Henrissat, *Nucleic acids research* **2009**, 27, 233-238. [3] D.H. Juers, R. Jacobson, D. Wigley, X. Zhang, R.E. Huber, D. Tronrud, B.W. Matthews. *Protein Science* **2000**, 9, 1685-1699. [4] T. Skálová, J. Dohnálek, V. Spiwok, P. Lipovová, E. Vondráčková, H. Petroková, J. Dušková, H. Strnad, B. Králová, J. Hašek. *Journal of Molecular Biology* **2005**, 353, 282-294.

**Keywords:** yeast, glycosidase, biotechnology

## MS93.P37

*Acta Cryst.* (2011) A67, C781

### Structural studies of $\beta$ -D-xylosidase from *Streptomyces thermoviolaceus* OPC-520

Koji Tomoo,<sup>a</sup> Kei Saito,<sup>a</sup> Tsunego Usui,<sup>a</sup> Toshimasa Ishida,<sup>a</sup> Katsushiro Miyamoto,<sup>b</sup> Hiroshi Tsujibo,<sup>b</sup> Raita Hirose,<sup>c</sup> Kensaku Hamada,<sup>c</sup> Yuki Nakamura,<sup>d</sup> Go Ueno,<sup>e</sup> Masaki Yamamoto,<sup>e</sup> <sup>a</sup>Department of Physical Chemistry and <sup>b</sup>Department of Microbiology, Osaka University of Pharmaceutical Sciences, (Japan). <sup>c</sup>X-ray Research Laboratory, Rigaku Corporation, (Japan). <sup>d</sup>Rigaku Corporation, (Japan). <sup>e</sup>SR Life Science Instrumentation Unit, RIKEN SPring-8 Center. (Japan). E-mail: tomoo@gly.oups.ac.jp.

Xylan is one of the major hemicellulose components of plant cells and is the second most abundant resource after cellulose. *Streptomyces thermoviolaceus* OPC-520 produces relatively high xylanase activity when grown in a medium containing xylan as a carbon source.

An intracellular  $\beta$ -D-xylosidase (Bx1A) from *Streptomyces thermoviolaceus* OPC-520, together with an extra-cellular Bx1E and the integral membrane proteins Bx1F and Bx1G, constitutes a xylanolytic system that participates in the intracellular transport of xylan degradation products and the production of xylose. To elucidate the hydrolytic mechanism of xylooligosaccharides to xylose at the atomic level, we attempted X-ray structural analysis of Bx1A. The recombinant Bx1A protein (molecular weight: 82kDa) was crystallized by the hanging-drop vapor-diffusion method at 293 K. The crystals

belonged to the monoclinic space group C2, with unit-cell parameters  $a=141.0$ ,  $b=129.5$ ,  $c=100.4$  Å,  $\beta=120.1^\circ$ , and contained two molecules per asymmetric unit ( $V_M=2.47$  Å<sup>3</sup>/Da). Diffraction data were collected to a resolution to 2.2 Å on the BL26B1 beam line at the SPring-8. The brilliant light source present at SPring-8 was necessary to obtain useful data from these very weakly diffracting crystals. The initial structure of Bx1A was determined by MAD method using the SeMet-labeled crystal. The structure of Bx1A was built from typical catalytic domain and additional c-terminal domain. We now progress the refinement of structure to elucidate the detailed mechanism of glycoside hydrolysis of Bx1A at the atomic level.

**Keywords:** xylosidase, MAD phasing, crystallographic structure

## MS93.P38

*Acta Cryst.* (2011) A67, C781

### Structural study of the mutants of FDTS from *T. maritima*

Irimpan I. Mathews,<sup>a</sup> Todd Fleischmann,<sup>b</sup> Amnon Kohen,<sup>b</sup> and Scott A. Lesley,<sup>c</sup> <sup>a</sup>2575 Sand Hill Road, Stanford Synchrotron Radiation Lightsource, SLAC, Stanford University, Menlo Park, California 94025, (USA). <sup>b</sup>Department of Chemistry, University of Iowa, Iowa City, Iowa 52242, (USA). <sup>c</sup>The Joint Center for Structural Genomics at the Genomics Institute of Novartis Research Foundation, San Diego, California 92121, (USA). E-mail: iimathew@slac.stanford.edu

Like thymidylate synthase (TS) in eukaryotes, the flavin-dependent thymidylate synthase (FDTS) is essential for cell survival of many prokaryotes for the *de novo* synthesis of thymidylate (dTMP). TS and FDTS inhibition stops DNA production, eventually leading to cell death. The two enzyme families present no structural or sequence homology. While both convert dUMP to dTMP, in contrast to the classical TS enzymes, the FDTS enzymes employ 5,10-methylene-5,6,7,8-tetrahydrofolate (CH<sub>2</sub>H<sub>4</sub>folate) as the methyl donor but not as reducing agent. Furthermore, recent studies showed that, in contrast to TS, FDTS doesn't use enzymatic nucleophile for the reaction and that the catalytic mechanism of TSs and FDTSs is substantially different [1]. These differences offer the possibility of developing specific FDTS inhibitors as antibiotic drugs with low toxicity.

We report the structural study of five mutants of FDTS from *Thermotoga maritima*. All of the mutants crystallize in the orthorhombic space group P2<sub>1</sub>2<sub>1</sub>2<sub>1</sub> with unit cell dimensions of  $a\sim 55$ Å,  $b\sim 116$ Å, and  $c\sim 141$ Å. A comparison of the structures with the wild type enzyme and a comparison of the structures of the mutant enzymes with and without the substrate dUMP show several features that are important for the FDTS reaction. The structural information is also compared with the activity data to get a better understanding of the reaction mechanism.

[1] E.M. Koehn, I. Fleischmann, J.A. Conrad, B.A. Palfey, S.A. Lesley, I.I. Mathews, A. Kohen, *Nature* **2009**, 458, 919-923.

**Keywords:** flavin, thymidylate, synthase

## MS93.P39

*Acta Cryst.* (2011) A67, C781-C782

### Structure determination of levan fructotransferase

Jinseo Park, Myung-II Kim, Sangkee Rhee, Department of Agricultural Biotechnology, Seoul National University, Seoul 151-921, (Korea). E-mail: sun44@snu.ac.kr

Levan is fructan composed of linear and branched polymers of