

## MS38-P14 Combination of solid state NMR, DFT and XRPD for determination of a theophylline : 4-aminobenzoic acid cocrystal phase


José A.P. Fernandes<sup>1</sup>, Mariana Sardo<sup>1</sup>, Luís Mafra<sup>1</sup>, Duane Choquesillo-Lazarte<sup>2</sup>, Norberto Masciocchi<sup>3</sup>

1. CICECO, Universidade de Aveiro, Campus de Santiago, 3810-193 Aveiro, Portugal
2. Laboratorio de Estudios Cristalográficos, IACT, CSIC-Universidad de Granada, Av. de las Palmeras 4, E-18100 Armilla, Granada, Spain
3. Dipartimento di Scienza e Alta Tecnologia, Università dell'Insubria, 22100 Como, Italy

email: jafernandes@ua.pt

Theophylline and 4-aminobenzoic acid in 1:1 ratio selectively, and quantitatively, generate, by Liquid Assisted Grinding (LAG), a new cocrystal phase (**I**). Its structure was retrieved by conventional laboratory X-ray powder diffraction methods, using the programme TOPAS academic for solving and refining the structure. The unusual absence of hydrogen-bond interactions involving the amino groups required further validation, performed by solid state NMR and IR characterization.

The use of 1D/2D <sup>1</sup>H high-resolution solid-state NMR techniques provided structural insight on local length scales revealing internuclear proximities and relative orientations between the building blocks of the title compound, thus providing information on the type of hydrogen bond synthons formed. DFT calculations were also employed to generate meaningful structures and calculate NMR <sup>1</sup>H and <sup>13</sup>C chemical shifts to further validate the XRPD model.



**Figure 1.** a) Final Rietveld refinement plot. Blue and black bars refer to the angular positions of Bragg reflections for 1 and the minor 4-aminobenzoic acid (<5%) contaminant, respectively; b) <sup>1</sup>H MAS NMR spectrum of 1 ( $B_0=16.4$  T; 30 kHz MAS); c) Overlap of the structures obtained by XRPD and DFT.

**Keywords:** powder diffraction, solid state nuclear magnetic resonance, DFT, TOPAS, CASTEP

## MS38-P15 ADPs simulations as an additional source of fine structural information

Francesco Punzo<sup>1</sup>, Giuseppe M. Lombardo<sup>2</sup>

1. Dipartimento di Scienze del Farmaco Università degli Studi di Catania
2. Dipartimento di Scienze del Farmaco Università degli Studi di Catania

email: fpunzo@unict.it

The effect of the atomic thermal motion about their equilibrium positions,  $\mathbf{u}$ , enters in the calculation of the structure-factor, which is a function of the Miller indexes defined by the vector  $\mathbf{h}$ , as the Debye-Waller factor,  $T(\mathbf{h})$ . The calculation of the ADPs from the MD simulation ( $\mathbf{U}_{MD,cif}^{ADP}$ ) was carried out as follows: the first step involved the calculation of the symmetric tensor  $\mathbf{U}^*$  from the MD trajectory by converting, in each of the  $N$  snapshots, the atomic Cartesian coordinates ( $\mathbf{r}$ ) into fractional coordinates ( $\mathbf{r}^*$ ), and applying to the fractional coordinates the usual definitions of mean coordinates and variance-covariance matrix; thereafter, the calculation of the average values of the cell parameters and the determination of the averaged reciprocal lattice parameters lengths were carried out. These values, as calculated from the MD trajectory, were used to finally obtain  $\mathbf{U}_{MD,cif}^{ADP}$ . Apart from the possible comparison between X-ray and neutron inferred ADP data – the latter being more trustable for several physical reasons – valuable information could be obtained from their detailed analysis with a particular emphasis not only on their size but on their shape and spatial distribution. In this contribution we provide several examples – spanning, among others, from the study of polymorphism and pseudosymmetry to the energy ranking and stability of crystalline arrangements (1-4) – on extremely different chemical species thus confirming the wide pertinence of the proposed approach.

1. Lombardo G. M., Rescifina A., Punzo F. Functional hybrid co-crystals of humic substances: a growth forecast. CrystEngComm 2014, 16, 5917-5923.

2. Lombardo G. M., Punzo F. False asymmetry, pseudosymmetry, disorder, polymorphism and atomic displacement parameters. J. Mol. Struct. 2014, 1078, 158-164.

3. Lombardo G. M., Portalone G., Chiacchio U., Rescifina A., Punzo F. Potassium caffeate/caffein acid co-crystal: the rat race between cathecolic and carboxylic moiety in an atypical co-crystal. Dalton Trans., 2012, 41, 14337-14344.

4. Lombardo G. M., Thompson A. L., Ballistreri F. P., Pappalardo A., Trusso Sfrazzetto G., Tomasselli G. A., Toscano R. M., Punzo F. An integrated X-ray and Molecular Dynamics study of uranyl-salens structures and properties. Dalton Trans. 2012, 41, 1951-1960.