MS18-P05 | COMBINATION OF EXAFS AND XRD FOR STUDIES OF THE ORTHORHOMBIC-TETRAGONAL PHASE TRANSFORMATION IN MAPBI_{3-x}CL_x PEROVSKITES

Schuck, Götz (Helmholtz-Zentrum Berlin für Materialien und Energie GmbH, Berlin, GER); Többens, Daniel M. (Helmholtz-Zentrum Berlin für Materialien und Energie GmbH, Berlin, GER); Lehmann, Frederike (Helmholtz-Zentrum Berlin für Materialien und Energie GmbH, Berlin, GER); Schorr, Susan (Helmholtz-Zentrum Berlin für Materialien und Energie GmbH, Berlin, GER)

Research interest has increasingly focused on hybrid perovskites MABX₃ like $[CH_3NH_3]^+$ (MA), B = Pb and X = I or Cl as a future photovoltaic material. In QENS studies it could be shown that chlorine substitution has a large influence on the rotational dynamics of the MA molecule in MAPbl_{3-x}Cl_x perovskites. The QENS results show that chlorine substitution in the low temperature orthorhombic phase leads to a weakening of the hydrogen bridge bonds (these bonds connect the MA molecules with the $[PbX_6]^-$ octahedra host structure) since the characteristic relaxation times of C₃ rotation at 70 K in MAPbCl₃ (135 ps) and MAPbI_{2.94}Cl_{0.06} (485 ps) are much shorter than in MAPbl₃ (1635 ps).[1] The structural counterpart to the changes in the MA molecule dynamics caused by the chlorine substitution is the influence of the chlorine substitution on the tilting and distortion of the $[PbX_6]^-$ octahedra in the orthorhombic-tetragonal phase transformation temperature range. By a combination of EXAFS Pb L3 edge investigations (local structure) and XRD (long-range order) the influence of the chlorine substitution on the distortions of the $[PbX_6]^-$ octahedra was investigated. The temperature dependent XAFS measurements as well as the temperature dependent XRD measurements were performed at KMC-2 at Bessy II.

[1] G. Schuck, F. Lehmann, J. Ollivier, H. Mutka, S. Schorr, J. Phys. Chem. C (2019), DOI: 10.1021/acs.jpcc.9b01238.