

## addenda and errata

**Parameter-space screening: a powerful tool for high-throughput crystal structure determination. Corrigendum**


Zhi-Jie Liu,\* Dawei Lin, Wolfram Tempel, Jeremy L. Praissman, John P. Rose and Bi-Cheng Wang\*

Southeast Collaboratory for Structural Genomics, Department of Biochemistry and Molecular Biology, The University of Georgia, Athens, Georgia, USA. Correspondence e-mail: liu@secsg.uga.edu, wang@bcl1.bmb.uga.edu

Fig. 4 in the article by Liu *et al.* [(2005), *Acta Cryst. D* **61**, 520–527] was labelled incorrectly. A corrected version of the figure is given here. Also in §3.1.3 of the original article the Cr  $K\alpha$  wavelength was given incorrectly. It should be 2.29 Å.

**References**

Liu, Z.-J., Lin, D., Tempel, W., Praissman, J., Rose, J. P. & Wang, B.-C. (2005). *Acta Cryst. D* **61**, 520–527.

**Figure 4**

A graphical representation of pipeline success space for the PA-L1 example. A total of 55 SOLVE/RESOLVE phase sets were used as input to ARP/wARP. The colour scheme used represents success (number of residues fitted), with red indicating a near-complete model and blue/cyan representing cases where model building failed. An interesting and unexpected feature is that success space is not continuous with regions of low success sandwiched between regions of high success.