

Paula Delgado,^a Silvia Cruz,^b
Justo Cobo,^c John N. Low^d and
Christopher Glidewell^{e*}^aGrupo de Investigación de Compuestos Heterocíclicos, Departamento de Química, Universidad de Valle, AA 25360 Cali, Colombia, ^bDepartamento de Química, Universidad de Nariño, Ciudad Universitaria, Torobajo, AA 1175 Pasto, Colombia, ^cDepartamento de Química Inorgánica y Orgánica, Universidad de Jaén, 23071 Jaén, Spain, ^dDepartment of Chemistry, University of Aberdeen, Meston Walk, Old Aberdeen AB24 3UE, Scotland, and ^eSchool of Chemistry, University of St Andrews, Fife KY16 9ST, Scotland

Correspondence e-mail: cg@st-andrews.ac.uk

Key indicators



Single-crystal X-ray study
 $T = 120\text{ K}$
Mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$
Disorder in solvent or counterion
 R factor = 0.051
 wR factor = 0.141
Data-to-parameter ratio = 16.4For details of how these key indicators were automatically derived from the article, see <http://journals.iucr.org/e>.3-Methyl-4-(4-nitrophenyl)-1-phenyl-1,7-dihydro-6H-pyrazolo[3,4-*b*]thiazolo[5,4-*e*]pyridine-6-thione–dimethylformamide (1/1)The title compound is a stoichiometric solvate, $\text{C}_{20}\text{H}_{13}\text{N}_5\text{O}_2\text{S}_2 \cdot \text{C}_3\text{H}_7\text{NO}$, in which the two components are linked by an $\text{N}-\text{H} \cdots \text{O}$ hydrogen bond. The heterocyclic molecules are linked into chains by a combination of a $\text{C}-\text{H} \cdots \text{S}=\text{C}$ hydrogen bond and a $\pi-\pi$ stacking interaction.

Received 25 October 2005

Accepted 28 October 2005

Online 5 November 2005

Comment

With the aim of preparing new classes of fused thiazolo systems, we have synthesized a novel series of 5-arylmethylene-2-thioxothiazolidin-4-ones (Delgado *et al.*, 2005) as intermediates for cyclocondensation reactions. We report here the structure of 3-methyl-4-(4-nitrophenyl)-1-phenyl-1,7-dihydro-6H-pyrazolo[3,4-*b*]thiazolo[5,4-*e*]pyridin-6-thione, formed by the reaction of 5-(4-nitrobenzyliden)-2-thioxothiazolidin-4-one with 5-amino-3-methyl-1-phenylpyrazole, and crystallized as its dimethylformamide solvate, (I).The title compound (Fig. 1) is a stoichiometric solvate in which the two independent components are linked by an almost linear $\text{N}-\text{H} \cdots \text{O}$ hydrogen bond (Table 2).




Figure 1

The independent components of (I), showing displacement ellipsoids drawn at the 30% probability level. For the sake of clarity, only the major orientation of the dimethylformamide component is shown, with only one orientation of the methyl groups. The dashed line indicates a hydrogen bond.

Within the heterocyclic component, the bond distances (Table 1) indicate electronic delocalization within the pyridine ring, with strong bond fixation in the pyrazole ring. The dihedral angle between the unsubstituted phenyl ring, C11–C16, and the pyrazole ring is only $6.1(2)^\circ$, and this near planarity may be associated with the two intramolecular C–H \cdots N contacts (Table 2). On the other hand, the nitrated phenyl ring, C41–C46, makes a dihedral angle of $57.7(2)^\circ$ with the pyridine ring, thus precluding the development of quinoid forms such as (II).

There are two weak intermolecular interactions between the heterocyclic molecules, which may be of structural significance. Aryl atom C46 in the molecule at (x, y, z) acts as a hydrogen-bond donor to the thione atom S6 in the molecule at $(\frac{1}{2} + x, \frac{1}{2} - y, \frac{1}{2} + z)$, thereby forming a $C(8)$ (Bernstein *et al.*, 1995) chain running parallel to the [101] direction and generated by the n -glide plane at $y = 0.25$ (Fig. 2). This chain is reinforced by a π – π stacking interaction between the unsubstituted phenyl ring in the molecule at (x, y, z) and the pyridine ring in the molecule at $(\frac{1}{2} + x, \frac{1}{2} - y, \frac{1}{2} + z)$; these rings are nearly parallel, with a dihedral angle between them of only $5.8(2)^\circ$. The ring-centroid separation is $3.545(2) \text{ \AA}$ and the interplanar spacing is *ca* 3.41 \AA , corresponding to a ring-centroid offset of *ca* 0.97 \AA . Two [101] chains pass through each unit cell, but there are no direction-specific interactions between adjacent chains.

Experimental

A solution containing equimolar quantities (1 mmol of each component) of 5-amino-3-methyl-1-phenylpyrazole and 5-(4-nitrobenzylidene)-2-thioxothiazolidin-4-one in dimethylformamide (DMF; 3 ml) was heated under reflux for 6 h. The reaction mixture was cooled, and ethanol was added before the combined solvents were removed under reduced pressure. The resulting solid product was recrystallized from DMF to yield **orange** crystals suitable for single-

crystal X-ray diffraction. Yield 90%; m.p. 530 K. MS (70 eV) m/z (%) 421 (13, $M^+ + 2$); 420 (28, $M^+ + 1$); 419 (100, M^+); 372 (6).

Crystal data

$C_{20}H_{13}N_5O_2S_2 \cdot C_3H_7NO$
 $M_r = 492.57$
 Monoclinic, $P2_1/n$
 $a = 9.3023(2) \text{ \AA}$
 $b = 25.5447(7) \text{ \AA}$
 $c = 10.3794(2) \text{ \AA}$
 $\beta = 113.7660(13)^\circ$
 $V = 2257.25(9) \text{ \AA}^3$
 $Z = 4$

$D_x = 1.449 \text{ Mg m}^{-3}$
 Mo $K\alpha$ radiation
 Cell parameters from 5163 reflections
 $\theta = 3.4\text{--}27.6^\circ$
 $\mu = 0.28 \text{ mm}^{-1}$
 $T = 120(2) \text{ K}$
 Block, yellow
 $0.62 \times 0.44 \times 0.22 \text{ mm}$

Data collection

Bruker–Nonius KappaCCD diffractometer
 φ and ω scans
 Absorption correction: multi-scan (SADABS; Sheldrick, 2003)
 $T_{\min} = 0.848$, $T_{\max} = 0.942$
 25526 measured reflections

5163 independent reflections
 4088 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.004$
 $\theta_{\text{max}} = 27.6^\circ$
 $h = -12 \rightarrow 12$
 $k = -32 \rightarrow 33$
 $l = -13 \rightarrow 13$

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.051$
 $wR(F^2) = 0.141$
 $S = 1.04$
 5163 reflections
 315 parameters
 H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0667P)^2 + 2.3558P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 1.12 \text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.80 \text{ e \AA}^{-3}$

Table 1

Selected geometric parameters (\AA , $^\circ$).

N1–N2	1.383 (3)	N7–C7A	1.380 (3)
N2–C3	1.322 (3)	C7A–N8	1.328 (3)
C3–C3A	1.433 (3)	N8–C8A	1.339 (3)
C3A–C4	1.408 (3)	C8A–N1	1.373 (3)
C4–C4A	1.387 (3)	C3A–C8A	1.412 (3)
C4A–S5	1.745 (2)	C4A–C7A	1.414 (3)
S5–C6	1.749 (2)	C6–S6	1.656 (2)
C6–N7	1.350 (3)		
N2–N1–C11–C12	3.8 (3)	C3A–C4–C41–C42	128.2 (2)
C8A–N1–C11–C12	–174.1 (2)	C4A–C4–C41–C42	–56.1 (3)
N2–N1–C11–C16	–177.4 (2)	C3A–C4–C41–C46	–56.6 (3)
C8A–N1–C11–C16	4.8 (4)	C4A–C4–C41–C46	119.1 (2)

Table 2

Hydrogen-bond geometry (\AA , $^\circ$).

$D\text{--}H\cdots A$	$D\text{--}H$	$H\cdots A$	$D\cdots A$	$D\text{--}H\cdots A$
N7–H7 \cdots O21	0.88	1.84	2.711 (3)	170
C12–H12 \cdots N2	0.95	2.41	2.758 (4)	102
C16–H16 \cdots N8	0.95	2.32	2.975 (4)	126
C46–H46 \cdots S6 ⁱ	0.95	2.80	3.743 (3)	176

Symmetry code: (i) $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$.

All H atoms were located in difference maps and subsequently treated as riding atoms with C–H = 0.95 (CH) or 0.98 \AA (CH₃), and N–H = 0.88 \AA , and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$, or $1.5U_{\text{eq}}(\text{C})$ for the methyl groups. It was apparent from an early stage that the methyl groups of the dimethylformamide component were disordered over two sets of sites, corresponding to two conformations of this molecule. This disorder was modelled using common sites for the N–




Figure 2

Stereoview of part of the crystal structure of (I), showing the formation of a [101] chain built from C—H...S=C hydrogen bonds (dashed lines) and π - π stacking interactions. For the sake of clarity, the solvent molecules and the H atoms not involved in the motif shown have been omitted.

CHO fragment in the two conformations and two distinct sets of sites for the methyl C atoms; the refined values of the site occupancy factors were 0.598 (8) and 0.402 (8). In addition, it was necessary to model each of the methyl groups using six half-occupancy H-atom sites, offset from one another by 60°. The crystals of (I) were very fragile, and attempts to cut small fragments from larger crystals consistently led to the shattering of the crystals. The highest peak in the difference map is located 1.05 Å from one of the methyl H atoms in the minor orientation of the disordered solvent component.

Data collection: *COLLECT* (Hooft, 1999); cell refinement: *DENZO* (Otwinowski & Minor, 1997) and *COLLECT*; data reduction: *DENZO* and *COLLECT*; program(s) used to solve structure: *OSCAIL* (McArdle, 2003) and *SHELXS97* (Sheldrick, 1997);

program(s) used to refine structure: *OSCAIL* and *SHELXL97* (Sheldrick, 1997); molecular graphics: *PLATON* (Spek, 2003); software used to prepare material for publication: *SHELXL97* and *PRPKAPPA* (Ferguson, 1999).

X-ray data were collected at the EPSRC X-ray Crystallographic Service, University of Southampton, England. JC thanks the Consejería de Innovación, Ciencia y Empresa (Junta de Andalucía, Spain) and the Universidad de Jaén for financial support. PD and SC thank COLCIENCIAS, UNIVALLE (Universidad del Valle, Colombia) and Universidad de Nariño for financial support.

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Delgado, P., Quiroga, J., Cobo, J., Low, J. N. & Glidewell, C. (2005). *Acta Cryst.* **C61**, o477–o482.
- Ferguson, G. (1999). *PRPKAPPA*. University of Guelph, Canada.
- Hooft, R. W. W. (1999). *COLLECT*. Nonius BV, Delft, The Netherlands.
- McArdle, P. (2003). *OSCAIL* for Windows. Version 10. Crystallography Centre, Chemistry Department, NUI Galway, Ireland.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography, Part A*, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
- Sheldrick, G. M. (2003). *SADABS*. Version 2.10. University of Göttingen, Germany.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.