

1-Isopropyl-6,6,8a-trimethyl-1,3a,5,6,7,8a-hexahydro-3H-1-benzofuro[2,3-b]pyrrole-2,4-dione

R. N. Narasegowda,^a H. S. Yathirajan,^a D. E. Lynch,^b T. Narasimhamurthy^c and R. S. Rathore^{d*}

^aDepartment of Studies in Chemistry, University of Mysore, Mysore 570 006, India, ^bFaculty of Health and Life Sciences, Coventry University, Coventry CV1 5FB, England, ^cIndian Institute of Science, Bangalore 560 012, India, and ^dOriental Organization of Molecular and Structural Biology, 204, Agarwal Bhavan, Malleshwaram, Bangalore 560 055, India

Correspondence e-mail:
ravindranath_rathore@yahoo.com

Key indicators


Single-crystal X-ray study
 $T = 120$ K
 Mean $\sigma(\text{C}-\text{C}) = 0.002$ Å
 R factor = 0.039
 wR factor = 0.101
 Data-to-parameter ratio = 10.7

For details of how these key indicators were automatically derived from the article, see <http://journals.iucr.org/e>.

The crystal structure of the title benzofuran derivative, $\text{C}_{16}\text{H}_{23}\text{NO}_3$, has been elucidated. The tricyclic core, *i.e.* the tetrahydrobenzo-dihydrofuro-pyrrolidine ring system, is non-planar owing to the folding of the five-membered rings at their *cis* junction. The cyclohexene ring assumes a half-chair conformation, while the dihydrofuran and pyrrolidine rings each adopt an envelope conformation. Intramolecular C—H \cdots O hydrogen bonds form *S*(6) closed patterns.

Comment

The title compound, (I), has been shown to exhibit a moderate hypoglycemic activity in a previous structure–activity relationship study (Nagarajan *et al.*, 1988). Compound (I) is a new tricyclic benzofuran derivative containing linearly fused tetrahydrobenzo-dihydrofuro-pyrrolidine (*A–B–C*) rings. This chiral molecule formally derives from a perhydro-furo (or -pyrrolo)-benzofuran system (Nagarajan *et al.*, 1988) and is structurally related to a structure containing a tetrahydrobenzo-furo-furan ring system, which we recently published (Nagaraj *et al.*, 2005).



The molecular structure is shown in Fig. 1. The *BC* ring-junction is *cis* (Bucourt, 1974). The shape of the tricyclic core is non-planar owing to the folding at the *BC* junction. The torsion angles at this junction, namely $\text{N1}-\text{C1}-\text{C4}-\text{C5}$ and $\text{O1}-\text{C1}-\text{C4}-\text{C3}$, are -99.69 (11) and 132.78 (11) $^\circ$, respectively. The structure of the analogous molecule based on a chiral tetrahydrobenzo-furo-furan core (Nagaraj *et al.*, 2005) also has a non-planar shape for its tricyclic core, and the equivalent torsion angles are 103.98 (10) and -127.17 (10) $^\circ$, respectively. The torsion angle $\text{C1}-\text{N1}-\text{C14}-\text{C15}$ in (I), describing the conformation of the *N*-isopropyl substituent, is 121.51 (14) $^\circ$. The internal torsion angles of individual rings are shown in Fig. 1. Ring *A* (cyclohexene) adopts a half-chair (C_2) conformation (Bucourt, 1974) with the following values of puckering parameters (Cremer & Pople, 1975): $q_2 = 0.347$ (2), $q_3 = -0.275$ (2) Å, $\varphi_2 = 345.7$ (3), $\theta_2 = 128.4$ (2) $^\circ$ and $Q = 0.443$ (2) Å. Rings *B* and *C* adopt envelope (C_s) conforma-

Received 23 December 2005
 Accepted 20 February 2006




Figure 1

The molecular structure of (I). Displacement ellipsoids are drawn at the 30% probability level and numerical values refer to the internal torsion angles ($^{\circ}$) of individual rings (s.u. values lie in the range 0.1–0.2 $^{\circ}$). Intramolecular C—H \cdots O hydrogen bonds are displayed with dashed lines.

tions (Fuchs, 1978) with atoms C1 and C4, respectively, at the flap positions. Atoms C1 and C4 are 0.24 (1) and 0.33 (1) Å out of the mean planes formed by the remaining ring atoms. The puckering amplitudes q (Å) and the phase angles φ ($^{\circ}$) of the five-membered rings are 0.146 (2) and 39.7 (7), and 0.210 (2) and 80.3 (5), respectively. Two $S(6)$ hydrogen-bonded closed patterns (Bernstein *et al.*, 1995) are formed by C15—H \cdots O2 and C16—H \cdots O2 intramolecular contacts (Table 1). The lengths and directionality suggest that these hydrogen bonds are very weak. The crystal packing is entirely due to van der Waals interactions.

Experimental

The synthesis of (I) was described by Nagarajan *et al.* (1988). Suitable single crystals were obtained by slow evaporation of a benzene–hexane (1:1) solution.

Crystal data

| | |
|-----------------------------------|---|
| $C_{16}H_{23}NO_3$ | $Z = 2$ |
| $M_r = 277.35$ | $D_x = 1.237 \text{ Mg m}^{-3}$ |
| Triclinic, $P\bar{1}$ | Mo $K\alpha$ radiation |
| $a = 8.9725$ (3) Å | Cell parameters from 3368 reflections |
| $b = 10.2472$ (3) Å | $\theta = 2.9\text{--}27.5^{\circ}$ |
| $c = 10.4372$ (3) Å | $\mu = 0.09 \text{ mm}^{-1}$ |
| $\alpha = 101.733$ (2) $^{\circ}$ | $T = 120$ (2) K |
| $\beta = 109.290$ (1) $^{\circ}$ | Plate, colourless |
| $\gamma = 115.765$ (1) $^{\circ}$ | $0.22 \times 0.18 \times 0.05 \text{ mm}$ |
| $V = 744.67$ (4) Å 3 | |

Data collection

| | |
|---|--|
| Nonius KappaCCD diffractometer | 2548 reflections with $I > 2\sigma(I)$ |
| φ and ω scans | $R_{\text{int}} = 0.038$ |
| Absorption correction: multi-scan (SADABS; Sheldrick, 2003) | $\theta_{\text{max}} = 26.0^{\circ}$ |
| $T_{\text{min}} = 0.890$, $T_{\text{max}} = 0.996$ | $h = -11 \rightarrow 11$ |
| 14657 measured reflections | $k = -12 \rightarrow 12$ |
| 2911 independent reflections | $l = -12 \rightarrow 12$ |

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.101$
 $S = 1.09$
 2911 reflections
 273 parameters
 All H-atom parameters refined

$$w = 1/[\sigma^2(F_o^2) + (0.0394P)^2 + 0.3291P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.21 \text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.26 \text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (Å, $^{\circ}$).

| $D\text{—}H\cdots A$ | $D\text{—}H$ | $H\cdots A$ | $D\cdots A$ | $D\text{—}H\cdots A$ |
|----------------------|--------------|-------------|-------------|----------------------|
| C15—H153 \cdots O2 | 0.99 (3) | 2.54 (2) | 3.122 (2) | 118 (2) |
| C16—H161 \cdots O2 | 1.02 (3) | 2.58 (2) | 3.186 (2) | 118 (1) |

Larger than expected values of residual electron density were observed and attributed to the presence of a few poorly fitting reflections ($\bar{1}22$, $\bar{1}\bar{1}1$, 011 , $2\bar{2}\bar{1}$, $\bar{2}22$ and $\bar{1}\bar{1}1$). The application of an extinction correction [extinction parameter = 0.57 (3)] further degraded the model quality. In the absence of any obvious cause, these reflections were omitted during the last cycles of refinement. Residual electron density was then featureless and the residual factor R dropped from 0.056 to 0.039 for observed data. H atoms were located in a difference map and were refined freely [$C\text{—}H = 0.97$ (2)–1.02 (2) Å].

Data collection: *COLLECT* (Hooft, 1998); cell refinement: *DENZO* (Otwinowski & Minor, 1997) and *COLLECT*; data reduction: *DENZO* and *COLLECT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *PLATON* (Spek, 2003); software used to prepare material for publication: *SHELXL97* and *PLATON*.

The authors thank the EPSRC National Crystallography Service, Southampton, England.

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bucourt, R. (1974). *Topics in Stereochemistry*, edited by E. L. Eliel & N. L. Allinger, Vol. 8, pp. 159–224. New York: Interscience.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Fuchs, B. (1978). *Topics in Stereochemistry*, edited by E. L. Eliel & N. L. Allinger, Vol. 10, pp. 1–94. New York: Interscience.
- Hooft, R. W. W. (1998). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Nagarajan, K., Talwalker, P. K., Nagana Goud, A., Shah, R. K., Shenoy, S. J. & Desai, N. D. (1988). *Indian J. Chem. Sect. B*, **27**, 1113–1123.
- Nagaraj, B., Yathirajan, H. S., Nagaraja, P. & Lynch, D. E. (2005). *Acta Cryst. E61*, o1041–o1042.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
- Sheldrick, G. M. (2003). *SADABS*. Version 2.10. Bruker AXS Inc., Madison, Wisconsin, USA.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.