

Kevin R. Flower* and Robin G. Pritchard

 School of Chemistry, University of Manchester,
 Sackville Street, Manchester, England

 Correspondence e-mail:
 k.r.flower@manchester.ac.uk

Key indicators

 Single-crystal X-ray study
 $T = 293\text{ K}$
 Mean $\sigma(\text{C}-\text{C}) = 0.011\text{ \AA}$
 R factor = 0.049
 wR factor = 0.092
 Data-to-parameter ratio = 18.0

 For details of how these key indicators were
 automatically derived from the article, see
<http://journals.iucr.org/e>.

Bis[2-(2-methylphenylimino)phenyl]mercury(II)


 The structure of the cyclomercurated 2-phenyliminophenyl title compound, $[\text{Hg}(\text{C}_{14}\text{H}_{12}\text{N})_2]$, shows that the mercury coordination is essentially square planar

Received 19 May 2006

Accepted 29 May 2006

Comment

The structure of the title compound, (I), is shown in Fig. 1. Organomercurials are often used as transmetallation reagents in the synthesis of organometallic complexes (Roper & Wright, 1977). Several years ago we reported a synthetic route for the preparation of a range of functionalized 1-mercurio-2-phenyliminophenyls (Flower *et al.*, 2002) and from the structural data obtained concurred with a previous report of Batsanov (1998) that the van der Waals radius of mercury is in the range 2.0–2.2 Å, rather than the often quoted value of 1.55 Å (Bondi, 1964). Here, and in the following paper (Flower & Pritchard, 2006), we report two additional structures of this type of compound. All of the bond lengths and angles in the two structures are as expected. The Hg–N distances in (I) and bis-2-(2-isopropylphenyliminophenyl)-mercury, (II), range from 2.787 (10) to 2.850 (10) Å and are comfortably within the sum of the van der Waals radii (3.5–3.7 Å), if the van der Waals radius of Hg is considered to be 2.0–2.2 Å, indicating significant Hg–N interactions. This gives rise to an overall distorted square-planar geometry at Hg in both cases. Other examples of square planar Hg^{II} complexes are known (Balasubramani *et al.*, 2005; Haid *et al.*, 2003; Cheng *et al.*, 1994).


Experimental

Caution: preparation of an organomercurial. Organomercurials are extremely toxic. To $\text{Hg}(\text{C}_6\text{H}_4\text{-2-CHO})_2$ (1 g, 2.4 mmol) dissolved in ethanol (10 ml) containing *p*-toluenesulfonic acid (10 mg, 0.05 mmol) was added 2-methylaniline (0.56 g, 6 mmol) and the solution was refluxed for 5 h, during which time white crystals of (I) precipitated.

The crystalline material was collected by filtration, washed with water and dried in a desiccator. Yield 0.93 g, 68%. An analytically pure sample was obtained through recrystallization from hot ethanol, and crystals suitable for the diffraction study were grown by dissolving approximately 10 mg of (I) in CH_2Cl_2 (0.2 ml) in a small vial (1 × 5 cm), layering ethanol (5 ml) on top and leaving the vial to stand for 24 h. Elemental analysis $\text{C}_{28}\text{H}_{24}\text{HgN}_2$ requires: C 57.56, H 4.11, N 4.76%; found: C 57.79, H 4.22, N 4.91%.

Crystal data

$[\text{Hg}(\text{C}_{14}\text{H}_{12}\text{N})_2]$	$Z = 4$
$M_r = 589.08$	$D_x = 1.736 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 11.9925 (3) \text{ \AA}$	$\mu = 6.85 \text{ mm}^{-1}$
$b = 11.3864 (3) \text{ \AA}$	$T = 293 (2) \text{ K}$
$c = 16.6542 (5) \text{ \AA}$	Plate, yellow
$\beta = 97.6730 (10)^\circ$	$0.2 \times 0.15 \times 0.05 \text{ mm}$
$V = 2253.79 (11) \text{ \AA}^3$	

Data collection

Nonius KappaCCD diffractometer	14711 measured reflections
φ and ω scans	5084 independent reflections
Absorption correction: multi-scan (<i>SORTAV</i> ; Blessing, 1995)	3476 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.254$, $T_{\max} = 0.707$	$R_{\text{int}} = 0.086$
	$\theta_{\text{max}} = 27.4^\circ$

Refinement

Refinement on F^2	$w = 1/[\sigma^2(F_o^2) + (0.0284P)^2 + 5.7069P]$
$R[F^2 > 2\sigma(F^2)] = 0.050$	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.092$	$(\Delta/\sigma)_{\text{max}} = 0.001$
$S = 1.02$	$\Delta\rho_{\text{max}} = 1.54 \text{ e \AA}^{-3}$
5084 reflections	$\Delta\rho_{\text{min}} = -1.64 \text{ e \AA}^{-3}$
283 parameters	Extinction correction: <i>SHELXL97</i>
H-atom parameters constrained	Extinction coefficient: 0.00103 (15)

H atoms were positioned geometrically and treated as riding, with C—H = 0.93 and 0.96 Å, and with $U_{\text{iso}}(\text{H})$ values of 1.2 and 1.5 times $U_{\text{eq}}(\text{C})$. The highest residual peak is located 1.03 Å from Hg1 and deepest hole is located 0.92 Å from Hg1.

Data collection: *COLLECT* (Nonius, 2000); cell refinement: *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *SCALEPACK* and *DENZO* (Otwinowski & Minor, 1997);




Figure 1

The molecular structure of (I), showing the atomic numbering scheme. Displacement ellipsoids are shown at the 30% probability level.

program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

We thank the EPSRC Crystallographic Service, University of Southampton, for collecting the data.

References

- Balasubramani, K., Thomas, P., Bocelli, G. & Cantoni, A. (2005). *J. Coord. Chem.* **58**, 1689–1694.
- Batsanov, A. S. (1998). *J. Chem. Soc. Dalton Trans.* pp. 1541–1546.
- Blessing, R. H. (1995). *Acta Cryst.* **A51**, 33–38.
- Bondi, A. (1964). *J. Phys. Chem.* **68**, 441–451.
- Cheng, Y., Emge, T. J. & Brennan, J. G. (1994). *Inorg. Chem.* **33**, 3711–3714.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–888.
- Flower, K. R., Howard, V. J., Naguthney, S., Pritchard, R. G., Warren, J. E. & McGown, A. T. (2002). *Inorg. Chem.* **41**, 1907–1912.
- Flower, K. R. & Pritchard, R. G. (2006). *Acta Cryst.* **E62**, m1469–m1470.
- Haid, R., Gutmann, R., Czermak, G., Langes, C., Oberhauser, W., Kopacka, H., Ongania, K.-H. & Bruggeller, P. (2003). *Inorg. Chem. Commun.* **6**, 61–67.
- Nonius (2000). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Roper, W. R. & Wright, L. J. (1977). *J. Organomet. Chem.* **142**, C1–C6.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.