

Jaime Portilla,^a José M. de la Torre,^b Justo Cobo,^b John N. Low^c and Christopher Glidewell^{d*}

^aGrupo de Investigación de Compuestos Heterocíclicos, Departamento de Química, Universidad de Valle, AA 25360 Cali, Colombia, ^bDepartamento de Química Inorgánica y Orgánica, Universidad de Jaén, 23071 Jaén, Spain, ^cDepartment of Chemistry, University of Aberdeen, Meston Walk, Old Aberdeen AB24 3UE, Scotland, and ^dSchool of Chemistry, University of St Andrews, St Andrews, Fife KY16 9ST, Scotland

Correspondence e-mail: cg@st-andrews.ac.uk

Key indicators

Single-crystal X-ray study
 $T = 120\text{ K}$
 $\text{Mean } \sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$
 $R \text{ factor} = 0.045$
 $wR \text{ factor} = 0.121$
Data-to-parameter ratio = 16.3

For details of how these key indicators were automatically derived from the article, see <http://journals.iucr.org/e>.


6-(2-Hydroxybenzoyl)-2-(4-methylphenyl)-5-methyl-7-phenylpyrazolo[1,5-a]pyrimidine: complex sheets built from $\text{C}-\text{H}\cdots\text{N}$ and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds

Received 29 June 2006
Accepted 29 June 2006

The molecules of the title compound, $C_{27}H_{21}N_3O_2$, contain an intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond, and they are linked into complex sheets by a combination of two $\text{C}-\text{H}\cdots\text{N}$ hydrogen bonds and two $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds.

Comment


We report here the structure of the title new pyrazolo[1,5-a]pyrimidine, (I) (Fig. 1), prepared under solvent-free conditions from the reaction between a 5-aminopyrazole and a 2-benzoylchromone.




The bond distances in the pyrazolo[1,5-a]pyrimidine fragment (Table 1) are typical of this ring system (Portilla *et al.*, 2005, 2006) and they are consistent with 10- π electron delocalization reminiscent of the naphthalene type. The aryl ring at C2 is almost coplanar with the heterocyclic system, but the substituents at C6 and C7 are considerably twisted out of this plane, probably to avoid mutually repulsive interactions between their aryl rings (Table 1).

There is a fairly short intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond, but the supramolecular aggregation of (I) depends solely on rather long, and thus fairly weak, $\text{C}-\text{H}\cdots\text{N}$ and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds (Table 2), which link the molecules into complex sheets. The formation of the sheet can be readily analysed in terms of its simple sub-structures.


Aryl atom C74 in the molecule at (x, y, z) acts as hydrogen-bond donor to hydroxyl atom O62 in the molecule at $(1 - x, -y, 1 - z)$, so generating by inversion a centrosymmetric $R_2^2(22)$ (Bernstein *et al.*, 1995) dimer centred at $(\frac{1}{2}, 0, \frac{1}{2})$ (Fig. 2), and these dimers are linked by the $\text{C}-\text{H}\cdots\text{N}$ hydrogen bonds. Atoms C3 and C22 in the molecule at (x, y, z) and $(1 - x, -y, 1 - z)$ both act as hydrogen-bond donors to the atoms N4 in the molecules at $(\frac{3}{2} - x, \frac{1}{2} + y, \frac{3}{2} - z)$ and $(-\frac{1}{2} + x, -\frac{1}{2} - y, -\frac{1}{2} + z)$, which themselves lie in the $R_2^2(22)$ dimers centred at $(1, \frac{1}{2}, 1)$.

**Figure 1**

The molecular structure of compound (I), showing the atom-labelling scheme and the intramolecular $O-H\cdots O$ hydrogen bond (dashed line). Displacement ellipsoids are drawn at the 30% probability level.

**Figure 2**

Part of the crystal structure of compound (I), showing the formation of an $R_2^2(22)$ dimer. For the sake of clarity, H atoms bonded to C atoms but not involved in the motifs shown have been omitted. Similarly, the unit-cell outline has been omitted. Atoms marked with an asterisk (*) are at the symmetry position ($1-x, -y, 1-z$).

**Figure 3**

A stereoscopic view of part of the crystal structure of compound (I), showing the formation of a chain of rings along [111]. For the sake of clarity, H atoms bonded to C atoms but not involved in the motifs shown have been omitted.

and $(0, -\frac{1}{2}, 0)$, respectively. Propagation of these hydrogen bonds then generates a chain of rings along [111], in which $R_2^2(22)$ rings alternate with $R_2^1(7)$ rings (Fig. 3).

Finally, aryl atom C76 in the molecule at (x, y, z) acts as hydrogen-bond donor to carbonyl atom O67 in the molecule at $(x, 1+y, z)$, so generating by translation a $C(7)$ chain running parallel to the [010] direction. The combination of [010] and [111] chains generates a sheet parallel to $(10\bar{1})$, but there are no direction-specific interactions between adjacent sheets.

Experimental

Equimolar quantities (1.0 mmol of each component) of 3-(4-methylphenyl)-5-amino-1*H*-pyrazole and 3-benzoyl-2-methylbenzo-4-pyrone were placed in an open Pyrex glass vessel and heated in an oil-bath at 373 K for 2 min. The reaction mixture was then cooled and extracted with ethanol. After removal of the solvent, the product, (I), was recrystallized from dimethylformamide to give yellow crystals suitable for single-crystal X-ray diffraction (m.p. 471–473 K, yield 60%).

Crystal data

$C_{27}H_{21}N_3O_2$	$Z = 8$
$M_r = 419.47$	$D_x = 1.338 \text{ Mg m}^{-3}$
Monoclinic, $C2/c$	Mo $K\alpha$ radiation
$a = 38.7649 (11) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$b = 6.7864 (2) \text{ \AA}$	$T = 120 (2) \text{ K}$
$c = 16.7002 (5) \text{ \AA}$	Lath, yellow
$\beta = 108.599 (2)^\circ$	$0.80 \times 0.50 \times 0.20 \text{ mm}$
$V = 4163.9 (2) \text{ \AA}^3$	

Data collection

Bruker Nonius KappaCCD area-detector diffractometer	30599 measured reflections
φ and ω scans	4751 independent reflections
Absorption correction: multi-scan (SADABS; Sheldrick, 2003)	3797 reflections with $I > 2\sigma(I)$
$R_{\text{int}} = 0.036$	
$\theta_{\text{max}} = 27.5^\circ$	
$T_{\min} = 0.964$, $T_{\max} = 0.983$	

Refinement

Refinement on F^2	$w = 1/[\sigma^2(F_o^2) + (0.0575P)^2 + 3.7732P]$
$R[F^2 > 2\sigma(F^2)] = 0.045$	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.121$	$(\Delta/\sigma)_{\text{max}} = 0.001$
$S = 1.02$	$\Delta\rho_{\text{max}} = 0.33 \text{ e \AA}^{-3}$
4751 reflections	$\Delta\rho_{\text{min}} = -0.25 \text{ e \AA}^{-3}$
291 parameters	
H-atom parameters constrained	

Table 1
Selected geometric parameters (\AA , $^\circ$).

N1—C2	1.3505 (18)	C5—C6	1.4342 (19)
C2—C3	1.4013 (19)	C6—C7	1.376 (2)
C3—C3A	1.377 (2)	C7—N7A	1.3675 (18)
C3A—N4	1.3525 (18)	N7A—N1	1.3603 (16)
N4—C5	1.3141 (18)	C3A—N7A	1.4011 (17)
N1—C2—C21—C22		C6—C67—C61—C62	
C5—C6—C67—C61		−176.77 (13)	
−110.80 (15)		C6—C7—C71—C72	
		43.8 (2)	

Table 2
Hydrogen-bond geometry (\AA , $^\circ$).

$D\cdots H\cdots A$	$D\cdots H$	$H\cdots A$	$D\cdots A$	$D\cdots H\cdots A$
O62—H62 \cdots O67	0.96	1.68	2.554 (2)	150
C3—H3 \cdots N4 ⁱ	0.95	2.61	3.543 (2)	168
C22—H22 \cdots N4 ⁱ	0.95	2.57	3.513 (2)	174
C74—H74 \cdots O62 ⁱⁱ	0.95	2.55	3.422 (2)	154
C76—H76 \cdots O67 ⁱⁱⁱ	0.95	2.56	3.444 (2)	155

Symmetry codes: (i) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{3}{2}$; (ii) $-x + 1, -y, -z + 1$; (iii) $x, y + 1, z$.

All H atoms were located in a difference map and then treated as riding, with C—H distances of 0.95 (aromatic) or 0.98 \AA (methyl) and an O—H distance of 0.96 \AA , and with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C,O})$, where $x = 1.5$ for the methyl groups and 1.2 for all other H.

Data collection: *COLLECT* (Nonius, 1999); cell refinement: *DENZO* (Otwinowski & Minor, 1997) and *COLLECT*; data reduction: *DENZO* and *COLLECT*; program(s) used to solve structure: *SIR2004* (Burla *et al.*, 2005); program(s) used to refine structure: *OSCAIL* (McArdle, 2003) and *SHELXL97* (Sheldrick, 1997); molecular graphics: *PLATON* (Spek, 2003); software used to prepare material for publication: *SHELXL97* and *PRPKAPPA* (Ferguson, 1999).

X-ray data were collected at the EPSRC X-ray Crystallographic Service, University of Southampton, UK. JC and JT

thank the Consejería de Innovación, Ciencia y Empresa (Junta de Andalucía, Spain) and the Universidad de Jaén for financial support. JT also thanks the Universidad de Jaén for a research scholarship supporting a short stay at the EPSRC X-ray Crystallographic Service, University of Southampton, UK. JP thanks COLCIENCIAS, UNIVALLE (Universidad del Valle, Colombia), for financial support.

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
 Burla, M. C., Caliandro, R., Camalli, M., Carrozzini, B., Cascarano, G. L., De Caro, L., Giacovazzo, C., Polidori, G. & Spagna, R. (2005). *J. Appl. Cryst.* **38**, 381–388.
 Ferguson, G. (1999). *PRPKAPPA*. University of Guelph, Canada.
 McArdle, P. (2003). *OSCAIL for Windows*. Version 10. Crystallography Centre, Chemistry Department, NUI Galway, Ireland.
 Nonius (1999). *COLLECT*. Nonius BV, Delft, The Netherlands.
 Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
 Portilla, J., Quiroga, J., Cobo, J., Low, J. N. & Glidewell, C. (2005). *Acta Cryst. C* **61**, o452–o456.
 Portilla, J., Quiroga, J., Cobo, J., Low, J. N. & Glidewell, C. (2006). *Acta Cryst. C* **62**, o186–o189.
 Sheldrick, G. M. (1997). *SHELXL97*. University of Göttingen, Germany.
 Sheldrick, G. M. (2003). *SADABS*. Version 2.10. University of Göttingen, Germany.
 Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

supporting information

Acta Cryst. (2006). E62, o3195–o3197 [https://doi.org/10.1107/S1600536806025220]

6-(2-Hydroxybenzoyl)-2-(4-methylphenyl)-5-methyl-7-phenylpyrazolo[1,5-a]pyrimidine: complex sheets built from C—H…N and C—H…O hydrogen bonds

Jaime Portilla, José M. de la Torre, Justo Cobo, John N. Low and Christopher Glidewell

6-(2-Hydroxybenzoyl)-2-(4-methylphenyl)-5-methyl-7-phenylpyrazolo[1,5-a]pyrimidine

Crystal data

$C_{27}H_{21}N_3O_2$
 $M_r = 419.47$
Monoclinic, $C2/c$
Hall symbol: -C 2yc
 $a = 38.7649 (11)$ Å
 $b = 6.7864 (2)$ Å
 $c = 16.7002 (5)$ Å
 $\beta = 108.599 (2)^\circ$
 $V = 4163.9 (2)$ Å³
 $Z = 8$

$F(000) = 1760$
 $D_x = 1.338 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 4751 reflections
 $\theta = 2.5\text{--}27.5^\circ$
 $\mu = 0.09 \text{ mm}^{-1}$
 $T = 120 \text{ K}$
Lath, yellow
 $0.80 \times 0.50 \times 0.20 \text{ mm}$

Data collection

Bruker Nonius KappaCCD area-detector diffractometer
Radiation source: Bruker Nonius FR591 rotating anode
Graphite monochromator
Detector resolution: 9.091 pixels mm⁻¹
 φ and ω scans
Absorption correction: multi-scan (SADABS; Sheldrick, 2003)

$T_{\min} = 0.964$, $T_{\max} = 0.983$
30599 measured reflections
4751 independent reflections
3797 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$
 $\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 2.5^\circ$
 $h = -50 \rightarrow 50$
 $k = -8 \rightarrow 8$
 $l = -20 \rightarrow 21$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.121$
 $S = 1.02$
4751 reflections
291 parameters
0 restraints
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
Hydrogen site location: inferred from neighbouring sites
H-atom parameters constrained
 $w = 1/\sigma^2(F_o^2) + (0.0575P)^2 + 3.7732P$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.33 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.25 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. MS (70 eV) m/z (%): 420 (29), 419 (89, M^+), 418 (40), 404 (19), 298 (42), 299 (100), 300 (23), 77 (19), 39 (29).

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.64536 (3)	0.64378 (18)	0.73713 (7)	0.0241 (3)
C2	0.67725 (4)	0.7409 (2)	0.76857 (8)	0.0225 (3)
C21	0.67888 (4)	0.9261 (2)	0.81559 (8)	0.0228 (3)
C22	0.71209 (4)	1.0247 (2)	0.84866 (9)	0.0277 (3)
C23	0.71402 (4)	1.2014 (2)	0.89185 (9)	0.0289 (3)
C24	0.68331 (4)	1.2854 (2)	0.90335 (9)	0.0282 (3)
C241	0.68582 (5)	1.4727 (2)	0.95313 (11)	0.0382 (4)
C25	0.65025 (4)	1.1876 (2)	0.86945 (10)	0.0299 (3)
C26	0.64783 (4)	1.0101 (2)	0.82675 (9)	0.0268 (3)
C3	0.70596 (4)	0.6500 (2)	0.74876 (9)	0.0238 (3)
C3A	0.69070 (4)	0.4875 (2)	0.70129 (8)	0.0226 (3)
N4	0.70502 (3)	0.34454 (17)	0.66512 (7)	0.0234 (3)
C5	0.68290 (4)	0.2098 (2)	0.62001 (9)	0.0240 (3)
C51	0.69940 (4)	0.0567 (2)	0.57839 (10)	0.0310 (3)
C6	0.64457 (4)	0.2128 (2)	0.60786 (9)	0.0240 (3)
C67	0.62121 (4)	0.0628 (2)	0.54968 (9)	0.0267 (3)
O67	0.62591 (3)	-0.11379 (16)	0.56964 (7)	0.0373 (3)
C61	0.59523 (4)	0.1220 (2)	0.46813 (9)	0.0257 (3)
C62	0.57220 (4)	-0.0213 (2)	0.41643 (10)	0.0304 (3)
O62	0.57228 (3)	-0.21027 (18)	0.44091 (8)	0.0421 (3)
C63	0.54909 (4)	0.0291 (3)	0.33721 (10)	0.0380 (4)
C64	0.54886 (4)	0.2187 (3)	0.30835 (10)	0.0404 (4)
C65	0.57112 (5)	0.3631 (3)	0.35764 (10)	0.0365 (4)
C66	0.59382 (4)	0.3147 (2)	0.43721 (10)	0.0299 (3)
C7	0.63000 (4)	0.3528 (2)	0.64733 (8)	0.0235 (3)
C71	0.59079 (4)	0.3629 (2)	0.63969 (9)	0.0262 (3)
C72	0.57238 (4)	0.1894 (3)	0.64525 (9)	0.0323 (4)
C73	0.53492 (4)	0.1940 (3)	0.62985 (11)	0.0398 (4)
C74	0.51589 (4)	0.3687 (3)	0.60952 (11)	0.0413 (4)
C75	0.53420 (4)	0.5409 (3)	0.60536 (10)	0.0378 (4)
C76	0.57165 (4)	0.5397 (2)	0.62136 (9)	0.0308 (3)
N7A	0.65374 (3)	0.48960 (18)	0.69467 (7)	0.0226 (3)
H22	0.7335	0.9704	0.8415	0.033*
H23	0.7368	1.2661	0.9140	0.035*
H24A	0.6830	1.4424	1.0081	0.057*
H24B	0.6665	1.5633	0.9220	0.057*
H24C	0.7096	1.5342	0.9618	0.057*
H25	0.6288	1.2439	0.8757	0.036*
H26	0.6250	0.9456	0.8050	0.032*
H3	0.7306	0.6916	0.7647	0.029*
H51A	0.7248	0.0911	0.5860	0.046*

H51B	0.6857	0.0512	0.5179	0.046*
H51C	0.6984	-0.0722	0.6040	0.046*
H62	0.5916	-0.2188	0.4937	0.051*
H63	0.5334	-0.0671	0.3029	0.046*
H64	0.5332	0.2518	0.2536	0.049*
H65	0.5707	0.4937	0.3368	0.044*
H66	0.6087	0.4135	0.4716	0.036*
H72	0.5853	0.0688	0.6595	0.039*
H73	0.5223	0.0758	0.6334	0.048*
H74	0.4903	0.3705	0.5984	0.050*
H75	0.5211	0.6611	0.5914	0.045*
H76	0.5842	0.6593	0.6198	0.037*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0206 (6)	0.0306 (7)	0.0204 (6)	-0.0027 (5)	0.0057 (5)	-0.0040 (5)
C2	0.0190 (6)	0.0288 (8)	0.0174 (6)	-0.0027 (5)	0.0025 (5)	0.0012 (5)
C21	0.0218 (7)	0.0275 (7)	0.0181 (6)	-0.0007 (5)	0.0050 (5)	0.0022 (5)
C22	0.0221 (7)	0.0317 (8)	0.0291 (8)	-0.0020 (6)	0.0081 (6)	-0.0024 (6)
C23	0.0271 (7)	0.0288 (8)	0.0292 (8)	-0.0056 (6)	0.0068 (6)	-0.0021 (6)
C24	0.0346 (8)	0.0262 (8)	0.0239 (7)	0.0012 (6)	0.0095 (6)	0.0030 (6)
C241	0.0461 (10)	0.0320 (9)	0.0365 (9)	0.0011 (7)	0.0133 (7)	-0.0048 (7)
C25	0.0274 (8)	0.0332 (8)	0.0294 (8)	0.0064 (6)	0.0098 (6)	0.0026 (6)
C26	0.0207 (7)	0.0321 (8)	0.0258 (7)	-0.0006 (6)	0.0050 (6)	0.0010 (6)
C3	0.0182 (6)	0.0293 (8)	0.0218 (7)	-0.0030 (5)	0.0036 (5)	-0.0003 (6)
C3A	0.0172 (6)	0.0298 (8)	0.0195 (6)	-0.0008 (5)	0.0041 (5)	0.0020 (6)
N4	0.0202 (6)	0.0261 (6)	0.0222 (6)	-0.0001 (5)	0.0043 (5)	-0.0001 (5)
C5	0.0226 (7)	0.0266 (7)	0.0202 (7)	0.0003 (6)	0.0034 (5)	0.0034 (6)
C51	0.0279 (8)	0.0296 (8)	0.0334 (8)	0.0012 (6)	0.0069 (6)	-0.0055 (6)
C6	0.0218 (7)	0.0271 (8)	0.0201 (6)	-0.0032 (6)	0.0025 (5)	0.0013 (6)
C67	0.0248 (7)	0.0278 (8)	0.0275 (7)	-0.0047 (6)	0.0085 (6)	-0.0011 (6)
O67	0.0382 (6)	0.0295 (6)	0.0383 (6)	-0.0067 (5)	0.0040 (5)	0.0040 (5)
C61	0.0204 (7)	0.0315 (8)	0.0248 (7)	-0.0013 (6)	0.0065 (6)	-0.0036 (6)
C62	0.0243 (7)	0.0369 (9)	0.0306 (8)	-0.0080 (6)	0.0098 (6)	-0.0069 (7)
O62	0.0417 (7)	0.0394 (7)	0.0410 (7)	-0.0167 (5)	0.0072 (5)	-0.0067 (5)
C63	0.0282 (8)	0.0547 (11)	0.0288 (8)	-0.0130 (7)	0.0057 (6)	-0.0085 (8)
C64	0.0291 (8)	0.0612 (12)	0.0259 (8)	-0.0008 (8)	0.0017 (6)	0.0009 (8)
C65	0.0345 (9)	0.0411 (10)	0.0313 (8)	0.0041 (7)	0.0065 (7)	0.0043 (7)
C66	0.0275 (7)	0.0333 (8)	0.0272 (8)	-0.0005 (6)	0.0061 (6)	-0.0025 (6)
C7	0.0205 (7)	0.0293 (8)	0.0187 (6)	-0.0046 (5)	0.0035 (5)	0.0009 (6)
C71	0.0206 (7)	0.0372 (8)	0.0193 (7)	-0.0061 (6)	0.0039 (5)	-0.0048 (6)
C72	0.0273 (8)	0.0428 (9)	0.0260 (8)	-0.0091 (7)	0.0072 (6)	-0.0005 (7)
C73	0.0279 (8)	0.0571 (11)	0.0345 (9)	-0.0163 (8)	0.0101 (7)	-0.0012 (8)
C74	0.0194 (7)	0.0691 (13)	0.0337 (9)	-0.0083 (8)	0.0062 (6)	-0.0070 (8)
C75	0.0240 (8)	0.0532 (11)	0.0328 (9)	0.0010 (7)	0.0044 (6)	-0.0077 (8)
C76	0.0222 (7)	0.0414 (9)	0.0266 (7)	-0.0032 (6)	0.0047 (6)	-0.0052 (7)
N7A	0.0178 (6)	0.0292 (6)	0.0196 (6)	-0.0027 (5)	0.0045 (4)	-0.0018 (5)

Geometric parameters (\AA , $\text{^{\circ}}$)

N1—C2	1.3505 (18)	C51—H51C	0.98
C2—C3	1.4013 (19)	C6—C67	1.496 (2)
C3—C3A	1.377 (2)	C67—O67	1.2418 (19)
C3A—N4	1.3525 (18)	C67—C61	1.468 (2)
N4—C5	1.3141 (18)	C61—C66	1.400 (2)
C5—C6	1.4342 (19)	C61—C62	1.413 (2)
C6—C7	1.376 (2)	C62—O62	1.346 (2)
C7—N7A	1.3675 (18)	C62—C63	1.384 (2)
N7A—N1	1.3603 (16)	O62—H62	0.96
C3A—N7A	1.4011 (17)	C63—C64	1.373 (3)
C2—C21	1.472 (2)	C63—H63	0.95
C21—C26	1.397 (2)	C64—C65	1.389 (2)
C21—C22	1.399 (2)	C64—H64	0.95
C22—C23	1.389 (2)	C65—C66	1.379 (2)
C22—H22	0.95	C65—H65	0.95
C23—C24	1.387 (2)	C66—H66	0.95
C23—H23	0.95	C7—C71	1.4857 (19)
C24—C25	1.393 (2)	C71—C76	1.393 (2)
C24—C241	1.505 (2)	C71—C72	1.395 (2)
C241—H24A	0.98	C72—C73	1.392 (2)
C241—H24B	0.98	C72—H72	0.95
C241—H24C	0.98	C73—C74	1.380 (3)
C25—C26	1.388 (2)	C73—H73	0.95
C25—H25	0.95	C74—C75	1.380 (3)
C26—H26	0.95	C74—H74	0.95
C3—H3	0.95	C75—C76	1.390 (2)
C5—C51	1.502 (2)	C75—H75	0.95
C51—H51A	0.98	C76—H76	0.95
C51—H51B	0.98		
C2—N1—N7A	103.70 (11)	O67—C67—C61	120.77 (13)
N1—C2—C3	113.00 (13)	O67—C67—C6	118.40 (13)
N1—C2—C21	120.14 (12)	C61—C67—C6	120.68 (13)
C3—C2—C21	126.83 (12)	C66—C61—C62	118.34 (14)
C26—C21—C22	118.38 (14)	C66—C61—C67	122.14 (13)
C26—C21—C2	121.83 (13)	C62—C61—C67	119.42 (14)
C22—C21—C2	119.77 (13)	O62—C62—C63	117.67 (14)
C23—C22—C21	120.56 (14)	O62—C62—C61	122.05 (14)
C23—C22—H22	119.7	C63—C62—C61	120.28 (15)
C21—C22—H22	119.7	C62—O62—H62	105.2
C24—C23—C22	121.35 (14)	C64—C63—C62	119.82 (15)
C24—C23—H23	119.3	C64—C63—H63	120.1
C22—C23—H23	119.3	C62—C63—H63	120.1
C23—C24—C25	117.82 (14)	C63—C64—C65	121.32 (15)
C23—C24—C241	121.09 (14)	C63—C64—H64	119.3
C25—C24—C241	121.07 (14)	C65—C64—H64	119.3

C24—C241—H24A	109.5	C66—C65—C64	119.18 (16)
C24—C241—H24B	109.5	C66—C65—H65	120.4
H24A—C241—H24B	109.5	C64—C65—H65	120.4
C24—C241—H24C	109.5	C65—C66—C61	121.05 (15)
H24A—C241—H24C	109.5	C65—C66—H66	119.5
H24B—C241—H24C	109.5	C61—C66—H66	119.5
C26—C25—C24	121.68 (14)	N7A—C7—C6	116.24 (12)
C26—C25—H25	119.2	N7A—C7—C71	119.98 (12)
C24—C25—H25	119.2	C6—C7—C71	123.77 (13)
C25—C26—C21	120.20 (14)	C76—C71—C72	119.58 (14)
C25—C26—H26	119.9	C76—C71—C7	121.17 (13)
C21—C26—H26	119.9	C72—C71—C7	119.10 (14)
C3A—C3—C2	105.11 (12)	C73—C72—C71	119.65 (16)
C3A—C3—H3	127.4	C73—C72—H72	120.2
C2—C3—H3	127.4	C71—C72—H72	120.2
N4—C3A—C3	132.04 (13)	C74—C73—C72	120.58 (16)
N4—C3A—N7A	121.80 (12)	C74—C73—H73	119.7
C3—C3A—N7A	106.16 (12)	C72—C73—H73	119.7
C5—N4—C3A	118.00 (12)	C75—C74—C73	119.77 (15)
N4—C5—C6	121.89 (13)	C75—C74—H74	120.1
N4—C5—C51	116.77 (13)	C73—C74—H74	120.1
C6—C5—C51	121.29 (13)	C74—C75—C76	120.53 (17)
C5—C51—H51A	109.5	C74—C75—H75	119.7
C5—C51—H51B	109.5	C76—C75—H75	119.7
H51A—C51—H51B	109.5	C75—C76—C71	119.85 (15)
C5—C51—H51C	109.5	C75—C76—H76	120.1
H51A—C51—H51C	109.5	C71—C76—H76	120.1
H51B—C51—H51C	109.5	N1—N7A—C7	126.51 (11)
C7—C6—C5	120.48 (13)	N1—N7A—C3A	112.01 (11)
C7—C6—C67	121.65 (13)	C7—N7A—C3A	121.43 (12)
C5—C6—C67	117.86 (13)		
N7A—N1—C2—C3	-1.32 (15)	C67—C61—C62—O62	2.6 (2)
N7A—N1—C2—C21	176.72 (12)	C66—C61—C62—C63	0.3 (2)
N1—C2—C21—C26	-2.2 (2)	C67—C61—C62—C63	-176.11 (14)
C3—C2—C21—C26	175.57 (14)	O62—C62—C63—C64	-177.97 (15)
N1—C2—C21—C22	179.38 (13)	C61—C62—C63—C64	0.8 (2)
C3—C2—C21—C22	-2.9 (2)	C62—C63—C64—C65	-0.9 (3)
C26—C21—C22—C23	0.3 (2)	C63—C64—C65—C66	-0.1 (3)
C2—C21—C22—C23	178.82 (13)	C64—C65—C66—C61	1.2 (2)
C21—C22—C23—C24	-0.2 (2)	C62—C61—C66—C65	-1.3 (2)
C22—C23—C24—C25	-0.6 (2)	C67—C61—C66—C65	174.98 (14)
C22—C23—C24—C241	177.57 (14)	C5—C6—C7—N7A	2.6 (2)
C23—C24—C25—C26	1.1 (2)	C67—C6—C7—N7A	-175.93 (12)
C241—C24—C25—C26	-176.99 (14)	C5—C6—C7—C71	-178.45 (13)
C24—C25—C26—C21	-1.0 (2)	C67—C6—C7—C71	3.0 (2)
C22—C21—C26—C25	0.2 (2)	N7A—C7—C71—C76	47.23 (19)
C2—C21—C26—C25	-178.22 (13)	C6—C7—C71—C76	-131.71 (15)

N1—C2—C3—C3A	0.66 (16)	N7A—C7—C71—C72	−137.25 (14)
C21—C2—C3—C3A	−177.23 (13)	C6—C7—C71—C72	43.8 (2)
C2—C3—C3A—N4	−179.55 (14)	C76—C71—C72—C73	2.0 (2)
C2—C3—C3A—N7A	0.29 (15)	C7—C71—C72—C73	−173.62 (14)
C3—C3A—N4—C5	−177.43 (15)	C71—C72—C73—C74	−0.3 (2)
N7A—C3A—N4—C5	2.75 (19)	C72—C73—C74—C75	−0.7 (3)
C3A—N4—C5—C6	0.7 (2)	C73—C74—C75—C76	0.1 (3)
C3A—N4—C5—C51	178.12 (12)	C74—C75—C76—C71	1.6 (2)
N4—C5—C6—C7	−3.5 (2)	C72—C71—C76—C75	−2.6 (2)
C51—C5—C6—C7	179.21 (13)	C7—C71—C76—C75	172.87 (13)
N4—C5—C6—C67	175.04 (13)	C2—N1—N7A—C7	−176.01 (13)
C51—C5—C6—C67	−2.2 (2)	C2—N1—N7A—C3A	1.50 (14)
C7—C6—C67—O67	−116.63 (16)	C6—C7—N7A—N1	178.09 (12)
C5—C6—C67—O67	64.82 (18)	C71—C7—N7A—N1	−0.9 (2)
C7—C6—C67—C61	67.74 (19)	C6—C7—N7A—C3A	0.80 (19)
C5—C6—C67—C61	−110.80 (15)	C71—C7—N7A—C3A	−178.21 (12)
O67—C67—C61—C66	−168.53 (14)	N4—C3A—N7A—N1	178.71 (12)
C6—C67—C61—C66	7.0 (2)	C3—C3A—N7A—N1	−1.15 (15)
O67—C67—C61—C62	7.7 (2)	N4—C3A—N7A—C7	−3.6 (2)
C6—C67—C61—C62	−176.77 (13)	C3—C3A—N7A—C7	176.50 (12)
C66—C61—C62—O62	178.99 (14)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O62—H62···O67	0.96	1.68	2.554 (2)	150
C3—H3···N4 ⁱ	0.95	2.61	3.543 (2)	168
C22—H22···N4 ⁱ	0.95	2.57	3.513 (2)	174
C74—H74···O62 ⁱⁱ	0.95	2.55	3.422 (2)	154
C76—H76···O67 ⁱⁱⁱ	0.95	2.56	3.444 (2)	155

Symmetry codes: (i) $-x+3/2, y+1/2, -z+3/2$; (ii) $-x+1, -y, -z+1$; (iii) $x, y+1, z$.