

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# Bis(2,2'-bipyridyl- $\kappa^2N,N'$ )(carbonato- $\kappa^2O,O'$ )cobalt(III) bromide trihydrate

Peng-Tao Ma, Yu-Xia Wang, Guo-Qian Zhang and Ming-Xue Li\*

Institute of Molecular and Crystal Engineering, College of Chemistry and Chemical Engineering, Henan University, Kaifeng 475004, People's Republic of China  
Correspondence e-mail: mptonline@163.com


Received 20 November 2007; accepted 23 November 2007

Key indicators: single-crystal X-ray study;  $T = 296$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.028;  $wR$  factor = 0.084; data-to-parameter ratio = 13.7.

The title complex,  $[\text{Co}(\text{CO}_3)(\text{C}_{10}\text{H}_8\text{N}_2)_2]\text{Br}\cdot 3\text{H}_2\text{O}$ , is isostructural with the chloride analogue. The six-coordinated octahedral  $[\text{Co}(2,2'\text{-bipy})_2\text{CO}_3]^+$  cation (2,2'-bipy is 2,2'-bipyridyl), bromide ion and water molecules are linked together *via*  $\text{O}-\text{H}\cdots\text{Br}$  and  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bonds, generating a one-dimensional chain.

## Related literature

For related literature, see: Das *et al.* (1993); Thirumurugan & Natarajan (2004).



## Experimental

### Crystal data

$[\text{Co}(\text{CO}_3)(\text{C}_{10}\text{H}_8\text{N}_2)_2]\text{Br}\cdot 3\text{H}_2\text{O}$   $a = 9.1281$  (1) Å  
 $M_r = 565.27$   $b = 9.6652$  (2) Å  
 Triclinic,  $P\bar{1}$   $c = 13.0732$  (2) Å

$\alpha = 92.054$  (1)°  
 $\beta = 102.315$  (1)°  
 $\gamma = 91.448$  (1)°  
 $V = 1125.48$  (3) Å<sup>3</sup>  
 $Z = 2$

Mo  $K\alpha$  radiation  
 $\mu = 2.58$  mm<sup>-1</sup>  
 $T = 296$  (2) K  
 $0.18 \times 0.12 \times 0.09$  mm

### Data collection

Bruker SMART APEX CCD  
 area-detector diffractometer  
 Absorption correction: multi-scan  
 (*SADABS*; Sheldrick, 1996)  
 $T_{\text{min}} = 0.654$ ,  $T_{\text{max}} = 0.801$

14734 measured reflections  
 4411 independent reflections  
 4000 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.020$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.028$   
 $wR(F^2) = 0.084$   
 $S = 1.08$   
 4411 reflections  
 323 parameters  
 9 restraints

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\text{max}} = 0.31$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.56$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O4}-\text{H4B}\cdots\text{O5}^i$	0.841 (10)	1.997 (11)	2.832 (3)	172 (3)
$\text{O4}-\text{H4A}\cdots\text{O5}^{ii}$	0.839 (10)	2.016 (12)	2.847 (3)	171 (4)
$\text{O5}-\text{H5A}\cdots\text{O3}^{iii}$	0.835 (10)	1.907 (12)	2.735 (3)	171 (3)
$\text{O5}-\text{H5B}\cdots\text{Br1}$	0.837 (10)	2.422 (13)	3.247 (2)	169 (3)
$\text{O6}-\text{H6B}\cdots\text{Br1}$	0.840 (10)	2.514 (12)	3.345 (2)	170 (3)
$\text{O6}-\text{H6A}\cdots\text{Br1}^{iv}$	0.841 (10)	2.541 (12)	3.378 (3)	173 (3)

Symmetry codes: (i)  $x - 1, y, z$ ; (ii)  $-x + 1, -y + 1, -z + 1$ ; (iii)  $x + 1, y + 1, z$ ; (iv)  $-x + 1, -y + 1, -z$ .

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINT-Plus* (Bruker, 2001); data reduction: *SAINT-Plus*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *SHELXTL* (Bruker, 2001); software used to prepare material for publication: *SHELXTL*.

This work was supported by the Basic Research Foundation for Natural Science of Henan University (grant No. 05ZDZR002).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: AT2506).

## References

- Bruker (2001). *SMART*, *SAINT-Plus* and *SHELXTL*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Das, K., Sinha, U. C., Chatterjee, C. & Mishnev, A. (1993). *Z. Kristallogr.* **205**, 316–318.  
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.  
 Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.  
 Thirumurugan, A. & Natarajan, S. (2004). *Dalton Trans.* pp. 2923–2928.

**supplementary materials**

*Acta Cryst.* (2008). E64, m14 [ doi:10.1107/S1600536807062551 ]

## Bis(2,2'-bipyridyl- $\kappa^2N,N'$ )(carbonato- $\kappa^2O,O'$ )cobalt(III) bromide trihydrate

P.-T. Ma, Y.-X. Wang, G.-Q. Zhang and M.-X. Li

### Comment

Recently, the design and assembly of metal coordination polymers continues attracting chemist's interests and constitutes an important area of research (Thirumurugan & Natarajan, 2004). During the past decades, lots of such compounds have been reported, which present predictable one-, two, three-dimensional frameworks by covalent bonds or hydrogen bonds interactions. Herein, we report the title compound (I).

The title complex (I),  $[\text{Co}(2,2'\text{-bipy})_2\text{CO}_3]\text{Br}\cdot 3\text{H}_2\text{O}$ , contains a  $[\text{Co}(2,2'\text{-bipy})_2\text{CO}_3]^+$  complex cation, a bromine ion, and three water molecules (Fig.1), which is isostructural with its chloride analogue (Das *et al.*, 1993). In the molecular structure, Co atom resides in a distorted octahedral environment, which is defined by four nitrogen atoms from two 2,2'-bipyridyl ligands, two oxygen donors from the carbonate anion. In addition, the  $[\text{Co}(2,2'\text{-bipy})_2\text{CO}_3]^+$  cation, bromine ion, and water molecules in the complex are linked together *via*  $\text{O}-\text{H}\cdots\text{Br}$  and  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bonds generating a one-dimensional chain (Fig.2, Table 1).

### Experimental

Solid  $\text{CoSO}_4\cdot 7\text{H}_2\text{O}$  (0.5 mmol, 0.141 g),  $\text{KBr}$  (1.0 mmol, 0.119 g) and 2,2'-bipy (1 mmol, 0.156 g) was dissolved in 20 ml of the mixed solvent of ethanol and water in a ratio of 1:4(v/v). Under continuous stirring, 5 ml (1 mol/L) solution of  $\text{Na}_2\text{CO}_3$  was added dropwise until a purple solution resulted. The solution was filtered and left at room temperature. After slow evaporation over a period of a week, block red crystals of (I) were obtained.

### Refinement

The H atoms of the water molecules were located in a difference synthesis and refined with distance restraints  $\text{O}-\text{H} = 0.85$  (1)Å and  $\text{H}\cdots\text{H} = 1.34$  (2) Å. The remaining H atoms were positioned geometrically with  $\text{C}-\text{H} = 0.93$  Å, and were refined as riding with  $U_{\text{iso}}(\text{H})=1.2U_{\text{eq}}(\text{C})$ .

### Figures




Fig. 1. The molecular structure of (I), showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. One-dimensional structure of (I) linked by hydrogen bonds.

## Bis(2,2'-bipyridyl- $\kappa^2N,N'$ )(carbonato- $\kappa^2O,O'$ )cobalt(III) bromide trihydrate

### Crystal data

$[\text{Co}(\text{CO}_3)(\text{C}_{10}\text{H}_8\text{N}_2)_2]\text{Br}\cdot 3\text{H}_2\text{O}$	$Z = 2$
$M_r = 565.27$	$F_{000} = 572$
Triclinic, $P\bar{1}$	$D_x = 1.668 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation
$a = 9.1281 (1) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 9.6652 (2) \text{ \AA}$	Cell parameters from 9295 reflections
$c = 13.0732 (2) \text{ \AA}$	$\theta = 2.6\text{--}26.7^\circ$
$\alpha = 92.054 (1)^\circ$	$\mu = 2.58 \text{ mm}^{-1}$
$\beta = 102.315 (1)^\circ$	$T = 296 (2) \text{ K}$
$\gamma = 91.448 (1)^\circ$	Block, red
$V = 1125.48 (3) \text{ \AA}^3$	$0.18 \times 0.12 \times 0.09 \text{ mm}$

### Data collection

Bruker SMART APEX CCD area-detector diffractometer	4411 independent reflections
Radiation source: fine-focus sealed tube	4000 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.020$
$T = 296(2) \text{ K}$	$\theta_{\text{max}} = 26.0^\circ$
$\varphi$ and $\omega$ scans	$\theta_{\text{min}} = 2.1^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -10 \rightarrow 11$
$T_{\text{min}} = 0.654$ , $T_{\text{max}} = 0.801$	$k = -11 \rightarrow 11$
14734 measured reflections	$l = -16 \rightarrow 16$

### Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.028$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.084$	$w = 1/[\sigma^2(F_o^2) + (0.0499P)^2 + 0.4188P]$
$S = 1.08$	where $P = (F_o^2 + 2F_c^2)/3$
4411 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
323 parameters	$\Delta\rho_{\text{max}} = 0.31 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.56 \text{ e \AA}^{-3}$

9 restraints

Extinction correction: SHELXL,  
 $F_c^* = kF_c [1 + 0.001 \times F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Primary atom site location: structure-invariant direct methods

Extinction coefficient: 0.0160 (12)

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes)

are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and

goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based

on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of

$F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is

not relevant to the choice of reflections for refinement.  $R$ -factors based

on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -

factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	0.26787 (3)	0.01455 (3)	0.244258 (18)	0.02444 (10)
Br1	0.68362 (3)	0.61581 (3)	0.14986 (2)	0.06088 (12)
C1	0.3267 (2)	-0.2651 (2)	0.31331 (18)	0.0361 (5)
H1	0.2315	-0.2896	0.2742	0.043*
C2	0.4099 (3)	-0.3639 (2)	0.37077 (19)	0.0421 (5)
H2	0.3717	-0.4542	0.3702	0.051*
C3	0.5510 (3)	-0.3268 (2)	0.42920 (18)	0.0422 (5)
H3	0.6083	-0.3917	0.4694	0.051*
C4	0.6063 (2)	-0.1933 (2)	0.42761 (17)	0.0369 (5)
H4	0.7016	-0.1672	0.4660	0.044*
C5	0.5184 (2)	-0.0982 (2)	0.36814 (14)	0.0278 (4)
C6	0.5627 (2)	0.0465 (2)	0.35739 (14)	0.0278 (4)
C7	0.7009 (2)	0.1073 (2)	0.40451 (17)	0.0366 (5)
H7	0.7727	0.0569	0.4478	0.044*
C8	0.7303 (3)	0.2439 (3)	0.38614 (19)	0.0436 (5)
H8	0.8224	0.2865	0.4169	0.052*

## supplementary materials

---

C9	0.6221 (3)	0.3167 (2)	0.3219 (2)	0.0430 (5)
H9	0.6406	0.4087	0.3086	0.052*
C10	0.4857 (2)	0.2511 (2)	0.27730 (17)	0.0354 (5)
H10	0.4126	0.3004	0.2342	0.042*
C11	0.3835 (2)	-0.1361 (2)	0.07871 (17)	0.0331 (4)
H11	0.4386	-0.1880	0.1312	0.040*
C12	0.3882 (3)	-0.1675 (2)	-0.02410 (18)	0.0390 (5)
H12	0.4445	-0.2404	-0.0409	0.047*
C13	0.3086 (3)	-0.0894 (3)	-0.10162 (17)	0.0399 (5)
H13	0.3101	-0.1097	-0.1715	0.048*
C14	0.2262 (2)	0.0193 (2)	-0.07523 (15)	0.0341 (5)
H14	0.1740	0.0744	-0.1266	0.041*
C15	0.2232 (2)	0.0443 (2)	0.02917 (15)	0.0263 (4)
C16	0.1368 (2)	0.1543 (2)	0.06733 (15)	0.0265 (4)
C17	0.0429 (2)	0.2405 (2)	0.00477 (17)	0.0338 (4)
H17	0.0303	0.2332	-0.0677	0.041*
C18	-0.0327 (2)	0.3386 (2)	0.0514 (2)	0.0413 (5)
H18	-0.0990	0.3964	0.0106	0.050*
C19	-0.0085 (3)	0.3493 (2)	0.1585 (2)	0.0447 (6)
H19	-0.0560	0.4167	0.1910	0.054*
C20	0.0864 (3)	0.2598 (2)	0.21795 (18)	0.0395 (5)
H20	0.1021	0.2673	0.2906	0.047*
O3	-0.0230 (2)	-0.0747 (2)	0.37222 (16)	0.0604 (5)
C21	0.0755 (2)	-0.0408 (2)	0.32724 (17)	0.0368 (5)
N1	0.45605 (18)	0.11867 (17)	0.29456 (13)	0.0282 (3)
N2	0.37927 (18)	-0.13463 (17)	0.31232 (12)	0.0281 (3)
N3	0.30138 (17)	-0.03247 (17)	0.10536 (12)	0.0265 (3)
N4	0.15670 (18)	0.16221 (17)	0.17294 (13)	0.0292 (4)
O1	0.08496 (15)	-0.08699 (15)	0.23240 (11)	0.0321 (3)
O2	0.18851 (16)	0.04514 (16)	0.36505 (11)	0.0341 (3)
O4	0.1357 (3)	0.4381 (3)	0.43419 (18)	0.0666 (6)
O5	0.9249 (3)	0.6454 (3)	0.37190 (19)	0.0676 (6)
O6	0.3364 (3)	0.4743 (3)	0.1049 (2)	0.0754 (7)
H4A	0.109 (4)	0.409 (3)	0.4873 (19)	0.084 (12)*
H4B	0.075 (3)	0.498 (3)	0.410 (2)	0.061 (9)*
H6A	0.331 (3)	0.445 (4)	0.0429 (13)	0.074 (12)*
H5A	0.939 (3)	0.7312 (12)	0.379 (2)	0.055 (9)*
H6B	0.423 (2)	0.510 (4)	0.125 (2)	0.082 (12)*
H5B	0.856 (3)	0.630 (3)	0.3190 (19)	0.086 (12)*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Co1	0.02552 (15)	0.02518 (16)	0.02032 (14)	0.00563 (10)	-0.00088 (10)	0.00176 (10)
Br1	0.05214 (18)	0.0591 (2)	0.0695 (2)	0.00644 (13)	0.00614 (14)	0.01561 (15)
C1	0.0349 (11)	0.0304 (11)	0.0403 (12)	0.0004 (8)	0.0015 (9)	0.0057 (9)
C2	0.0496 (13)	0.0294 (11)	0.0486 (13)	0.0054 (9)	0.0114 (11)	0.0107 (10)
C3	0.0474 (13)	0.0386 (13)	0.0398 (12)	0.0159 (10)	0.0036 (10)	0.0143 (10)

C4	0.0358 (11)	0.0412 (13)	0.0302 (10)	0.0108 (9)	-0.0025 (8)	0.0049 (9)
C5	0.0287 (9)	0.0325 (11)	0.0212 (9)	0.0064 (8)	0.0022 (7)	0.0016 (8)
C6	0.0295 (9)	0.0318 (11)	0.0210 (9)	0.0051 (8)	0.0027 (7)	0.0001 (8)
C7	0.0312 (10)	0.0439 (13)	0.0311 (10)	0.0020 (9)	-0.0008 (8)	-0.0013 (9)
C8	0.0387 (12)	0.0451 (14)	0.0438 (13)	-0.0090 (10)	0.0043 (10)	-0.0082 (10)
C9	0.0501 (13)	0.0308 (12)	0.0478 (13)	-0.0054 (10)	0.0111 (11)	-0.0022 (10)
C10	0.0403 (11)	0.0289 (11)	0.0361 (11)	0.0032 (9)	0.0054 (9)	0.0035 (9)
C11	0.0323 (10)	0.0320 (11)	0.0349 (11)	0.0084 (8)	0.0064 (8)	0.0028 (8)
C12	0.0407 (12)	0.0376 (12)	0.0410 (12)	0.0079 (9)	0.0142 (10)	-0.0039 (10)
C13	0.0415 (12)	0.0504 (14)	0.0286 (10)	0.0009 (10)	0.0109 (9)	-0.0057 (9)
C14	0.0323 (10)	0.0435 (12)	0.0245 (10)	0.0009 (9)	0.0010 (8)	0.0048 (9)
C15	0.0230 (9)	0.0283 (10)	0.0261 (9)	0.0007 (7)	0.0017 (7)	0.0027 (8)
C16	0.0235 (9)	0.0263 (10)	0.0279 (9)	0.0004 (7)	0.0009 (7)	0.0034 (8)
C17	0.0303 (10)	0.0341 (11)	0.0351 (11)	0.0037 (8)	0.0008 (8)	0.0110 (9)
C18	0.0345 (11)	0.0324 (12)	0.0550 (14)	0.0109 (9)	0.0016 (10)	0.0155 (10)
C19	0.0439 (12)	0.0327 (12)	0.0583 (15)	0.0143 (10)	0.0116 (11)	0.0008 (11)
C20	0.0456 (12)	0.0354 (12)	0.0377 (11)	0.0143 (10)	0.0082 (9)	-0.0017 (9)
O3	0.0540 (11)	0.0702 (13)	0.0655 (12)	-0.0069 (9)	0.0345 (10)	-0.0080 (10)
C21	0.0353 (11)	0.0412 (12)	0.0339 (11)	0.0074 (9)	0.0065 (9)	0.0011 (9)
N1	0.0304 (8)	0.0269 (9)	0.0257 (8)	0.0042 (7)	0.0023 (6)	0.0010 (6)
N2	0.0292 (8)	0.0282 (9)	0.0251 (8)	0.0036 (6)	0.0012 (6)	0.0031 (6)
N3	0.0264 (8)	0.0259 (8)	0.0256 (8)	0.0043 (6)	0.0015 (6)	0.0015 (6)
N4	0.0295 (8)	0.0291 (9)	0.0274 (8)	0.0063 (7)	0.0019 (6)	0.0005 (7)
O1	0.0260 (7)	0.0374 (8)	0.0298 (7)	0.0018 (6)	-0.0003 (5)	-0.0014 (6)
O2	0.0387 (8)	0.0394 (8)	0.0229 (7)	0.0030 (6)	0.0040 (6)	-0.0022 (6)
O4	0.0692 (14)	0.0753 (16)	0.0579 (13)	0.0224 (12)	0.0159 (11)	0.0070 (11)
O5	0.0678 (14)	0.0646 (15)	0.0665 (14)	0.0084 (11)	0.0022 (11)	0.0222 (11)
O6	0.0618 (14)	0.0878 (18)	0.0730 (17)	-0.0079 (12)	0.0045 (11)	0.0232 (14)

*Geometric parameters (Å, °)*

Co1—O1	1.8896 (14)	C11—C12	1.377 (3)
Co1—O2	1.8897 (14)	C11—H11	0.9300
Co1—N2	1.9195 (16)	C12—C13	1.376 (3)
Co1—N4	1.9238 (16)	C12—H12	0.9300
Co1—N1	1.9440 (17)	C13—C14	1.383 (3)
Co1—N3	1.9447 (16)	C13—H13	0.9300
Co1—C21	2.314 (2)	C14—C15	1.384 (3)
C1—N2	1.339 (3)	C14—H14	0.9300
C1—C2	1.378 (3)	C15—N3	1.352 (2)
C1—H1	0.9300	C15—C16	1.474 (3)
C2—C3	1.381 (3)	C16—N4	1.353 (3)
C2—H2	0.9300	C16—C17	1.373 (3)
C3—C4	1.377 (3)	C17—C18	1.385 (3)
C3—H3	0.9300	C17—H17	0.9300
C4—C5	1.384 (3)	C18—C19	1.370 (4)
C4—H4	0.9300	C18—H18	0.9300
C5—N2	1.353 (2)	C19—C20	1.379 (3)
C5—C6	1.467 (3)	C19—H19	0.9300

## supplementary materials

---

C6—N1	1.357 (2)	C20—N4	1.342 (3)
C6—C7	1.387 (3)	C20—H20	0.9300
C7—C8	1.380 (3)	O3—C21	1.219 (3)
C7—H7	0.9300	C21—O2	1.307 (3)
C8—C9	1.378 (4)	C21—O1	1.324 (3)
C8—H8	0.9300	O4—H4A	0.839 (10)
C9—C10	1.384 (3)	O4—H4B	0.841 (10)
C9—H9	0.9300	O5—H5A	0.835 (10)
C10—N1	1.339 (3)	O5—H5B	0.837 (10)
C10—H10	0.9300	O6—H6A	0.841 (10)
C11—N3	1.346 (3)	O6—H6B	0.840 (10)
O1—Co1—O2	69.26 (6)	N3—C11—H11	119.1
O1—Co1—N2	91.31 (7)	C12—C11—H11	119.1
O2—Co1—N2	88.81 (7)	C13—C12—C11	119.1 (2)
O1—Co1—N4	88.69 (7)	C13—C12—H12	120.4
O2—Co1—N4	92.44 (7)	C11—C12—H12	120.4
N2—Co1—N4	178.66 (7)	C12—C13—C14	119.7 (2)
O1—Co1—N1	165.27 (7)	C12—C13—H13	120.2
O2—Co1—N1	96.93 (7)	C14—C13—H13	120.2
N2—Co1—N1	83.06 (7)	C13—C14—C15	118.66 (19)
N4—Co1—N1	97.25 (7)	C13—C14—H14	120.7
O1—Co1—N3	97.79 (6)	C15—C14—H14	120.7
O2—Co1—N3	166.44 (7)	N3—C15—C14	121.61 (18)
N2—Co1—N3	95.69 (7)	N3—C15—C16	114.35 (16)
N4—Co1—N3	82.99 (7)	C14—C15—C16	124.04 (18)
N1—Co1—N3	96.31 (7)	N4—C16—C17	121.78 (18)
O1—Co1—C21	34.90 (7)	N4—C16—C15	113.11 (16)
O2—Co1—C21	34.39 (7)	C17—C16—C15	125.10 (18)
N2—Co1—C21	89.09 (7)	C16—C17—C18	118.9 (2)
N4—Co1—C21	91.68 (7)	C16—C17—H17	120.6
N1—Co1—C21	131.00 (7)	C18—C17—H17	120.6
N3—Co1—C21	132.64 (7)	C19—C18—C17	119.2 (2)
N2—C1—C2	121.8 (2)	C19—C18—H18	120.4
N2—C1—H1	119.1	C17—C18—H18	120.4
C2—C1—H1	119.1	C18—C19—C20	119.7 (2)
C1—C2—C3	118.9 (2)	C18—C19—H19	120.1
C1—C2—H2	120.5	C20—C19—H19	120.1
C3—C2—H2	120.5	N4—C20—C19	121.3 (2)
C4—C3—C2	119.6 (2)	N4—C20—H20	119.4
C4—C3—H3	120.2	C19—C20—H20	119.4
C2—C3—H3	120.2	O3—C21—O2	125.6 (2)
C3—C4—C5	119.1 (2)	O3—C21—O1	124.9 (2)
C3—C4—H4	120.4	O2—C21—O1	109.44 (18)
C5—C4—H4	120.4	O3—C21—Co1	177.39 (19)
N2—C5—C4	121.01 (19)	O2—C21—Co1	54.74 (10)
N2—C5—C6	113.60 (16)	O1—C21—Co1	54.75 (10)
C4—C5—C6	125.38 (18)	C10—N1—C6	119.14 (17)
N1—C6—C7	121.44 (19)	C10—N1—Co1	126.97 (14)
N1—C6—C5	114.04 (17)	C6—N1—Co1	113.81 (14)

C7—C6—C5	124.51 (18)	C1—N2—C5	119.56 (17)
C8—C7—C6	118.9 (2)	C1—N2—Co1	125.30 (14)
C8—C7—H7	120.5	C5—N2—Co1	115.00 (14)
C6—C7—H7	120.5	C11—N3—C15	119.05 (17)
C9—C8—C7	119.6 (2)	C11—N3—Co1	127.02 (14)
C9—C8—H8	120.2	C15—N3—Co1	113.76 (12)
C7—C8—H8	120.2	C20—N4—C16	119.12 (17)
C8—C9—C10	119.1 (2)	C20—N4—Co1	125.47 (14)
C8—C9—H9	120.4	C16—N4—Co1	115.00 (13)
C10—C9—H9	120.4	C21—O1—Co1	90.36 (12)
N1—C10—C9	121.8 (2)	C21—O2—Co1	90.87 (12)
N1—C10—H10	119.1	H4A—O4—H4B	106.4 (16)
C9—C10—H10	119.1	H5A—O5—H5B	106.9 (16)
N3—C11—C12	121.83 (19)	H6A—O6—H6B	105.4 (16)

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
O4—H4B $\cdots$ O5 <sup>i</sup>	0.841 (10)	1.997 (11)	2.832 (3)	172 (3)
O4—H4A $\cdots$ O5 <sup>ii</sup>	0.839 (10)	2.016 (12)	2.847 (3)	171 (4)
O5—H5A $\cdots$ O3 <sup>iii</sup>	0.835 (10)	1.907 (12)	2.735 (3)	171 (3)
O5—H5B $\cdots$ Br1	0.837 (10)	2.422 (13)	3.247 (2)	169 (3)
O6—H6B $\cdots$ Br1	0.840 (10)	2.514 (12)	3.345 (2)	170 (3)
O6—H6A $\cdots$ Br1 <sup>iv</sup>	0.841 (10)	2.541 (12)	3.378 (3)	173 (3)

Symmetry codes: (i)  $x-1, y, z$ ; (ii)  $-x+1, -y+1, -z+1$ ; (iii)  $x+1, y+1, z$ ; (iv)  $-x+1, -y+1, -z$ .

Fig. 1




Fig. 2

