

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## $\mu$ -Oxido-bis{chlorido[tris(2-pyridyl-methyl)amine]manganese(III)} bis(hexafluoridophosphate)

Qingguo Meng,\* Lintong Wang, Yanzhen Liu and Yan Pang

College of Chemistry and Chemical Engineering, Weifang University, Weifang 261061, People's Republic of China

Correspondence e-mail: qgmeng\_weifang@yahoo.cn


Received 11 November 2007; accepted 11 December 2007

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.029;  $wR$  factor = 0.076; data-to-parameter ratio = 14.4.

In the title compound,  $[\text{Mn}_2\text{O}(\text{C}_{18}\text{H}_{18}\text{N}_4)_2](\text{PF}_6)_2$ , the Mn atom is chelated by a tetradentate ligand *via* four N atoms, and further bonded to one chloride ion and one bridging oxide, to give a centrosymmetric cation and distorted octahedral coordination geometry.

### Related literature

For related literature, see: Scapin *et al.* (1997); Okabe *et al.* (2000); Serre *et al.* (2005).



### Experimental

#### Crystal data

$[\text{Mn}_2\text{O}(\text{C}_{18}\text{H}_{18}\text{N}_4)_2](\text{PF}_6)_2$   
 $M_r = 1067.45$   
 Triclinic,  $P\bar{1}$   
 $a = 8.5517$  (12) Å  
 $b = 11.3128$  (18) Å  
 $c = 12.914$  (2) Å  
 $\alpha = 115.51$  (2)°  
 $\beta = 107.44$  (2)°

$\gamma = 91.49$  (2)°  
 $V = 1058.1$  (3) Å<sup>3</sup>  
 $Z = 1$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.89$  mm<sup>-1</sup>  
 $T = 293$  (2) K  
 $0.28 \times 0.22 \times 0.18$  mm

#### Data collection

Bruker APEXII CCD diffractometer  
 Absorption correction: multi-scan (SADABS; Bruker, 2001)  
 $T_{\min} = 0.788$ ,  $T_{\max} = 0.856$

9180 measured reflections  
 4125 independent reflections  
 3801 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.021$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.029$   
 $wR(F^2) = 0.076$   
 $S = 1.00$   
 4125 reflections

287 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.27$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.31$  e Å<sup>-3</sup>

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT-Plus (Bruker, 2001); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXS97 (Sheldrick, 1997); program(s) used to refine structure: SHELXL97 (Sheldrick, 1997); molecular graphics: SHELXTL (Bruker, 2001); software used to prepare material for publication: SHELXTL.

The authors thank Liaocheng University for financial support and Professor Jianmin Dou for his help.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CF2174).

### References

- Bruker (2001). SADABS, SAINT-Plus and SHELXTL. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Bruker (2004). APEX2. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Okabe, N. & Oya, N. (2000). *Acta Cryst.* **C56**, 1416–1417.  
 Scapin, G., Reddy, S. G., Zheng, R. & Blanchard, J. S. (1997). *Biochemistry*, **36**, 15081–15088.  
 Serre, C., Marrot, J. & Ferey, G. (2005). *Inorg. Chem.* **44**, 654–658.  
 Sheldrick, G. M. (1997). SHELXS97 and SHELXL97. University of Göttingen, Germany.

**supplementary materials**

*Acta Cryst.* (2008). E64, m204 [ doi:10.1107/S1600536807066512 ]

**$\mu$ -Oxido-bis{chlorido[tris(2-pyridylmethyl)amine]manganese(III)} bis(hexafluoridophosphate)**

**Q. Meng, L. Wang, Y. Liu and Y. Pang**

**Comment**

In recent years, many symmetrical polypyridine ligands and their coordination complexes have been synthesized (Scapin *et al.*, 1997; Okabe *et al.*, 2000; Serre *et al.*, 2005). In this paper, we report the structure of the title compound, (I), containing an unsymmetrical polypyridine ligand.

As shown in Fig. 1, the Mn atom is chelated by the tetradentate ligand *via* four N atoms, and further bonded to one chloride ion and one bridging oxide, to give a centrosymmetric cation and distorted octahedral coordination geometry.

**Experimental**

A mixture of manganese(III) acetate (1 mmol) and tris(2-pyridylmethyl)amine (1 mmol) in 20 ml methanol was refluxed for two hours. The cooled solution was filtered and the filtrate allowed to evaporate at room temperature. Two days later, pink blocks of (I) were obtained with a yield of 30%. Anal. Calc. for  $C_{36}H_{36}Cl_2F_{12}Mn_2N_8OP_2$ : C 40.48, H 3.37, N 10.50%; Found: C 40.42, H 3.38, N 10.44%.

**Refinement**

All H atoms were placed in calculated positions with C—H = 0.93 Å and refined as riding with  $U_{iso}(H) = 1.2U_{eq}(\text{carrier})$ .

**Figures**




Fig. 1. The molecular structure of (I), drawn with 30% probability displacement ellipsoids for the non-hydrogen atoms.

**$\mu$ -Oxido-bis{chlorido[tris(2-pyridylmethyl)amine]manganese(III)} bis(hexafluoridophosphate)**

*Crystal data*

$[Mn_2O(C_{18}H_{18}Cl_1N_4)_2](PF_6)_2$

$M_r = 1067.45$

Triclinic,  $P\bar{1}$

Hall symbol: -P 1

$Z = 1$

$F_{000} = 538$

$D_x = 1.675 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation

$\lambda = 0.71073 \text{ \AA}$

# supplementary materials

---

$a = 8.5517 (12) \text{ \AA}$	Cell parameters from 4125 reflections
$b = 11.3128 (18) \text{ \AA}$	$\theta = 3.0\text{--}26.0^\circ$
$c = 12.914 (2) \text{ \AA}$	$\mu = 0.89 \text{ mm}^{-1}$
$\alpha = 115.51 (2)^\circ$	$T = 293 (2) \text{ K}$
$\beta = 107.44 (2)^\circ$	Block, pink
$\gamma = 91.49 (2)^\circ$	$0.28 \times 0.22 \times 0.18 \text{ mm}$
$V = 1058.1 (3) \text{ \AA}^3$	

## Data collection

Bruker APEXII CCD diffractometer	4125 independent reflections
Radiation source: fine-focus sealed tube	3801 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.021$
$T = 293(2) \text{ K}$	$\theta_{\text{max}} = 26.0^\circ$
$\varphi$ and $\omega$ scans	$\theta_{\text{min}} = 3.0^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2001)	$h = -8 \rightarrow 10$
$T_{\text{min}} = 0.788, T_{\text{max}} = 0.856$	$k = -13 \rightarrow 13$
9180 measured reflections	$l = -15 \rightarrow 15$

## Refinement

Refinement on $F^2$	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.029$	$w = 1/[\sigma^2(F_o^2) + (0.0355P)^2 + 0.508P]$
$wR(F^2) = 0.076$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.00$	$(\Delta/\sigma)_{\text{max}} = 0.001$
4125 reflections	$\Delta\rho_{\text{max}} = 0.27 \text{ e \AA}^{-3}$
287 parameters	$\Delta\rho_{\text{min}} = -0.31 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: SHELXL97 (Sheldrick, 1997), $F_c^* = kFc[1+0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$
Secondary atom site location: difference Fourier map	Extinction coefficient: 0.0273 (16)

## Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Mn1	1.03744 (3)	0.34680 (2)	0.39271 (2)	0.0591 (3)
C16	0.9137 (3)	0.5294 (2)	0.15470 (19)	0.0579 (6)
H16	0.8144	0.5473	0.1147	0.070*
C15	1.0631 (4)	0.5736 (2)	0.1515 (2)	0.0698 (7)
H15	1.0656	0.6240	0.1109	0.084*
C12	0.7465 (3)	0.1753 (2)	0.14497 (19)	0.0547 (5)
H12A	0.7451	0.2011	0.0822	0.066*
H12B	0.6370	0.1255	0.1212	0.066*
C14	1.2072 (4)	0.5437 (3)	0.2077 (2)	0.0693 (7)
H14	1.3080	0.5723	0.2048	0.083*
C13	1.2008 (3)	0.4705 (2)	0.2687 (2)	0.0586 (5)
H13	1.2989	0.4497	0.3070	0.070*
C9	0.9549 (3)	-0.1237 (2)	0.0608 (2)	0.0591 (6)
H9	0.9333	-0.2116	0.0007	0.071*
C7	1.1344 (3)	0.0604 (2)	0.2403 (2)	0.0506 (5)
H7	1.2376	0.0959	0.3026	0.061*
C10	0.8352 (3)	-0.0448 (2)	0.05877 (19)	0.0503 (5)
H10	0.7312	-0.0789	-0.0027	0.060*
C8	1.1067 (3)	-0.0700 (2)	0.1531 (2)	0.0589 (6)
H8	1.1897	-0.1211	0.1567	0.071*
C6	0.6719 (2)	0.2876 (2)	0.33136 (19)	0.0473 (5)
H6A	0.5632	0.2359	0.2740	0.057*
H6B	0.6564	0.3760	0.3821	0.057*
C3	0.7291 (3)	0.1172 (2)	0.5339 (2)	0.0583 (5)
H3	0.6660	0.0777	0.5619	0.070*
C1	0.9874 (2)	0.18989 (19)	0.53077 (17)	0.0432 (4)
H1	1.1028	0.1989	0.5575	0.052*
C17	0.9146 (3)	0.45733 (19)	0.21901 (16)	0.0442 (4)
C2	0.8988 (3)	0.1304 (2)	0.57405 (19)	0.0509 (5)
H2	0.9530	0.0998	0.6292	0.061*
C4	0.6514 (3)	0.1631 (2)	0.4512 (2)	0.0528 (5)
H4	0.5359	0.1536	0.4227	0.063*
C18	0.7604 (3)	0.41495 (19)	0.23735 (18)	0.0450 (4)
H18A	0.7428	0.4883	0.3054	0.054*
H18B	0.6641	0.3918	0.1647	0.054*
C5	0.7463 (2)	0.22291 (18)	0.41146 (17)	0.0400 (4)
C11	0.8713 (2)	0.08589 (18)	0.14917 (16)	0.0398 (4)
C11	1.31393 (6)	0.35086 (5)	0.48444 (5)	0.05319 (15)
F3	0.5826 (2)	0.6606 (2)	0.1912 (2)	0.1062 (6)
F5	0.6040 (2)	0.71991 (18)	0.04862 (16)	0.0904 (5)
F2	0.4876 (2)	0.84879 (19)	0.27957 (14)	0.0865 (5)
F1	0.73275 (18)	0.8610 (2)	0.24889 (16)	0.0976 (6)
F6	0.5112 (2)	0.90781 (15)	0.13872 (16)	0.0796 (4)
F4	0.36039 (17)	0.71075 (14)	0.08069 (13)	0.0705 (4)
N3	1.0566 (2)	0.42831 (16)	0.27464 (15)	0.0455 (4)

## supplementary materials

---

N1	0.77789 (18)	0.29832 (15)	0.26229 (14)	0.0395 (3)
N4	1.01895 (19)	0.13923 (15)	0.23967 (14)	0.0415 (3)
N2	0.91299 (18)	0.23567 (15)	0.45101 (14)	0.0386 (3)
O1	1.0000	0.5000	0.5000	0.0399 (4)
P1	0.54877 (7)	0.78428 (6)	0.16552 (5)	0.05074 (15)

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Mn1	0.0639 (6)	0.0610 (6)	0.0547 (6)	0.0008 (4)	0.0176 (5)	-0.0068 (4)
C16	0.0825 (17)	0.0485 (11)	0.0445 (11)	0.0145 (11)	0.0199 (11)	0.0241 (9)
C15	0.109 (2)	0.0528 (13)	0.0546 (13)	0.0012 (13)	0.0329 (14)	0.0282 (11)
C12	0.0490 (12)	0.0473 (11)	0.0438 (11)	0.0091 (9)	-0.0021 (9)	0.0119 (9)
C14	0.0823 (18)	0.0680 (15)	0.0664 (15)	-0.0023 (13)	0.0354 (14)	0.0328 (13)
C13	0.0575 (13)	0.0637 (13)	0.0599 (13)	0.0039 (10)	0.0251 (11)	0.0302 (11)
C9	0.0717 (15)	0.0427 (11)	0.0615 (13)	0.0130 (10)	0.0313 (12)	0.0171 (10)
C7	0.0432 (11)	0.0545 (11)	0.0568 (12)	0.0165 (9)	0.0196 (9)	0.0259 (10)
C10	0.0524 (12)	0.0457 (10)	0.0480 (11)	0.0023 (9)	0.0171 (9)	0.0179 (9)
C8	0.0620 (14)	0.0535 (12)	0.0700 (14)	0.0251 (10)	0.0329 (12)	0.0285 (11)
C6	0.0285 (9)	0.0554 (11)	0.0601 (12)	0.0089 (8)	0.0103 (8)	0.0316 (10)
C3	0.0582 (13)	0.0635 (13)	0.0647 (14)	0.0047 (10)	0.0273 (11)	0.0360 (11)
C1	0.0405 (10)	0.0443 (10)	0.0450 (10)	0.0095 (8)	0.0116 (8)	0.0226 (8)
C17	0.0552 (12)	0.0401 (9)	0.0348 (9)	0.0102 (8)	0.0153 (8)	0.0152 (8)
C2	0.0555 (12)	0.0529 (11)	0.0510 (11)	0.0101 (9)	0.0184 (9)	0.0297 (10)
C4	0.0355 (10)	0.0582 (12)	0.0680 (13)	0.0054 (9)	0.0189 (9)	0.0314 (11)
C18	0.0469 (11)	0.0461 (10)	0.0402 (10)	0.0158 (8)	0.0092 (8)	0.0218 (8)
C5	0.0315 (9)	0.0392 (9)	0.0446 (10)	0.0053 (7)	0.0113 (7)	0.0164 (8)
C11	0.0410 (10)	0.0412 (9)	0.0388 (9)	0.0051 (7)	0.0143 (7)	0.0197 (8)
Cl1	0.0309 (2)	0.0587 (3)	0.0585 (3)	0.0126 (2)	0.0094 (2)	0.0207 (2)
F3	0.0981 (14)	0.1209 (15)	0.1660 (19)	0.0536 (12)	0.0649 (13)	0.1092 (15)
F5	0.0931 (12)	0.0951 (12)	0.0853 (11)	0.0077 (9)	0.0564 (10)	0.0269 (9)
F2	0.0790 (11)	0.1126 (13)	0.0617 (9)	0.0061 (9)	0.0342 (8)	0.0286 (9)
F1	0.0443 (8)	0.1515 (17)	0.0847 (11)	-0.0079 (9)	0.0016 (7)	0.0586 (11)
F6	0.0760 (10)	0.0695 (9)	0.0981 (11)	0.0058 (7)	0.0193 (8)	0.0500 (9)
F4	0.0522 (8)	0.0690 (8)	0.0724 (9)	-0.0055 (6)	0.0123 (7)	0.0238 (7)
N3	0.0483 (9)	0.0479 (9)	0.0439 (9)	0.0075 (7)	0.0187 (7)	0.0225 (7)
N1	0.0332 (8)	0.0421 (8)	0.0402 (8)	0.0094 (6)	0.0078 (6)	0.0193 (7)
N4	0.0391 (8)	0.0433 (8)	0.0436 (8)	0.0111 (6)	0.0153 (7)	0.0205 (7)
N2	0.0325 (8)	0.0400 (8)	0.0429 (8)	0.0076 (6)	0.0113 (6)	0.0197 (7)
O1	0.0350 (9)	0.0409 (9)	0.0411 (9)	0.0076 (7)	0.0104 (7)	0.0185 (8)
P1	0.0402 (3)	0.0644 (3)	0.0516 (3)	0.0071 (2)	0.0144 (2)	0.0312 (3)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Mn1—O1	1.8034 (5)	C6—N1	1.485 (3)
Mn1—N2	2.1227 (16)	C6—C5	1.514 (3)
Mn1—N3	2.1341 (17)	C6—H6A	0.970
Mn1—N1	2.2280 (16)	C6—H6B	0.970
Mn1—N4	2.2893 (17)	C3—C2	1.367 (3)

Mn1—C11	2.2944 (7)	C3—C4	1.386 (3)
C16—C15	1.379 (4)	C3—H3	0.930
C16—C17	1.390 (3)	C1—N2	1.348 (2)
C16—H16	0.930	C1—C2	1.375 (3)
C15—C14	1.361 (4)	C1—H1	0.930
C15—H15	0.930	C17—N3	1.341 (3)
C12—N1	1.491 (2)	C17—C18	1.509 (3)
C12—C11	1.495 (3)	C2—H2	0.930
C12—H12A	0.970	C4—C5	1.378 (3)
C12—H12B	0.970	C4—H4	0.930
C14—C13	1.377 (3)	C18—N1	1.488 (2)
C14—H14	0.930	C18—H18A	0.970
C13—N3	1.346 (3)	C18—H18B	0.970
C13—H13	0.930	C5—N2	1.343 (2)
C9—C8	1.372 (3)	C11—N4	1.339 (2)
C9—C10	1.377 (3)	F3—P1	1.5854 (18)
C9—H9	0.930	F5—P1	1.5906 (16)
C7—N4	1.348 (3)	F2—P1	1.5908 (16)
C7—C8	1.375 (3)	F1—P1	1.5799 (16)
C7—H7	0.930	F6—P1	1.5966 (16)
C10—C11	1.385 (3)	F4—P1	1.6046 (15)
C10—H10	0.930	O1—Mn1 <sup>i</sup>	1.8034 (5)
C8—H8	0.930		
O1—Mn1—N2	90.82 (4)	N2—C1—H1	118.8
O1—Mn1—N3	92.87 (5)	C2—C1—H1	118.8
N2—Mn1—N3	154.78 (6)	N3—C17—C16	121.2 (2)
O1—Mn1—N1	91.90 (5)	N3—C17—C18	116.09 (17)
N2—Mn1—N1	78.71 (6)	C16—C17—C18	122.6 (2)
N3—Mn1—N1	76.24 (6)	C1—C2—C3	118.3 (2)
O1—Mn1—N4	166.67 (4)	C1—C2—H2	120.8
N2—Mn1—N4	82.20 (6)	C3—C2—H2	120.8
N3—Mn1—N4	88.79 (6)	C5—C4—C3	119.6 (2)
N1—Mn1—N4	75.64 (6)	C5—C4—H4	120.2
O1—Mn1—C11	102.88 (3)	C3—C4—H4	120.2
N2—Mn1—C11	103.45 (5)	N1—C18—C17	110.49 (15)
N3—Mn1—C11	100.01 (5)	N1—C18—H18A	109.6
N1—Mn1—C11	164.97 (4)	C17—C18—H18A	109.6
N4—Mn1—C11	89.83 (5)	N1—C18—H18B	109.6
C15—C16—C17	118.5 (2)	C17—C18—H18B	109.6
C15—C16—H16	120.7	H18A—C18—H18B	108.1
C17—C16—H16	120.7	N2—C5—C4	120.59 (18)
C16—C15—C14	120.3 (2)	N2—C5—C6	116.63 (17)
C16—C15—H15	119.9	C4—C5—C6	122.56 (17)
C14—C15—H15	119.9	N4—C11—C10	122.48 (18)
N1—C12—C11	114.72 (15)	N4—C11—C12	117.29 (16)
N1—C12—H12A	108.6	C10—C11—C12	120.17 (18)
C11—C12—H12A	108.6	C17—N3—C13	119.35 (19)
N1—C12—H12B	108.6	C17—N3—Mn1	114.96 (13)

## supplementary materials

---

C11—C12—H12B	108.6	C13—N3—Mn1	124.77 (15)
H12A—C12—H12B	107.6	C12—N1—C6	112.81 (17)
C13—C14—C15	118.8 (2)	C12—N1—C18	109.20 (15)
C13—C14—H14	120.6	C6—N1—C18	112.67 (15)
C15—C14—H14	120.6	C12—N1—Mn1	113.32 (12)
N3—C13—C14	121.8 (2)	C6—N1—Mn1	104.27 (11)
N3—C13—H13	119.1	C18—N1—Mn1	104.25 (11)
C14—C13—H13	119.1	C7—N4—C11	117.38 (17)
C8—C9—C10	118.7 (2)	C7—N4—Mn1	125.96 (14)
C8—C9—H9	120.7	C11—N4—Mn1	116.09 (12)
C10—C9—H9	120.7	C5—N2—C1	119.34 (17)
N4—C7—C8	123.0 (2)	C5—N2—Mn1	114.73 (12)
N4—C7—H7	118.5	C1—N2—Mn1	125.63 (13)
C8—C7—H7	118.5	Mn1 <sup>i</sup> —O1—Mn1	180
C11—C10—C9	119.3 (2)	F3—P1—F1	91.72 (12)
C11—C10—H10	120.3	F3—P1—F2	90.05 (11)
C9—C10—H10	120.3	F1—P1—F2	90.83 (10)
C9—C8—C7	119.1 (2)	F3—P1—F6	179.00 (10)
C9—C8—H8	120.4	F1—P1—F6	89.26 (10)
C7—C8—H8	120.4	F2—P1—F6	89.71 (10)
N1—C6—C5	112.64 (15)	F3—P1—F5	90.98 (11)
N1—C6—H6A	109.1	F1—P1—F5	90.78 (10)
C5—C6—H6A	109.1	F2—P1—F5	178.06 (10)
N1—C6—H6B	109.1	F6—P1—F5	89.23 (10)
C5—C6—H6B	109.1	F3—P1—F4	90.10 (11)
H6A—C6—H6B	107.8	F1—P1—F4	178.14 (11)
C2—C3—C4	119.6 (2)	F2—P1—F4	88.82 (9)
C2—C3—H3	120.2	F6—P1—F4	88.92 (9)
C4—C3—H3	120.2	F5—P1—F4	89.54 (9)
N2—C1—C2	122.44 (19)		

Symmetry codes: (i)  $-x+2, -y+1, -z+1$ .

Fig. 1

