

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# Poly[[ $(\mu_2$ -2-amino-4,5-dimethylbenzene-sulfonato- $\kappa^2$ N:O)( $\mu_2$ -2-methylpyrazine- $\kappa^2$ N:N')silver(I)] monohydrate]

Xian-Wu Dong\* and Yu-Jie Li

Jilin Agricultural Science and Technology College, Jilin 132000, People's Republic of China

Correspondence e-mail: hljwuhua@163.com


Received 23 November 2007; accepted 27 November 2007

Key indicators: single-crystal X-ray study;  $T = 292$  K; mean  $\sigma(\text{C}-\text{C}) = 0.007$  Å; H-atom completeness 89%;  $R$  factor = 0.032;  $wR$  factor = 0.084; data-to-parameter ratio = 17.7.

In the title compound,  $[\text{Ag}(\text{C}_8\text{H}_{10}\text{NO}_3\text{S})(\text{C}_7\text{H}_6\text{N}_2)] \cdot \text{H}_2\text{O}$ , each  $\text{Ag}^{\text{I}}$  cation is four-coordinated by three N atoms from two different 2-methylpyrazine ligands and one  $-\text{NH}_2$  group of a 2-amino-4,5-dimethylbenzenesulfonate ligand, and by one sulfonate O atom, in a distorted tetrahedral coordination geometry. The  $\text{Ag}^{\text{I}}$  centres are bridged by both types of ligands, forming a two-dimensional network.  $\text{N}-\text{H} \cdots \text{O}$  hydrogen bonds and  $\text{O} \cdots \text{O}$  interactions complete the structure.

## Related literature

For related literature, see: Cote & Shimizu (2004); Li *et al.* (2005); Liu *et al.* (2007).



## Experimental

## Crystal data

$[\text{Ag}(\text{C}_8\text{H}_{10}\text{NO}_3\text{S})(\text{C}_7\text{H}_6\text{N}_2)] \cdot \text{H}_2\text{O}$   
 $M_r = 420.23$   
 Orthorhombic,  $P2_12_12_1$   
 $a = 7.2340$  (4) Å  
 $b = 11.7610$  (5) Å  
 $c = 18.913$  (1) Å

$V = 1609.10$  (14) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 1.40$  mm<sup>-1</sup>  
 $T = 292$  (2) K  
 $0.35 \times 0.29 \times 0.25$  mm

## Data collection

Rigaku R-AXIS RAPID diffractometer

Absorption correction: multi-scan (ABSCOR; Higashi, 1995)  
 $T_{\text{min}} = 0.615$ ,  $T_{\text{max}} = 0.711$

13881 measured reflections  
 3667 independent reflections

3083 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.053$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$   
 $wR(F^2) = 0.085$   
 $S = 1.02$   
 3667 reflections  
 207 parameters  
 4 restraints

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\text{max}} = 0.50$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.58$  e Å<sup>-3</sup>  
 Absolute structure: Flack (1983), 1369 Friedel pairs  
 Flack parameter: 0.00 (3)

Table 1

Selected geometric parameters (Å, °).

Ag1—N2	2.243 (3)	Ag1—N3 <sup>i</sup>	2.469 (4)
Ag1—N1	2.301 (4)	Ag1—O3 <sup>ii</sup>	2.525 (3)
N2—Ag1—N1	141.78 (13)	N2—Ag1—O3 <sup>ii</sup>	125.98 (12)
N2—Ag1—N3 <sup>i</sup>	102.19 (12)	N1—Ag1—O3 <sup>ii</sup>	87.53 (14)
N1—Ag1—N3 <sup>i</sup>	98.44 (13)	N3 <sup>i</sup> —Ag1—O3 <sup>ii</sup>	84.57 (12)

Symmetry codes: (i)  $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$ ; (ii)  $x + 1, y, z$ .

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
N1—H2N $\cdots$ O2	0.82 (3)	2.36 (4)	2.982 (5)	133 (4)
N1—H1N $\cdots$ O1W <sup>ii</sup>	0.82 (5)	2.15 (5)	2.946 (5)	164 (5)

Symmetry code: (ii)  $x + 1, y, z$ .

Data collection: *PROCESS-AUTO* (Rigaku, 1998); cell refinement: *PROCESS-AUTO*; data reduction: *PROCESS-AUTO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *SHELXTL-Plus* (Sheldrick, 1990); software used to prepare material for publication: *SHELXL97*.

The authors thank the Science Foundation for Young Teachers of Northeast Normal University (grant No. 20070314) for support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2532).

## References

- Cote, A. P. & Shimizu, G. K. H. (2004). *Inorg. Chem.* **43**, 6663–6673.  
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.  
 Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.  
 Li, F.-F., Ma, J.-F., Song, S.-Y., Yang, J., Liu, Y.-Y. & Su, Z.-M. (2005). *Inorg. Chem.* **44**, 9374–9383.  
 Liu, H.-Y., Ma, J.-C. & Yang, J. (2007). *Acta Cryst.* **E63**, m2707.  
 Rigaku (1998). *PROCESS-AUTO*. Rigaku Corporation, Tokyo, Japan.  
 Sheldrick, G. M. (1990). *SHELXTL-Plus*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.  
 Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.

**supplementary materials**

*Acta Cryst.* (2008). E64, m80 [ doi:10.1107/S1600536807063672 ]

## Poly[[ $(\mu_2$ -2-amino-4,5-dimethylbenzenesulfonato- $\kappa^2N:O$ )( $\mu_2$ -2-methylpyrazine- $\kappa^2N:N'$ )silver(I)] monohydrate]

X.-W. Dong and Y.-J. Li

### Comment

Silver(I) sulfonate coordination polymers have received much attention for their interesting structural features and potential applications (Cote & Shimizu, 2004). Recently, silver(I) sulfonate compounds with nitrogen-based secondary ligands have been reported (Li *et al.*, 2005). We report here the crystal structure of the title compound.

Selected geometric parameters are listed in Table 1. The Ag<sup>I</sup> cation is four-coordinated by three N atoms from two different 2-methylpyrazine ligands and one –NH<sub>2</sub> group of anion, and one sulfonate O atom in a distorted tetrahedral coordination geometry (Fig.1). The Ag—N distances in the title compound are similar to those in related compounds (Liu *et al.*, 2007). The Ag<sup>I</sup> centers are doubly bridged by both types of ligands to form a two-dimensional network (Fig.2), which are linked via N—H···O hydrogen bonds (Table 2) and O···O interactions into a three-dimensional framework (Fig.3).

### Experimental

An aqueous solution (10 ml) of 2-amino-4,5-dimethylbenzenesulfonic acid (1 mmol) was added to solid Ag<sub>2</sub>CO<sub>3</sub> (0.5 mmol) and stirred for several minutes until no further CO<sub>2</sub> was given off. 2-Methylpyrazine (1 mmol) was then added and a precipitate was formed. The precipitate was dissolved by ammonium hydroxide. Crystals of the title compound were obtained by slow evaporation of the solution at room temperature for 7 d.

### Refinement

All H atoms on C atoms were positioned geometrically (C—H = 0.93–0.96 Å) and refined as riding, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  or  $1.5U_{\text{eq}}(\text{C}_{\text{methyl}})$ . The amino H-atoms were located in a difference Fourier map and its positional parameters were refined, with the N—H distances restrained to 0.82 (2) Å, and with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$ . H atoms bonded to water molecules could not be located and were therefore omitted.

### Figures




Fig. 1. The coordination environment of atom Ag1 in the title compound, showing 30% probability displacement ellipsoids [Symmetry codes: (i)  $-x + 1, y + 1/2, -z + 1/2$ ; (ii)  $x + 1, y, z$ ].




Fig. 2. View of a two-dimensional network in the title compound.




Fig. 3. Part of the three-dimensional network of the title compound. Hydrogen bonds are shown as dashed lines. H atoms not involved in hydrogen-bonding interactions have been omitted.

**Poly[[ $(\mu_2$ -2-amino-4,5-dimethylbenzenesulfonato- $\kappa^2$ N:O)(\mu\_2-2-methylpyrazine- $\kappa^2$ N:N')silver(I)] monohydrate]**

*Crystal data*

[Ag(C<sub>8</sub>H<sub>10</sub>NO<sub>3</sub>S)(C<sub>7</sub>H<sub>6</sub>N<sub>2</sub>)]·H<sub>2</sub>O

$M_r$  = 420.23

Orthorhombic,  $P2_12_12_1$

Hall symbol: P 2ac 2ab

$a$  = 7.2340 (4) Å

$b$  = 11.7610 (5) Å

$c$  = 18.913 (1) Å

$V$  = 1609.10 (14) Å<sup>3</sup>

$Z$  = 4

$F_{000}$  = 848

$D_x$  = 1.735 Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation

$\lambda$  = 0.71069 Å

Cell parameters from 3667 reflections

$\theta$  = 2.0–27.5°

$\mu$  = 1.40 mm<sup>-1</sup>

$T$  = 292 (2) K

Block, yellow

0.35 × 0.29 × 0.25 mm

*Data collection*

Rigaku R-Axis RAPID  
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T$  = 292(2) K

$\omega$  scans

Absorption correction: multi-scan  
(ABSCOR; Higashi, 1995)

$T_{\min}$  = 0.615,  $T_{\max}$  = 0.711

13881 measured reflections

3667 independent reflections

3083 reflections with  $I > 2\sigma(I)$

$R_{\text{int}}$  = 0.053

$\theta_{\max}$  = 27.5°

$\theta_{\min}$  = 2.0°

$h$  = -9→9

$k$  = -15→15

$l$  = -24→24

*Refinement*

Refinement on  $F^2$

Least-squares matrix: full

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$	$w = 1/[\sigma^2(F_o^2) + (0.0492P)^2]$
$wR(F^2) = 0.085$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.02$	$(\Delta/\sigma)_{\max} = 0.001$
3667 reflections	$\Delta\rho_{\max} = 0.50 \text{ e } \text{\AA}^{-3}$
207 parameters	$\Delta\rho_{\min} = -0.58 \text{ e } \text{\AA}^{-3}$
4 restraints	Extinction correction: none
Primary atom site location: structure-invariant direct methods	Absolute structure: Flack (1983), with 1369 Friedel pairs
Secondary atom site location: difference Fourier map	Flack parameter: 0.00 (3)

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Ag1	0.57509 (5)	0.32306 (3)	0.368222 (17)	0.04386 (11)
C1	0.1920 (5)	0.5165 (3)	0.4190 (2)	0.0308 (8)
C2	0.1248 (6)	0.6189 (4)	0.3919 (2)	0.0393 (10)
H2	0.0045	0.6211	0.3746	0.047*
C3	0.2286 (7)	0.7160 (4)	0.3898 (2)	0.0444 (11)
C4	0.4079 (8)	0.7131 (4)	0.4174 (2)	0.0455 (11)
C5	0.4746 (6)	0.6123 (4)	0.4454 (2)	0.0408 (10)
H5	0.5927	0.6113	0.4647	0.049*
C6	0.3719 (5)	0.5128 (3)	0.4457 (2)	0.0316 (9)
C7	0.5321 (8)	0.8154 (5)	0.4164 (3)	0.0684 (16)
H7A	0.4744	0.8765	0.4418	0.103*
H7B	0.5535	0.8383	0.3684	0.103*
H7C	0.6478	0.7967	0.4384	0.103*
C8	0.1486 (9)	0.8237 (4)	0.3565 (3)	0.0723 (16)
H8A	0.1767	0.8879	0.3859	0.108*
H8B	0.0169	0.8160	0.3521	0.108*
H8C	0.2018	0.8347	0.3105	0.108*
C9	0.3884 (6)	0.1510 (4)	0.2665 (2)	0.0455 (12)
H10	0.2963	0.2058	0.2704	0.055*
C10	0.3556 (7)	0.0582 (4)	0.2262 (3)	0.0493 (12)
H11	0.2415	0.0505	0.2040	0.059*

## supplementary materials

---

C11	0.6419 (6)	-0.0062 (4)	0.2511 (2)	0.0413 (10)
H12	0.7349	-0.0602	0.2459	0.050*
C12	0.6755 (6)	0.0873 (4)	0.2934 (2)	0.0410 (10)
C13	0.8545 (8)	0.1001 (6)	0.3321 (4)	0.078 (2)
H24A	0.8528	0.1694	0.3590	0.117*
H24B	0.9545	0.1024	0.2988	0.117*
H24C	0.8712	0.0368	0.3635	0.117*
N1	0.4536 (5)	0.4089 (3)	0.46717 (18)	0.0371 (8)
N2	0.5462 (5)	0.1673 (3)	0.30075 (16)	0.0366 (8)
N3	0.4829 (5)	-0.0226 (3)	0.21749 (19)	0.0435 (9)
O1	0.1454 (4)	0.3189 (3)	0.36351 (17)	0.0529 (8)
O2	0.0582 (5)	0.3433 (3)	0.48525 (17)	0.0518 (8)
O3	-0.1281 (4)	0.4287 (3)	0.3932 (2)	0.0573 (9)
O1W	-0.2470 (5)	0.4027 (3)	0.57266 (16)	0.0484 (8)
S1	0.05493 (14)	0.39276 (9)	0.41453 (6)	0.0369 (2)
H1N	0.539 (7)	0.421 (4)	0.495 (2)	0.055*
H2N	0.383 (6)	0.362 (3)	0.485 (2)	0.055*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Ag1	0.04702 (19)	0.03504 (16)	0.04951 (17)	0.00277 (16)	-0.00239 (16)	-0.00726 (15)
C1	0.029 (2)	0.030 (2)	0.0339 (19)	-0.0029 (17)	0.0002 (16)	0.0016 (16)
C2	0.036 (2)	0.036 (2)	0.046 (2)	0.0036 (18)	-0.0017 (17)	-0.0005 (18)
C3	0.057 (3)	0.030 (2)	0.046 (2)	0.003 (2)	-0.003 (2)	0.0004 (18)
C4	0.054 (3)	0.037 (2)	0.045 (2)	-0.011 (2)	0.001 (2)	-0.0067 (18)
C5	0.036 (2)	0.041 (2)	0.045 (2)	-0.006 (2)	0.0013 (18)	-0.0043 (18)
C6	0.027 (2)	0.034 (2)	0.0334 (19)	0.0037 (17)	-0.0002 (15)	-0.0014 (16)
C7	0.078 (4)	0.050 (3)	0.077 (3)	-0.026 (3)	-0.005 (3)	0.002 (3)
C8	0.087 (4)	0.033 (2)	0.097 (4)	0.002 (3)	-0.014 (3)	0.012 (3)
C9	0.038 (3)	0.049 (3)	0.049 (2)	0.009 (2)	-0.0029 (19)	-0.012 (2)
C10	0.040 (2)	0.058 (3)	0.050 (3)	0.007 (2)	-0.012 (2)	-0.016 (2)
C11	0.038 (2)	0.036 (2)	0.050 (2)	0.0044 (19)	-0.001 (2)	-0.007 (2)
C12	0.039 (2)	0.038 (2)	0.046 (2)	-0.001 (2)	0.000 (2)	-0.001 (2)
C13	0.050 (3)	0.066 (4)	0.118 (5)	0.006 (3)	-0.031 (3)	-0.032 (4)
N1	0.031 (2)	0.0407 (19)	0.0398 (18)	0.0009 (16)	-0.0075 (16)	0.0044 (15)
N2	0.0350 (19)	0.0370 (18)	0.0377 (16)	-0.0004 (19)	0.0021 (14)	-0.0052 (15)
N3	0.047 (2)	0.041 (2)	0.0423 (19)	-0.0042 (19)	-0.0043 (17)	-0.0079 (16)
O1	0.0533 (18)	0.0402 (16)	0.0653 (19)	-0.0123 (17)	0.0106 (17)	-0.0140 (19)
O2	0.0434 (18)	0.049 (2)	0.0626 (18)	-0.0032 (17)	0.0058 (15)	0.0194 (15)
O3	0.0284 (16)	0.048 (2)	0.095 (3)	-0.0068 (14)	-0.0196 (16)	0.0175 (18)
O1W	0.0428 (18)	0.0499 (19)	0.0526 (18)	0.0038 (16)	-0.0054 (15)	0.0033 (15)
S1	0.0255 (5)	0.0328 (5)	0.0524 (6)	-0.0029 (4)	-0.0004 (5)	0.0053 (4)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Ag1—N2	2.243 (3)	C8—H8C	0.96
Ag1—N1	2.301 (4)	C9—N2	1.326 (6)

Ag1—N3 <sup>i</sup>	2.469 (4)	C9—C10	1.353 (7)
Ag1—O3 <sup>ii</sup>	2.525 (3)	C9—H10	0.9300
C1—C2	1.396 (6)	C10—N3	1.333 (6)
C1—C6	1.396 (6)	C10—H11	0.93
C1—S1	1.763 (4)	C11—N3	1.329 (5)
C2—C3	1.368 (6)	C11—C12	1.381 (6)
C2—H2	0.93	C11—H12	0.93
C3—C4	1.399 (8)	C12—N2	1.334 (6)
C3—C8	1.528 (7)	C12—C13	1.495 (7)
C4—C5	1.386 (6)	C13—H24A	0.96
C4—C7	1.501 (7)	C13—H24B	0.96
C5—C6	1.386 (6)	C13—H24C	0.96
C5—H5	0.93	N1—H1N	0.82 (5)
C6—N1	1.417 (5)	N1—H2N	0.82 (3)
C7—H7A	0.96	N3—Ag1 <sup>iii</sup>	2.469 (4)
C7—H7B	0.96	O1—S1	1.454 (3)
C7—H7C	0.96	O2—S1	1.459 (3)
C8—H8A	0.96	O3—S1	1.448 (3)
C8—H8B	0.96	O3—Ag1 <sup>iv</sup>	2.525 (3)
N2—Ag1—N1	141.78 (13)	N2—C9—H10	118.6
N2—Ag1—N3 <sup>i</sup>	102.19 (12)	C10—C9—H10	118.6
N1—Ag1—N3 <sup>i</sup>	98.44 (13)	N3—C10—C9	121.6 (4)
N2—Ag1—O3 <sup>ii</sup>	125.98 (12)	N3—C10—H11	119.2
N1—Ag1—O3 <sup>ii</sup>	87.53 (14)	C9—C10—H11	119.2
N3 <sup>i</sup> —Ag1—O3 <sup>ii</sup>	84.57 (12)	N3—C11—C12	123.1 (4)
C2—C1—C6	119.0 (4)	N3—C11—H12	118.5
C2—C1—S1	119.9 (3)	C12—C11—H12	118.5
C6—C1—S1	121.0 (3)	N2—C12—C11	119.9 (4)
C3—C2—C1	122.7 (4)	N2—C12—C13	119.0 (4)
C3—C2—H2	118.7	C11—C12—C13	121.1 (4)
C1—C2—H2	118.7	C12—C13—H24A	109.5
C2—C3—C4	118.6 (4)	C12—C13—H24B	109.5
C2—C3—C8	119.8 (5)	H24A—C13—H24B	109.5
C4—C3—C8	121.7 (4)	C12—C13—H24C	109.5
C5—C4—C3	119.1 (4)	H24A—C13—H24C	109.5
C5—C4—C7	118.8 (5)	H24B—C13—H24C	109.5
C3—C4—C7	122.1 (4)	C6—N1—Ag1	107.7 (2)
C4—C5—C6	122.5 (4)	C6—N1—H1N	111 (4)
C4—C5—H5	118.7	Ag1—N1—H1N	108 (3)
C6—C5—H5	118.7	C6—N1—H2N	116 (3)
C5—C6—C1	118.1 (4)	Ag1—N1—H2N	107 (3)
C5—C6—N1	120.4 (4)	H1N—N1—H2N	108 (4)
C1—C6—N1	121.3 (4)	C9—N2—C12	116.8 (4)
C4—C7—H7A	109.5	C9—N2—Ag1	118.4 (3)
C4—C7—H7B	109.5	C12—N2—Ag1	124.8 (3)
H7A—C7—H7B	109.5	C11—N3—C10	115.8 (4)
C4—C7—H7C	109.5	C11—N3—Ag1 <sup>iii</sup>	124.6 (3)

## supplementary materials

H7A—C7—H7C	109.5	C10—N3—Ag1 <sup>iii</sup>	119.2 (3)
H7B—C7—H7C	109.5	S1—O3—Ag1 <sup>iv</sup>	133.33 (19)
C3—C8—H8A	109.5	O3—S1—O1	113.6 (2)
C3—C8—H8B	109.5	O3—S1—O2	112.8 (2)
H8A—C8—H8B	109.5	O1—S1—O2	111.3 (2)
C3—C8—H8C	109.5	O3—S1—C1	106.66 (19)
H8A—C8—H8C	109.5	O1—S1—C1	105.76 (18)
H8B—C8—H8C	109.5	O2—S1—C1	106.0 (2)
N2—C9—C10	122.8 (4)		
C6—C1—C2—C3	0.6 (6)	C10—C9—N2—Ag1	-178.1 (4)
S1—C1—C2—C3	-175.8 (4)	C11—C12—N2—C9	0.6 (6)
C1—C2—C3—C4	-1.5 (7)	C13—C12—N2—C9	-178.7 (5)
C1—C2—C3—C8	177.4 (5)	C11—C12—N2—Ag1	179.1 (3)
C2—C3—C4—C5	0.4 (7)	C13—C12—N2—Ag1	-0.2 (6)
C8—C3—C4—C5	-178.4 (5)	N1—Ag1—N2—C9	63.6 (4)
C2—C3—C4—C7	179.3 (5)	N3 <sup>i</sup> —Ag1—N2—C9	-57.6 (3)
C8—C3—C4—C7	0.4 (7)	O3 <sup>ii</sup> —Ag1—N2—C9	-149.7 (3)
C3—C4—C5—C6	1.6 (7)	N1—Ag1—N2—C12	-114.9 (3)
C7—C4—C5—C6	-177.3 (4)	N3 <sup>i</sup> —Ag1—N2—C12	124.0 (3)
C4—C5—C6—C1	-2.5 (6)	O3 <sup>ii</sup> —Ag1—N2—C12	31.8 (4)
C4—C5—C6—N1	172.5 (4)	C12—C11—N3—C10	0.8 (7)
C2—C1—C6—C5	1.4 (6)	C12—C11—N3—Ag1 <sup>iii</sup>	173.4 (3)
S1—C1—C6—C5	177.7 (3)	C9—C10—N3—C11	0.4 (7)
C2—C1—C6—N1	-173.6 (4)	C9—C10—N3—Ag1 <sup>iii</sup>	-172.7 (4)
S1—C1—C6—N1	2.8 (5)	Ag1 <sup>iv</sup> —O3—S1—O1	62.2 (4)
N2—C9—C10—N3	-1.1 (8)	Ag1 <sup>iv</sup> —O3—S1—O2	-65.6 (4)
N3—C11—C12—N2	-1.3 (7)	Ag1 <sup>iv</sup> —O3—S1—C1	178.4 (3)
N3—C11—C12—C13	177.9 (5)	C2—C1—S1—O3	-10.7 (4)
C5—C6—N1—Ag1	-90.6 (4)	C6—C1—S1—O3	172.9 (3)
C1—C6—N1—Ag1	84.2 (4)	C2—C1—S1—O1	110.6 (3)
N2—Ag1—N1—C6	-116.6 (3)	C6—C1—S1—O1	-65.8 (4)
N3 <sup>i</sup> —Ag1—N1—C6	5.7 (3)	C2—C1—S1—O2	-131.2 (3)
O3 <sup>ii</sup> —Ag1—N1—C6	89.8 (3)	C6—C1—S1—O2	52.5 (4)
C10—C9—N2—C12	0.5 (7)		

Symmetry codes: (i)  $-x+1, y+1/2, -z+1/2$ ; (ii)  $x+1, y, z$ ; (iii)  $-x+1, y-1/2, -z+1/2$ ; (iv)  $x-1, y, z$ .

### Hydrogen-bond geometry ( $\text{\AA}, ^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H2N $\cdots$ O2	0.82 (3)	2.36 (4)	2.982 (5)	133 (4)
N1—H1N $\cdots$ O1W <sup>ii</sup>	0.82 (5)	2.15 (5)	2.946 (5)	164 (5)

Symmetry codes: (ii)  $x+1, y, z$ .

Fig. 1




Fig. 2




Fig. 3

