

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

cis-cis-trans-Bis(acetonitrile- κN)-dichloridobis(triphenylphosphine- κP)-ruthenium(II) acetonitrile disolvate

Ahmad M. Al-Far and LeGrande M. Slaughter*

Department of Chemistry, Oklahoma State University, Stillwater, OK 74078, USA

Correspondence e-mail: lms@chem.okstate.edu


Received 30 November 2007; accepted 6 December 2007

Key indicators: single-crystal X-ray study; $T = 170$ K; mean $\sigma(C-C) = 0.005$ Å; R factor = 0.035; wR factor = 0.072; data-to-parameter ratio = 21.9.

The title compound, $[\text{RuCl}_2(\text{C}_2\text{H}_3\text{N})_2(\text{C}_{18}\text{H}_{15}\text{P})_2] \cdot 2\text{C}_2\text{H}_3\text{N}$, was obtained upon stirring an acetonitrile/ethanol solution of $[\text{RuCl}_2(\text{PPh}_3)_3]$. In the crystal structure, each Ru^{II} ion is coordinated by two Cl [Ru—Cl = 2.4308 (7) and 2.4139 (7) Å], two N [Ru—N = 2.016 (2) and 2.003 (2) Å], and two P [Ru—P = 2.3688 (7) and 2.3887 (7) Å] atoms in a distorted octahedral geometry. Packing interactions include typical C—H $\cdots\pi$ contacts involving phenyl groups as well as weak hydrogen bonds between CH_3CN methyl H atoms and Cl or solvent CH_3CN N atoms.

Related literature

For the original synthesis, characterization and reactivity of the title compound and its precursor, see: Gilbert & Wilkinson (1969); Stephenson & Wilkinson (1966); Hallman *et al.* (1970); Caulton (1974).



Experimental

Crystal data

 $[\text{RuCl}_2(\text{C}_2\text{H}_3\text{N})_2(\text{C}_{18}\text{H}_{15}\text{P})_2] \cdot 2\text{C}_2\text{H}_3\text{N}$
 $M_r = 860.73$ Orthorhombic, $P2_12_12_1$ $a = 9.0622$ (9) Å $b = 18.0167$ (18) Å $c = 25.628$ (2) Å $V = 4184.3$ (7) Å³ $Z = 4$ Mo $K\alpha$ radiation $\mu = 0.61$ mm⁻¹ $T = 170$ (2) K $0.40 \times 0.35 \times 0.20$ mm

Data collection

 Bruker SMART APEXII CCD diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 2000)
 $T_{\text{min}} = 0.791$, $T_{\text{max}} = 0.887$

 25910 measured reflections
 10568 independent reflections
 9200 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.042$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.035$ $wR(F^2) = 0.072$ $S = 1.02$

10568 reflections

482 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.38$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.30$ e Å⁻³

Absolute structure: Flack (1983), 4387 Friedel pairs

Flack parameter: -0.02 (2)

Table 1

Hydrogen-bond geometry (Å, °).

Cg1 is the centroid of the C51–C56 phenyl ring.

D—H \cdots A	D—H	H \cdots A	D \cdots A	D—H \cdots A
C4—H4B \cdots Cl1 ⁱ	0.98	2.68	3.560 (3)	149
C101—H101 \cdots Cl1 ⁱⁱ	0.98	2.80	3.698 (4)	153
C2—H2C \cdots Cl2 ⁱⁱⁱ	0.98	2.57	3.544 (3)	175
C101—H102 \cdots Cl2	0.98	2.62	3.554 (4)	158
C2—H2A \cdots N100 ⁱ	0.98	2.60	3.519 (5)	155
C101—H103 \cdots N200	0.98	2.72	3.645 (6)	158
C201—H201 \cdots N200 ^{iv}	0.98	2.66	3.526 (7)	148
C64—H64 \cdots Cg1 ⁱⁱⁱ	0.95	2.96	3.715 (4)	138

Symmetry codes: (i) $-x + 2, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x - 1, y, z$; (iii) $x + 1, y, z$; (iv) $x - \frac{1}{2}, -y + \frac{1}{2}, -z$.

Data collection: APEX2 (Bruker, 2006); cell refinement: SAINT (Bruker, 2006); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2000); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL.

The authors thank Oklahoma State University for financial support and the Oklahoma State Regents for Higher Education for providing funds to purchase the APEXII diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2537).

References

- Bruker (2006). APEX2 (Version 2.0) and SAINT (Version 7.23A). Bruker AXS Inc., Madison, Wisconsin, USA.
- Caulton, K. G. (1974). *J. Am. Chem. Soc.* **96**, 3005–3006.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Gilbert, J. D. & Wilkinson, G. (1969). *J. Chem. Soc. A*, pp. 1749–1753.
- Hallman, P. S., Stephenson, T. A. & Wilkinson, G. (1970). *Inorg. Synth.* **12**, 237–240.
- Sheldrick, G. M. (2000). SADABS (Version 2.10) and SHELXTL (Version 6.14). Bruker AXS Inc., Madison, Wisconsin, USA.
- Stephenson, T. A. & Wilkinson, G. (1966). *J. Inorg. Nucl. Chem.* **28**, 945–956.

supplementary materials

Acta Cryst. (2008). E64, m184 [doi:10.1107/S1600536807065968]

***cis-cis-trans*-Bis(acetonitrile- κ N)dichloridobis(triphenylphosphine- κ P)ruthenium(II) acetonitrile disolvate**

A. M. Al-Far and L. M. Slaughter

Comment

[RuCl₂(PPh₃)₃] has been widely used as a convenient synthon for a variety of Ru^{II} complexes (Stephenson & Wilkinson, 1966; Hallman *et al.*, 1970). It readily loses one phosphine ligand in solution to give solvent adducts or chlorido-bridged Ru^{II} species that are potential catalyst precursors (Caulton, 1974). Gilbert & Wilkinson (1969) previously reported the synthesis of two isomers of [RuCl₂(CH₃CN)₂(PPh₃)₂] having either *cis* or *trans* orientations of the acetonitrile ligands as characterized by infrared spectroscopy. The *cis* isomer was obtained upon refluxing [RuCl₂(PPh₃)₃] in CH₃CN/acetone, whereas the *trans* isomer was formed upon refluxing in CH₃CN/toluene. We found that the *cis* isomer could also be obtained by stirring [RuCl₂(PPh₃)₃] in CH₃CN/ethanol at ambient temperature, confirming the importance of a polar co-solvent in favoring a *cis* geometry.

The crystal structure of the title compound contains one [RuCl₂(CH₃CN)₂(PPh₃)₂] complex and two acetonitriles of crystallization in the asymmetric unit. The Ru^{II} complex displays a *cis* orientation of both the chlorido and CH₃CN ligands and a *trans* orientation of the phosphine ligands (Fig. 1). The Ru—Cl [2.4308 (7), 2.4139 (7) Å], Ru—N [2.016 (2), 2.003 (2) Å], and Ru—P [2.3688 (7), 2.3887 (7) Å] distances are in the expected ranges, and the angles between coordinated atoms are in the range 90.02 (6)—93.83 (2)°. In addition to typical C—H \cdots π packing interactions involving phenyl rings, there are several weak hydrogen bonds between C—H bonds of coordinated or solvate acetonitriles and Cl ligands or solvate acetonitrile N atoms (Fig. 2). The H \cdots acceptor distances range from 2.57—2.80 Å, and the C \cdots acceptor distances range from 3.52—3.70 Å (Table 1).

Although it has been little investigated, [RuCl₂(CH₃CN)₂(PPh₃)₂] is a potentially useful precursor for catalytically active Ru species given the presence of two dissociable ligands in a *cis* arrangement.

Experimental

[RuCl₂(PPh₃)₃] (20 mg) was dissolved in a mixture of degassed absolute ethanol (2 ml) and freshly distilled CH₃CN (3 ml) and stirred for 15 min. During this time, the color of the solution changed from dark brown to yellow. The solvent was removed under vacuum, and the resulting yellow powder was dried for a further 2 h. A 10 mg portion of the solid was dissolved in 0.6 ml of acetonitrile and allowed to stand for 3 d under nitrogen. Large yellow-orange crystals of the title compound formed over this time. The crystals became opaque due to solvent loss within 20 min of removal from acetonitrile unless placed in a cold stream. The sample used in this study was cut from a larger (>1 mm) block, immersed in Paratone N oil in a 0.5 mm nylon loop, and placed in the nitrogen cold stream of an APEXII diffractometer at 170 (2) K for X-ray diffraction analysis.

Refinement

Phenyl H atoms were fixed at C—H distances of 0.95 Å and refined as riding, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. Methyl H atoms were placed with idealized threefold symmetry and fixed C—H distances of 0.98 Å, and they were refined in a riding model with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$. In order to assign the absolute structure, 4387 Friedel pairs (71% of all Friedel pairs) were measured, and Friedel opposites were not merged in the reflection list used for structure solution and refinement. The absolute structure parameter (Flack x) refined to -0.02 (2). For the inverted structure, Flack x refined to 1.02 (2), and increases in $R[F^2 > 2\sigma(F^2)]$ and $wR(F^2)$ of 0.33% and 1.29%, respectively, were observed.

Figures




Fig. 1. ORTEP view of the complex portion of the title compound, with displacement ellipsoids at the 50% probability level. Phenyl hydrogen atoms and acetonitriles of crystallization are omitted for clarity.




Fig. 2. Packing diagram showing a portion of the network of weak hydrogen bonds involving acetonitrile C—H bonds. Symmetry codes: (A) $2 - x, 1/2 + y, 1/2 - z$; (B) $-1 + x, y, z$; (C) $1 + x, y, z$; (D) $-1/2 + x, 1/2 - y, -z$. For solvent symmetry equivalents, N200 becomes N20A, etc.

cis-cis-trans-Bis(acetonitrile- κN)dichloridobis(triphenylphosphine- κP)ruthenium(II) acetonitrile disolvate

Crystal data

$[\text{RuCl}_2(\text{C}_2\text{H}_3\text{N})_2(\text{C}_{18}\text{H}_{15}\text{P})_2] \cdot 2\text{C}_2\text{H}_3\text{N}$

$M_r = 860.73$

Orthorhombic, $P2_12_12_1$

Hall symbol: P 2ac 2ab

$a = 9.0622$ (9) Å

$b = 18.0167$ (18) Å

$c = 25.628$ (2) Å

$V = 4184.3$ (7) Å³

$Z = 4$

$F_{000} = 1768$

$D_x = 1.366$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 7584 reflections

$\theta = 2.6\text{--}29.0^\circ$

$\mu = 0.61$ mm⁻¹

$T = 170$ (2) K

Block, orange

$0.40 \times 0.35 \times 0.20$ mm

Data collection

Bruker SMART APEXII CCD
diffractometer

10568 independent reflections

Radiation source: fine-focus sealed tube

9200 reflections with $I > 2\sigma(I)$

Monochromator: graphite $R_{\text{int}} = 0.042$
 Detector resolution: 0.75 pixels mm^{-1} $\theta_{\text{max}} = 29.0^\circ$
 $T = 170(2)$ K $\theta_{\text{min}} = 2.0^\circ$
 φ and ω scans $h = -12 \rightarrow 11$
 Absorption correction: multi-scan (SADABS; Sheldrick, 2000) $k = -23 \rightarrow 24$
 $T_{\text{min}} = 0.791$, $T_{\text{max}} = 0.887$ $l = -24 \rightarrow 34$
 25910 measured reflections

Refinement

Refinement on F^2 $w = 1/[\sigma^2(F_o^2) + (0.0197P)^2 + 1.2863P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 Least-squares matrix: full $(\Delta/\sigma)_{\text{max}} = 0.002$
 $R[F^2 > 2\sigma(F^2)] = 0.035$ $\Delta\rho_{\text{max}} = 0.38 \text{ e } \text{\AA}^{-3}$
 $wR(F^2) = 0.072$ $\Delta\rho_{\text{min}} = -0.30 \text{ e } \text{\AA}^{-3}$
 $S = 1.02$ Extinction correction: none
 10568 reflections Absolute structure: Flack (1983), 4387 Friedel pairs
 482 parameters Flack parameter: -0.02 (2)
 H-atom parameters constrained

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Ru1	0.98665 (2)	0.525404 (10)	0.202447 (8)	0.01835 (5)
N1	1.1935 (2)	0.56649 (11)	0.20310 (9)	0.0221 (4)
C1	1.3072 (3)	0.59216 (14)	0.20769 (11)	0.0260 (6)
C2	1.4541 (3)	0.62345 (18)	0.21633 (14)	0.0424 (9)
H2A	1.4572	0.6478	0.2505	0.064*
H2B	1.4759	0.6599	0.1890	0.064*
H2C	1.5278	0.5836	0.2152	0.064*
N2	0.9070 (2)	0.62612 (12)	0.21976 (8)	0.0220 (5)
C3	0.8554 (3)	0.68174 (16)	0.22992 (11)	0.0268 (6)
C4	0.7865 (4)	0.75235 (16)	0.24243 (12)	0.0371 (7)
H4A	0.7477	0.7749	0.2105	0.056*
H4B	0.8598	0.7855	0.2581	0.056*

supplementary materials

H4C	0.7055	0.7442	0.2671	0.056*
C11	1.07873 (8)	0.40246 (4)	0.18086 (3)	0.02873 (15)
C12	0.73218 (6)	0.48601 (4)	0.20527 (3)	0.02807 (14)
P1	0.95830 (7)	0.55102 (4)	0.11234 (3)	0.02106 (14)
P2	1.00686 (7)	0.49955 (3)	0.29358 (2)	0.02023 (12)
C11	0.8461 (3)	0.63406 (16)	0.10137 (10)	0.0234 (6)
C12	0.9038 (4)	0.70435 (16)	0.11118 (11)	0.0311 (7)
H12	1.0054	0.7095	0.1195	0.037*
C13	0.8153 (4)	0.76658 (17)	0.10896 (13)	0.0392 (8)
H13	0.8569	0.8141	0.1155	0.047*
C14	0.6677 (4)	0.76067 (19)	0.09743 (13)	0.0409 (8)
H14	0.6073	0.8037	0.0957	0.049*
C15	0.6078 (4)	0.69085 (19)	0.08828 (14)	0.0426 (8)
H15	0.5058	0.6861	0.0805	0.051*
C16	0.6965 (3)	0.62787 (17)	0.09043 (12)	0.0321 (7)
H16	0.6545	0.5803	0.0844	0.039*
C21	0.8672 (3)	0.48398 (16)	0.06914 (10)	0.0235 (6)
C22	0.8493 (3)	0.50217 (17)	0.01617 (11)	0.0304 (7)
H22	0.8862	0.5480	0.0033	0.036*
C23	0.7778 (4)	0.45353 (19)	-0.01746 (12)	0.0376 (7)
H23	0.7652	0.4664	-0.0531	0.045*
C24	0.7249 (4)	0.3863 (2)	0.00102 (13)	0.0407 (8)
H24	0.6764	0.3529	-0.0219	0.049*
C25	0.7430 (3)	0.36848 (18)	0.05247 (12)	0.0367 (7)
H25	0.7059	0.3225	0.0650	0.044*
C26	0.8147 (3)	0.41607 (16)	0.08707 (12)	0.0302 (7)
H26	0.8276	0.4023	0.1225	0.036*
C31	1.1296 (3)	0.56553 (17)	0.07510 (11)	0.0274 (6)
C32	1.2373 (3)	0.51060 (18)	0.07986 (12)	0.0356 (7)
H32	1.2235	0.4713	0.1041	0.043*
C33	1.3649 (4)	0.5126 (2)	0.04960 (13)	0.0429 (8)
H33	1.4369	0.4745	0.0529	0.052*
C34	1.3864 (4)	0.5699 (2)	0.01487 (13)	0.0474 (9)
H34	1.4726	0.5711	-0.0062	0.057*
C35	1.2835 (4)	0.6250 (2)	0.01071 (13)	0.0498 (10)
H35	1.3005	0.6652	-0.0125	0.060*
C36	1.1541 (4)	0.6232 (2)	0.03999 (12)	0.0408 (8)
H36	1.0825	0.6613	0.0360	0.049*
C41	0.9408 (3)	0.57923 (16)	0.33105 (10)	0.0287 (7)
C42	0.7897 (4)	0.5863 (2)	0.34125 (12)	0.0411 (8)
H42	0.7238	0.5470	0.3331	0.049*
C43	0.7367 (5)	0.6519 (2)	0.36353 (14)	0.0612 (13)
H43	0.6346	0.6567	0.3714	0.073*
C44	0.8310 (7)	0.7095 (2)	0.37417 (15)	0.0737 (16)
H44	0.7939	0.7540	0.3892	0.088*
C45	0.9794 (7)	0.7031 (2)	0.36316 (14)	0.0654 (13)
H45	1.0441	0.7432	0.3705	0.078*
C46	1.0343 (4)	0.63864 (18)	0.34158 (12)	0.0433 (9)
H46	1.1365	0.6348	0.3339	0.052*

C51	0.9060 (3)	0.42064 (16)	0.32141 (12)	0.0267 (6)
C52	0.8560 (3)	0.36297 (16)	0.28975 (13)	0.0337 (7)
H52	0.8711	0.3653	0.2531	0.040*
C53	0.7839 (3)	0.30185 (18)	0.31140 (17)	0.0461 (10)
H53	0.7493	0.2630	0.2895	0.055*
C54	0.7629 (4)	0.2979 (2)	0.36451 (18)	0.0509 (11)
H54	0.7147	0.2560	0.3792	0.061*
C55	0.8113 (4)	0.3542 (2)	0.39642 (15)	0.0464 (9)
H55	0.7962	0.3509	0.4330	0.056*
C56	0.8823 (3)	0.41577 (18)	0.37543 (12)	0.0338 (7)
H56	0.9148	0.4546	0.3977	0.041*
C61	1.1917 (3)	0.48226 (16)	0.32086 (11)	0.0266 (6)
C62	1.2245 (3)	0.49623 (18)	0.37348 (12)	0.0370 (7)
H62	1.1517	0.5174	0.3956	0.044*
C63	1.3620 (3)	0.4794 (2)	0.39327 (14)	0.0445 (8)
H63	1.3825	0.4880	0.4291	0.053*
C64	1.4698 (4)	0.45017 (19)	0.36158 (16)	0.0490 (9)
H64	1.5650	0.4398	0.3753	0.059*
C65	1.4394 (3)	0.43590 (18)	0.30962 (15)	0.0429 (9)
H65	1.5135	0.4156	0.2876	0.051*
C66	1.3001 (3)	0.45142 (16)	0.28976 (13)	0.0314 (7)
H66	1.2791	0.4406	0.2542	0.038*
N100	0.6221 (4)	0.2438 (2)	0.17890 (18)	0.0832 (13)
C100	0.5542 (4)	0.2940 (3)	0.16867 (16)	0.0563 (11)
C101	0.4699 (4)	0.3596 (2)	0.15297 (16)	0.0562 (10)
H103	0.4561	0.3593	0.1150	0.084*
H102	0.5237	0.4045	0.1632	0.084*
H101	0.3734	0.3591	0.1702	0.084*
N200	0.3740 (5)	0.3076 (2)	0.01978 (17)	0.0726 (11)
C200	0.2643 (5)	0.3231 (2)	0.00348 (18)	0.0559 (11)
C201	0.1228 (6)	0.3429 (3)	-0.0176 (3)	0.113 (2)
H203	0.1257	0.3389	-0.0558	0.169*
H202	0.0988	0.3940	-0.0077	0.169*
H201	0.0473	0.3092	-0.0038	0.169*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ru1	0.01907 (9)	0.01683 (8)	0.01915 (9)	0.00050 (8)	-0.00009 (8)	0.00030 (9)
N1	0.0268 (11)	0.0175 (10)	0.0219 (11)	0.0018 (8)	0.0020 (11)	0.0027 (11)
C1	0.0260 (13)	0.0215 (13)	0.0304 (15)	0.0033 (10)	0.0048 (12)	0.0006 (14)
C2	0.0214 (15)	0.0347 (16)	0.071 (2)	-0.0042 (12)	0.0039 (14)	-0.0045 (17)
N2	0.0266 (12)	0.0218 (11)	0.0176 (11)	0.0005 (9)	-0.0019 (9)	0.0036 (10)
C3	0.0361 (16)	0.0250 (15)	0.0192 (14)	0.0006 (12)	-0.0029 (12)	0.0004 (13)
C4	0.058 (2)	0.0217 (15)	0.0312 (16)	0.0095 (14)	-0.0018 (14)	-0.0020 (13)
Cl1	0.0313 (4)	0.0206 (3)	0.0342 (4)	0.0044 (3)	0.0042 (3)	-0.0014 (3)
Cl2	0.0233 (3)	0.0293 (3)	0.0316 (3)	-0.0013 (2)	-0.0012 (3)	-0.0008 (3)
P1	0.0222 (3)	0.0215 (3)	0.0195 (3)	-0.0017 (3)	-0.0008 (3)	-0.0003 (3)

supplementary materials

P2	0.0188 (3)	0.0210 (3)	0.0209 (3)	0.0012 (2)	0.0001 (3)	0.0029 (3)
C11	0.0292 (15)	0.0234 (14)	0.0176 (13)	-0.0022 (11)	-0.0036 (11)	0.0013 (12)
C12	0.0438 (18)	0.0250 (15)	0.0245 (15)	-0.0099 (13)	-0.0056 (13)	0.0047 (13)
C13	0.067 (2)	0.0201 (14)	0.0305 (16)	-0.0038 (16)	-0.0063 (15)	0.0059 (14)
C14	0.057 (2)	0.0299 (17)	0.0356 (18)	0.0124 (16)	-0.0032 (15)	0.0021 (16)
C15	0.0358 (19)	0.0411 (19)	0.051 (2)	0.0057 (15)	-0.0049 (15)	-0.0085 (18)
C16	0.0297 (16)	0.0246 (15)	0.0421 (18)	0.0022 (12)	-0.0026 (13)	-0.0073 (14)
C21	0.0220 (13)	0.0255 (15)	0.0229 (13)	0.0007 (11)	0.0003 (10)	-0.0039 (12)
C22	0.0348 (16)	0.0285 (15)	0.0278 (15)	0.0031 (12)	-0.0038 (12)	-0.0016 (13)
C23	0.0430 (18)	0.0436 (19)	0.0262 (15)	0.0044 (15)	-0.0070 (13)	-0.0106 (15)
C24	0.0390 (19)	0.047 (2)	0.0360 (18)	-0.0087 (16)	-0.0020 (15)	-0.0214 (17)
C25	0.0400 (18)	0.0325 (17)	0.0377 (18)	-0.0120 (14)	0.0066 (14)	-0.0113 (15)
C26	0.0353 (17)	0.0288 (16)	0.0265 (15)	-0.0020 (13)	0.0026 (12)	-0.0057 (13)
C31	0.0253 (15)	0.0366 (17)	0.0204 (13)	-0.0082 (12)	0.0006 (11)	0.0014 (13)
C32	0.0330 (16)	0.0366 (18)	0.0371 (17)	-0.0018 (13)	0.0112 (13)	-0.0009 (15)
C33	0.0334 (17)	0.052 (2)	0.0434 (19)	0.0000 (15)	0.0108 (14)	-0.0045 (18)
C34	0.0332 (19)	0.079 (3)	0.0303 (17)	-0.0140 (19)	0.0083 (14)	-0.0010 (19)
C35	0.043 (2)	0.079 (3)	0.0277 (17)	-0.015 (2)	0.0050 (15)	0.0214 (19)
C36	0.0340 (17)	0.060 (2)	0.0289 (16)	-0.0080 (16)	-0.0044 (13)	0.0162 (17)
C41	0.0408 (17)	0.0272 (15)	0.0182 (13)	0.0104 (12)	-0.0003 (11)	0.0039 (12)
C42	0.050 (2)	0.045 (2)	0.0290 (16)	0.0171 (16)	0.0111 (15)	0.0141 (16)
C43	0.085 (3)	0.066 (3)	0.0331 (19)	0.047 (2)	0.029 (2)	0.023 (2)
C44	0.153 (5)	0.043 (2)	0.026 (2)	0.044 (3)	0.003 (3)	-0.0040 (18)
C45	0.124 (4)	0.0335 (18)	0.0388 (19)	0.017 (3)	-0.025 (3)	-0.0080 (16)
C46	0.063 (2)	0.0347 (17)	0.0319 (16)	0.0037 (16)	-0.0126 (16)	-0.0022 (14)
C51	0.0171 (13)	0.0287 (15)	0.0345 (16)	0.0022 (11)	-0.0015 (11)	0.0107 (13)
C52	0.0248 (14)	0.0306 (15)	0.0458 (19)	0.0031 (11)	-0.0007 (14)	0.0083 (16)
C53	0.0288 (17)	0.0316 (17)	0.078 (3)	-0.0054 (13)	-0.0054 (17)	0.0155 (19)
C54	0.0262 (17)	0.043 (2)	0.083 (3)	0.0007 (15)	0.0091 (18)	0.039 (2)
C55	0.0317 (18)	0.056 (2)	0.052 (2)	0.0063 (16)	0.0099 (16)	0.028 (2)
C56	0.0280 (16)	0.0389 (18)	0.0346 (17)	0.0086 (13)	0.0062 (13)	0.0143 (15)
C61	0.0203 (13)	0.0268 (14)	0.0327 (15)	-0.0007 (12)	-0.0045 (10)	0.0073 (14)
C62	0.0336 (17)	0.0422 (18)	0.0351 (17)	-0.0014 (13)	-0.0066 (13)	0.0064 (15)
C63	0.0435 (19)	0.0426 (19)	0.0474 (19)	-0.0066 (17)	-0.0238 (15)	0.0127 (19)
C64	0.0262 (17)	0.0414 (18)	0.079 (3)	-0.0039 (14)	-0.0204 (17)	0.0235 (19)
C65	0.0259 (15)	0.0375 (18)	0.065 (2)	0.0039 (12)	0.0045 (15)	0.0139 (18)
C66	0.0255 (14)	0.0282 (14)	0.0405 (18)	0.0014 (11)	0.0018 (12)	0.0080 (14)
N100	0.064 (2)	0.071 (3)	0.115 (4)	0.008 (2)	-0.011 (2)	0.041 (3)
C100	0.041 (2)	0.072 (3)	0.056 (2)	-0.021 (2)	0.0030 (17)	0.004 (2)
C101	0.039 (2)	0.062 (2)	0.067 (2)	-0.0101 (18)	0.0026 (18)	-0.014 (2)
N200	0.061 (3)	0.065 (2)	0.092 (3)	-0.011 (2)	0.001 (2)	-0.012 (2)
C200	0.047 (3)	0.040 (2)	0.081 (3)	-0.0072 (18)	0.009 (2)	-0.007 (2)
C201	0.079 (4)	0.077 (4)	0.183 (7)	0.013 (3)	-0.035 (4)	-0.006 (4)

Geometric parameters (Å, °)

Ru1—N2	2.003 (2)	C33—H33	0.95
Ru1—N1	2.016 (2)	C34—C35	1.366 (5)
Ru1—P1	2.3688 (7)	C34—H34	0.95

Ru1—P2	2.3887 (7)	C35—C36	1.393 (5)
Ru1—Cl2	2.4139 (7)	C35—H35	0.95
Ru1—Cl1	2.4308 (7)	C36—H36	0.95
N1—C1	1.135 (3)	C41—C46	1.391 (4)
C1—C2	1.463 (4)	C41—C42	1.401 (4)
C2—H2A	0.98	C42—C43	1.398 (5)
C2—H2B	0.98	C42—H42	0.95
C2—H2C	0.98	C43—C44	1.371 (7)
N2—C3	1.136 (3)	C43—H43	0.95
C3—C4	1.453 (4)	C44—C45	1.380 (7)
C4—H4A	0.98	C44—H44	0.95
C4—H4B	0.98	C45—C46	1.378 (5)
C4—H4C	0.98	C45—H45	0.95
P1—C11	1.831 (3)	C46—H46	0.95
P1—C21	1.835 (3)	C51—C52	1.394 (4)
P1—C31	1.841 (3)	C51—C56	1.404 (4)
P2—C41	1.828 (3)	C52—C53	1.396 (4)
P2—C51	1.834 (3)	C52—H52	0.95
P2—C61	1.841 (3)	C53—C54	1.376 (6)
C11—C16	1.389 (4)	C53—H53	0.95
C11—C12	1.393 (4)	C54—C55	1.374 (6)
C12—C13	1.380 (4)	C54—H54	0.95
C12—H12	0.95	C55—C56	1.391 (4)
C13—C14	1.374 (5)	C55—H55	0.95
C13—H13	0.95	C56—H56	0.95
C14—C15	1.390 (5)	C61—C66	1.382 (4)
C14—H14	0.95	C61—C62	1.404 (4)
C15—C16	1.392 (4)	C62—C63	1.379 (4)
C15—H15	0.95	C62—H62	0.95
C16—H16	0.95	C63—C64	1.375 (5)
C21—C26	1.391 (4)	C63—H63	0.95
C21—C22	1.406 (4)	C64—C65	1.384 (5)
C22—C23	1.389 (4)	C64—H64	0.95
C22—H22	0.95	C65—C66	1.390 (4)
C23—C24	1.386 (5)	C65—H65	0.95
C23—H23	0.95	C66—H66	0.95
C24—C25	1.367 (4)	N100—C100	1.125 (5)
C24—H24	0.95	C100—C101	1.464 (6)
C25—C26	1.394 (4)	C101—H103	0.98
C25—H25	0.95	C101—H102	0.98
C26—H26	0.95	C101—H101	0.98
C31—C36	1.392 (4)	N200—C200	1.114 (5)
C31—C32	1.396 (4)	C200—C201	1.437 (7)
C32—C33	1.393 (4)	C201—H203	0.98
C32—H32	0.95	C201—H202	0.98
C33—C34	1.377 (5)	C201—H201	0.98
N2—Ru1—N1	90.03 (9)	C33—C32—H32	119.5
N2—Ru1—P1	90.02 (6)	C31—C32—H32	119.5
N1—Ru1—P1	92.15 (7)	C34—C33—C32	119.7 (3)

supplementary materials

N2—Ru1—P2	89.29 (6)	C34—C33—H33	120.1
N1—Ru1—P2	89.55 (7)	C32—C33—H33	120.1
P1—Ru1—P2	178.17 (2)	C35—C34—C33	120.0 (3)
N2—Ru1—C12	85.16 (7)	C35—C34—H34	120.0
N1—Ru1—C12	175.05 (6)	C33—C34—H34	120.0
P1—Ru1—C12	89.02 (2)	C34—C35—C36	121.0 (3)
P2—Ru1—C12	89.23 (2)	C34—C35—H35	119.5
N2—Ru1—C11	178.93 (7)	C36—C35—H35	119.5
N1—Ru1—C11	90.99 (6)	C31—C36—C35	120.1 (3)
P1—Ru1—C11	89.59 (3)	C31—C36—H36	120.0
P2—Ru1—C11	91.06 (2)	C35—C36—H36	120.0
C12—Ru1—C11	93.83 (2)	C46—C41—C42	119.3 (3)
C1—N1—Ru1	173.9 (2)	C46—C41—P2	120.5 (2)
N1—C1—C2	177.0 (3)	C42—C41—P2	119.3 (3)
C1—C2—H2A	109.5	C43—C42—C41	119.3 (4)
C1—C2—H2B	109.5	C43—C42—H42	120.3
H2A—C2—H2B	109.5	C41—C42—H42	120.3
C1—C2—H2C	109.5	C44—C43—C42	120.5 (4)
H2A—C2—H2C	109.5	C44—C43—H43	119.8
H2B—C2—H2C	109.5	C42—C43—H43	119.8
C3—N2—Ru1	176.7 (2)	C43—C44—C45	120.2 (4)
N2—C3—C4	178.8 (3)	C43—C44—H44	119.9
C3—C4—H4A	109.5	C45—C44—H44	119.9
C3—C4—H4B	109.5	C46—C45—C44	120.3 (4)
H4A—C4—H4B	109.5	C46—C45—H45	119.8
C3—C4—H4C	109.5	C44—C45—H45	119.8
H4A—C4—H4C	109.5	C45—C46—C41	120.4 (4)
H4B—C4—H4C	109.5	C45—C46—H46	119.8
C11—P1—C21	101.26 (12)	C41—C46—H46	119.8
C11—P1—C31	105.82 (14)	C52—C51—C56	118.5 (3)
C21—P1—C31	99.21 (12)	C52—C51—P2	120.9 (2)
C11—P1—Ru1	111.67 (9)	C56—C51—P2	120.6 (2)
C21—P1—Ru1	120.58 (9)	C51—C52—C53	120.6 (3)
C31—P1—Ru1	116.22 (9)	C51—C52—H52	119.7
C41—P2—C51	103.96 (13)	C53—C52—H52	119.7
C41—P2—C61	103.37 (13)	C54—C53—C52	119.9 (4)
C51—P2—C61	100.04 (12)	C54—C53—H53	120.1
C41—P2—Ru1	109.60 (9)	C52—C53—H53	120.1
C51—P2—Ru1	119.54 (10)	C55—C54—C53	120.5 (3)
C61—P2—Ru1	118.29 (9)	C55—C54—H54	119.8
C16—C11—C12	118.4 (3)	C53—C54—H54	119.8
C16—C11—P1	120.5 (2)	C54—C55—C56	120.4 (3)
C12—C11—P1	120.5 (2)	C54—C55—H55	119.8
C13—C12—C11	120.9 (3)	C56—C55—H55	119.8
C13—C12—H12	119.6	C55—C56—C51	120.1 (3)
C11—C12—H12	119.6	C55—C56—H56	119.9
C14—C13—C12	120.8 (3)	C51—C56—H56	119.9
C14—C13—H13	119.6	C66—C61—C62	118.4 (3)
C12—C13—H13	119.6	C66—C61—P2	119.7 (2)

C13—C14—C15	119.1 (3)	C62—C61—P2	121.8 (2)
C13—C14—H14	120.5	C63—C62—C61	120.4 (3)
C15—C14—H14	120.5	C63—C62—H62	119.8
C14—C15—C16	120.4 (3)	C61—C62—H62	119.8
C14—C15—H15	119.8	C64—C63—C62	120.6 (3)
C16—C15—H15	119.8	C64—C63—H63	119.7
C11—C16—C15	120.4 (3)	C62—C63—H63	119.7
C11—C16—H16	119.8	C63—C64—C65	119.9 (3)
C15—C16—H16	119.8	C63—C64—H64	120.1
C26—C21—C22	119.0 (3)	C65—C64—H64	120.1
C26—C21—P1	122.3 (2)	C64—C65—C66	119.7 (3)
C22—C21—P1	118.8 (2)	C64—C65—H65	120.1
C23—C22—C21	120.4 (3)	C66—C65—H65	120.1
C23—C22—H22	119.8	C61—C66—C65	121.0 (3)
C21—C22—H22	119.8	C61—C66—H66	119.5
C24—C23—C22	120.0 (3)	C65—C66—H66	119.5
C24—C23—H23	120.0	N100—C100—C101	177.2 (5)
C22—C23—H23	120.0	C100—C101—H103	109.5
C25—C24—C23	119.6 (3)	C100—C101—H102	109.5
C25—C24—H24	120.2	H103—C101—H102	109.5
C23—C24—H24	120.2	C100—C101—H101	109.5
C24—C25—C26	121.7 (3)	H103—C101—H101	109.5
C24—C25—H25	119.2	H102—C101—H101	109.5
C26—C25—H25	119.2	N200—C200—C201	179.8 (6)
C21—C26—C25	119.4 (3)	C200—C201—H203	109.5
C21—C26—H26	120.3	C200—C201—H202	109.5
C25—C26—H26	120.3	H203—C201—H202	109.5
C36—C31—C32	118.3 (3)	C200—C201—H201	109.5
C36—C31—P1	125.2 (2)	H203—C201—H201	109.5
C32—C31—P1	116.4 (2)	H202—C201—H201	109.5
C33—C32—C31	120.9 (3)		
N2—Ru1—P1—C11	-12.66 (12)	Ru1—P1—C31—C36	-133.4 (2)
N1—Ru1—P1—C11	-102.69 (11)	C11—P1—C31—C32	176.5 (2)
Cl2—Ru1—P1—C11	72.50 (10)	C21—P1—C31—C32	-78.9 (2)
Cl1—Ru1—P1—C11	166.34 (10)	Ru1—P1—C31—C32	52.0 (3)
N2—Ru1—P1—C21	-131.27 (12)	C36—C31—C32—C33	-1.2 (5)
N1—Ru1—P1—C21	138.70 (11)	P1—C31—C32—C33	173.9 (2)
Cl2—Ru1—P1—C21	-46.11 (10)	C31—C32—C33—C34	0.8 (5)
Cl1—Ru1—P1—C21	47.73 (10)	C32—C33—C34—C35	0.8 (5)
N2—Ru1—P1—C31	108.85 (13)	C33—C34—C35—C36	-2.0 (6)
N1—Ru1—P1—C31	18.82 (13)	C32—C31—C36—C35	0.0 (5)
Cl2—Ru1—P1—C31	-165.99 (11)	P1—C31—C36—C35	-174.6 (3)
Cl1—Ru1—P1—C31	-72.16 (11)	C34—C35—C36—C31	1.6 (5)
N2—Ru1—P2—C41	0.84 (12)	C51—P2—C41—C46	147.0 (2)
N1—Ru1—P2—C41	90.87 (12)	C61—P2—C41—C46	42.9 (3)
Cl2—Ru1—P2—C41	-84.33 (11)	Ru1—P2—C41—C46	-84.1 (2)
Cl1—Ru1—P2—C41	-178.15 (11)	C51—P2—C41—C42	-44.1 (3)
N2—Ru1—P2—C51	120.59 (12)	C61—P2—C41—C42	-148.2 (2)
N1—Ru1—P2—C51	-149.38 (12)	Ru1—P2—C41—C42	84.8 (2)

supplementary materials

C12—Ru1—P2—C51	35.42 (10)	C46—C41—C42—C43	-2.3 (4)
C11—Ru1—P2—C51	-58.40 (10)	P2—C41—C42—C43	-171.3 (2)
N2—Ru1—P2—C61	-117.22 (13)	C41—C42—C43—C44	1.6 (5)
N1—Ru1—P2—C61	-27.19 (12)	C42—C43—C44—C45	-0.3 (6)
C12—Ru1—P2—C61	157.61 (11)	C43—C44—C45—C46	-0.3 (6)
C11—Ru1—P2—C61	63.79 (11)	C44—C45—C46—C41	-0.5 (5)
C21—P1—C11—C16	32.4 (3)	C42—C41—C46—C45	1.8 (4)
C31—P1—C11—C16	135.5 (2)	P2—C41—C46—C45	170.7 (2)
Ru1—P1—C11—C16	-97.1 (2)	C41—P2—C51—C52	142.5 (2)
C21—P1—C11—C12	-156.8 (2)	C61—P2—C51—C52	-110.9 (2)
C31—P1—C11—C12	-53.8 (3)	Ru1—P2—C51—C52	19.9 (3)
Ru1—P1—C11—C12	73.6 (2)	C41—P2—C51—C56	-39.8 (3)
C16—C11—C12—C13	-1.4 (4)	C61—P2—C51—C56	66.8 (3)
P1—C11—C12—C13	-172.3 (2)	Ru1—P2—C51—C56	-162.40 (19)
C11—C12—C13—C14	0.5 (5)	C56—C51—C52—C53	0.0 (4)
C12—C13—C14—C15	0.4 (5)	P2—C51—C52—C53	177.7 (2)
C13—C14—C15—C16	-0.4 (5)	C51—C52—C53—C54	-0.6 (5)
C12—C11—C16—C15	1.4 (5)	C52—C53—C54—C55	0.6 (5)
P1—C11—C16—C15	172.3 (3)	C53—C54—C55—C56	-0.1 (5)
C14—C15—C16—C11	-0.5 (5)	C54—C55—C56—C51	-0.5 (5)
C11—P1—C21—C26	-125.0 (2)	C52—C51—C56—C55	0.5 (4)
C31—P1—C21—C26	126.7 (2)	P2—C51—C56—C55	-177.2 (2)
Ru1—P1—C21—C26	-1.3 (3)	C41—P2—C61—C66	-150.7 (2)
C11—P1—C21—C22	54.7 (2)	C51—P2—C61—C66	102.2 (2)
C31—P1—C21—C22	-53.6 (2)	Ru1—P2—C61—C66	-29.4 (3)
Ru1—P1—C21—C22	178.40 (18)	C41—P2—C61—C62	32.4 (3)
C26—C21—C22—C23	1.2 (4)	C51—P2—C61—C62	-74.7 (3)
P1—C21—C22—C23	-178.5 (2)	Ru1—P2—C61—C62	153.7 (2)
C21—C22—C23—C24	-0.6 (5)	C66—C61—C62—C63	0.0 (5)
C22—C23—C24—C25	0.3 (5)	P2—C61—C62—C63	176.9 (2)
C23—C24—C25—C26	-0.5 (5)	C61—C62—C63—C64	1.4 (5)
C22—C21—C26—C25	-1.3 (4)	C62—C63—C64—C65	-1.5 (5)
P1—C21—C26—C25	178.3 (2)	C63—C64—C65—C66	0.2 (5)
C24—C25—C26—C21	1.0 (5)	C62—C61—C66—C65	-1.3 (4)
C11—P1—C31—C36	-8.8 (3)	P2—C61—C66—C65	-178.3 (2)
C21—P1—C31—C36	95.7 (3)	C64—C65—C66—C61	1.2 (5)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C4—H4B \cdots C11 ⁱ	0.98	2.68	3.560 (3)	149
C101—H101 \cdots C11 ⁱⁱ	0.98	2.80	3.698 (4)	153
C2—H2C \cdots C12 ⁱⁱⁱ	0.98	2.57	3.544 (3)	175
C101—H102 \cdots C12	0.98	2.62	3.554 (4)	158
C2—H2A \cdots N100 ⁱ	0.98	2.60	3.519 (5)	155
C101—H103 \cdots N200	0.98	2.72	3.645 (6)	158
C201—H201 \cdots N200 ^{iv}	0.98	2.66	3.526 (7)	148
C64—H64 \cdots Cg1 ⁱⁱⁱ	0.95	2.96	3.715 (4)	138

Symmetry codes: (i) $-x+2, y+1/2, -z+1/2$; (ii) $x-1, y, z$; (iii) $x+1, y, z$; (iv) $x-1/2, -y+1/2, -z$.

Fig. 1

