

Retraction of articles

This article reports the retraction of articles published in *Acta Crystallographica Section E* between 2005 and 2009.

After further thorough investigation (see Harrison *et al.*, 2010), articles are retracted as a result of problems with the data sets or incorrect atom assignments. Full details of all the articles are given in Table 1.

Table 1

Details of articles to be retracted, in order of publication.

Title	Reference	DOI	Refcode
<i>Poly[diacquadi-μ₃-malonato-μ-pyrazine-dinickel(II)] catena-Poly[[[diaqua(6-carboxypyridine-2-carboxylato)samarium(II)]-μ-pyridine-2,6-dicarboxylato] tetrahydrate]</i>	Liu <i>et al.</i> (2005) Liu <i>et al.</i> (2006)	10.1107/S1600536805026358 10.1107/S1600536806038141	GATWAA FONCUH03
<i>Poly[[[μ₄-4,4'-carbonylbis(benzene-3,4-dicarboxylato)]tetrakis(1,10-phenanthroline)-dipalladium(II)] dihydrate]</i>	Li, Wang, Zhang & Yu (2007e)	10.1107/S1600536807039050	AFELAZ
<i>Poly[diacqua-μ₃-malonato-μ-pyrazine-diiron(II)]</i>	Li, Liu <i>et al.</i> (2007)	10.1107/S1600536807038743	AFELON
<i>Poly[diacqua-di-μ₃-malonato-μ-pyrazine-dimanganese(II)]</i>	Li, Wang, Zhang & Yu (2007f)	10.1107/S1600536807039773	VIJZAO
<i>Poly[[aqua(2,2-bipyridine)(μ₃-pyridine-3,4-dicarboxylato)cobalt(II)] monohydrate]</i>	Li, Wang, Zhang & Yu (2007g)	10.1107/S1600536807040275	VIKIC
<i>catena-Poly[[[diaqua(6-carboxypyridine-2-carboxylato)holmium(III)]-μ-pyridine-2,6-dicarboxylato] tetrahydrate]</i>	Li, Wang, Zhang & Yu (2007a)	10.1107/S1600536807041657	DILGEL
<i>catena-Poly[[[2,2'-bipyridine-κ²N,N']iron(II)]-μ-5-carboxy-4-carboxylatoimidazol-1-ido-κ⁴N³,O⁴:N¹,O²]</i>	Li, Wang, Zhang & Yu (2007h)	10.1107/S1600536807042122	XIKWAO
<i>Poly[[aqua(2,2'-bipyridine)(μ₃-pyridine-3,4-dicarboxylato)nickel(II)] monohydrate]</i>	Li, Wang, Zhang & Yu (2007b)	10.1107/S1600536807046466	LEVZAO01
<i>2-(Benzyliminomethyl)-6-methoxyphenol</i>	Li, Wang, Zhang & Yu (2007i)	10.1107/S1600536807042134	SILDEX
<i>Poly[aqua(2,2'-bipyridine)(μ₃-pyridine-2,4-dicarboxylato)palladium(II)]</i>	Li, Wang, Zhang & Yu (2007c)	10.1107/S1600536807047575	SILXAN
<i>μ-Oxido-bis[chlorido[tris(2-pyridylmethyl)amine]iron(III)] bis(hexafluoridophosphate)</i>	Liu, Dou, Li & Zhang (2007)	10.1107/S1600536807049665	TINRIS
<i>μ-Oxido-bis[(4,4'-dibromo-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato)-manganese(III)]</i>	Liu, Dou, Niu & Zhang (2007a)	10.1107/S1600536807051008	GIMZAE
<i>Bis[N-(8-quinolyl)pyridine-2-carboxamidato]iron(III) perchlorate monohydrate</i>	Li, Wang, Zhang & Yu (2007d)	10.1107/S1600536807048556	WIMZIC
<i>μ-Oxido-bis[(4,4'-dibromo-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato)-chromium(III)]</i>	Liu, Dou, Niu & Zhang (2007b)	10.1107/S1600536807057996	HIQFIX
<i>μ-Oxido-bis[chlorido[tris(2-pyridylmethyl)amine]chromium(III)] bis(hexafluoridophosphate)</i>	Li, Wang <i>et al.</i> (2008)	10.1107/S1600536807061296	MIRNAD
<i>μ-Oxido-bis[(4,4'-dibromo-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato)-iron(III)]</i>	Meng <i>et al.</i> (2008a)	10.1107/S1600536807063143	MIRWUG
<i>catena-Poly[[bis(1H-benzimidazole-κ³N,N')palladium(II)]-μ-benzene-1,4-dicarboxylato-κ²O¹:O²]</i>	Meng <i>et al.</i> (2008b)	10.1107/S1600536807065051	XISCAE
<i>Oxalato-bis(propene-1,3-diamine)manganese(II) chloride monohydrate</i>	Meng <i>et al.</i> (2008e)	10.1107/S1600536807065361	SISWIB
<i>μ-Oxido-bis[chlorido[tris(2-pyridylmethyl)amine]manganese(III)] bis(hexafluoridophosphate)</i>	Meng <i>et al.</i> (2008c)	10.1107/S1600536807066512	RISRIV
<i>Bis[N-(8-quinolyl)pyridine-2-carboxamidato-κ³N,N,N']manganese(III) perchlorate monohydrate</i>	Meng <i>et al.</i> (2008d)	10.1107/S1600536808000287	GISLEA
<i>Diaquabis(pyridine-2-carboxylato-κ²N,O)cobalt(II)</i>	Huang (2008)	10.1107/S1600536808010507	WIZPOL
<i>Tetra-μ-2,5-difluorobenzoato-bis[(2,2'-bipyridine)(2,5-difluorobenzoato)gadolinium(III)]</i>	Li, Zhang <i>et al.</i> (2008)	10.1107/S1600536808023507	BOFQIX
<i>catena-Poly[[[2,2'-bipyridine-κ²N,N']nickel(II)]-μ-oxalato-κ⁴O¹,O²:O¹,O²]</i>	Li, Yan <i>et al.</i> (2008)	10.1107/S1600536808028389	NOHYUF
<i>catena-Poly[[aqua(2,2'-bipyridyl)cobalt(II)]-μ-5-nitrosophthalato]</i>	Liu <i>et al.</i> (2008)	10.1107/S1600536808038178	AFIREN
<i>Diaquabis(pyridine-2-carboxylato-κ²N,O)iron(II)</i>	Xia & Sun (2009)	10.1107/S1600536809005765	RONFEG
<i>catena-Poly[[[diaquathulium(III)]-μ-6-carboxynicotinato-μ-pyridine-2,5-dicarboxylato] dihydrate]</i>	Li <i>et al.</i> (2009)	10.1107/S1600536809008836	NOQNIR
<i>1-Phenyl-3-(2,4,6-trimethoxyphenyl)prop-2-en-1-one</i>	Liu <i>et al.</i> (2009)	10.1107/S1600536809040227	PUGLOT

References

- Harrison, W. T. A., Simpson, J. & Weil, M. (2010). *Acta Cryst.* **E66**, e1–e2.
- Huang, G. S. (2008). *Acta Cryst.* **E64**, m685–m686.
- Li, S., Chen, Y., He, H.-M. & Ma, Y.-F. (2009). *Acta Cryst.* **E65**, m411.
- Li, S., Wang, S.-B., Zhang, F.-L. & Tang, K. (2008). *Acta Cryst.* **E64**, m2.
- Li, S., Yan, X.-L., Wang, S.-B. & Ma, Y.-F. (2008). *Acta Cryst.* **E64**, m1258.
- Li, S., Zhang, F.-L., Tang, K. & Ma, Y.-F. (2008). *Acta Cryst.* **E64**, m1142.
- Li, Z., Wang, S., Zhang, Q. & Yu, X. (2007a). *Acta Cryst.* **E63**, m2438–m2439.
- Li, Z., Wang, S., Zhang, Q. & Yu, X. (2007b). *Acta Cryst.* **E63**, m2604.
- Li, Z., Wang, S., Zhang, Q. & Yu, X. (2007c). *Acta Cryst.* **E63**, m2642.
- Li, Z., Wang, S., Zhang, Q. & Yu, X. (2007d). *Acta Cryst.* **E63**, m2781.
- Li, Z.-F., Liu, Y., Zhang, Q. & Yu, X.-J. (2007). *Acta Cryst.* **E63**, m2315.
- Li, Z.-F., Wang, S.-W., Zhang, Q. & Yu, X.-J. (2007e). *Acta Cryst.* **E63**, m2312.
- Li, Z.-F., Wang, S.-W., Zhang, Q. & Yu, X.-J. (2007f). *Acta Cryst.* **E63**, m2360.
- Li, Z.-F., Wang, S.-W., Zhang, Q. & Yu, X.-J. (2007g). *Acta Cryst.* **E63**, m2373.
- Li, Z.-F., Wang, S.-W., Zhang, Q. & Yu, X.-J. (2007h). *Acta Cryst.* **E63**, m2445.
- Li, Z.-F., Wang, S.-W., Zhang, Q. & Yu, X.-J. (2007i). *Acta Cryst.* **E63**, o3930.
- Liu, Y., Dou, J., Li, D. & Zhang, X. (2007). *Acta Cryst.* **E63**, m2722.
- Liu, Y., Dou, J., Niu, M. & Zhang, X. (2007a). *Acta Cryst.* **E63**, m2771.
- Liu, Y., Dou, J., Niu, M. & Zhang, X. (2007b). *Acta Cryst.* **E63**, m3032.
- Liu, Y., Dou, J., Wang, D., Ma, G. & Li, D. (2005). *Acta Cryst.* **E61**, m1834–m1836.
- Liu, Y., Dou, J.-M., Wang, D.-Q., Zhang, X.-X. & Zhou, L. (2006). *Acta Cryst.* **E62**, m2794–m2795.
- Liu, Y., He, Q., Zhang, X., Xue, Z. & Lv, C. (2008). *Acta Cryst.* **E64**, m1605–m1606.
- Liu, Y., Zhang, X., Xue, Z. & Lv, C. (2009). *Acta Cryst.* **E65**, o2724.
- Meng, Q., Wang, L., Liu, Y. & Pang, Y. (2008a). *Acta Cryst.* **E64**, m63.
- Meng, Q., Wang, L., Liu, Y. & Pang, Y. (2008b). *Acta Cryst.* **E64**, m133.
- Meng, Q., Wang, L., Liu, Y. & Pang, Y. (2008c). *Acta Cryst.* **E64**, m204.
- Meng, Q., Wang, L., Liu, Y. & Pang, Y. (2008d). *Acta Cryst.* **E64**, m332.
- Meng, Q.-G., Wang, L.-T., Liu, Y.-Z. & Pang, Y. (2008e). *Acta Cryst.* **E64**, m170–m171.
- Xia, G. & Sun, Z. (2009). *Acta Cryst.* **E65**, m315–m316.

Oxalatobis(propane-1,3-diamine)-manganese(II) chloride monohydrate

Qing-Guo Meng,* Lin-Tong Wang, Yan-Zhen Liu and Yan Pang

College of Chemistry and Chemical Engineering, Weifang University, Weifang 261061, People's Republic of China

Correspondence e-mail: qgmeng_weifang@yahoo.cn


Received 18 November 2007; accepted 4 December 2007

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.032; wR factor = 0.085; data-to-parameter ratio = 8.6.

In the asymmetric unit of the title compound, $[\text{Mn}(\text{C}_2\text{O}_4)(\text{C}_3\text{H}_7\text{N}_2)_2]\text{Cl}\cdot\text{H}_2\text{O}$, there are two independent Mn^{II} complexes, two Cl^- anions and two uncoordinated water molecules. Each Mn^{II} atom is hexacoordinated by four N atoms from two propane-1,3-diamine ligands and two O atoms from one oxalate ligand, resulting in a slightly distorted octahedral MnO_2N_4 geometry. Mn—O and Mn—N bond lengths are in the ranges 1.969 (2)–2.020 (3) and 2.068 (3)–2.113 (4) Å, respectively. There are weak intermolecular O—H...O, O—H...Cl, N—H...O and N—H...Cl hydrogen bonds with $D\cdots A$ distances in the range 2.831 (4)–3.423 (3) Å.

Related literature

For related literature, see: Chung *et al.* (1971); Church & Halvorson (1959); Okabe & Oya (2000); Pocker & Fong (1980); Poowell (1953); Scapin *et al.* (1997); Serre *et al.* (2005).



Experimental

Crystal data

$[\text{Mn}(\text{C}_2\text{O}_4)(\text{C}_3\text{H}_7\text{N}_2)_2]\text{Cl}\cdot\text{H}_2\text{O}$
 $M_r = 344.69$
 Monoclinic, Pn
 $a = 9.1286$ (17) Å
 $b = 11.807$ (2) Å
 $c = 13.912$ (3) Å
 $\beta = 100.037$ (14)°
 $V = 1476.6$ (5) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 1.10$ mm⁻¹
 $T = 293$ (2) K
 $0.43 \times 0.28 \times 0.23$ mm

Data collection

Bruker APEXII CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2001)
 $T_{\text{min}} = 0.650$, $T_{\text{max}} = 0.787$
 3373 measured reflections
 3060 independent reflections
 3032 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.085$
 $S = 1.00$
 3060 reflections
 356 parameters
 8 restraints
 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.53$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.43$ e Å⁻³
 Absolute structure: Flack (1983), 521 Friedel pairs
 Flack parameter: 0.040 (15)

Table 1
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O10—H3W...Cl2	0.81 (6)	2.70 (7)	3.205 (5)	122 (7)
O10—H4W...O6	0.81 (5)	2.71 (7)	3.078 (5)	110 (5)
O9—H2W...Cl1 ⁱ	0.83 (3)	2.55 (2)	3.348 (4)	163 (5)
O9—H1W...O10 ⁱⁱ	0.82 (5)	2.03 (5)	2.845 (6)	170 (6)
N8—H8B...Cl2 ⁱ	0.90	2.61	3.366 (3)	142
N8—H8A...O3 ⁱⁱⁱ	0.90	2.59	3.426 (4)	154
N7—H7B...O4 ⁱⁱⁱ	0.90	2.45	3.195 (4)	141
N7—H7A...O8 ^{iv}	0.90	2.13	3.014 (4)	169
N6—H6D...O9	0.90	2.20	3.054 (4)	157
N6—H6C...O8 ^{iv}	0.90	2.21	3.046 (4)	155
N5—H5D...O4 ⁱⁱⁱ	0.90	2.04	2.942 (4)	176
N5—H5C...Cl1 ^v	0.90	2.41	3.299 (3)	172
N4—H4D...Cl1 ^{vi}	0.90	2.53	3.423 (3)	170
N4—H4C...O2 ^{vii}	0.90	2.42	3.206 (4)	147
N3—H3D...Cl2	0.90	2.25	3.115 (3)	162
N3—H3C...O5 ^{viii}	0.90	2.02	2.831 (4)	149
N2—H2D...Cl1 ^{vi}	0.90	2.54	3.317 (4)	144
N2—H2C...O8 ^{viii}	0.90	2.15	3.024 (5)	163
N1—H1D...Cl2	0.90	2.71	3.404 (3)	135
N1—H1C...Cl1 ⁱⁱⁱ	0.90	2.55	3.388 (3)	155

Symmetry codes: (i) $x - 1, y, z$; (ii) $x - \frac{1}{2}, -y + 1, z - \frac{1}{2}$; (iii) $x + \frac{1}{2}, -y + 1, z + \frac{1}{2}$; (iv) $x - 1, y - 1, z$; (v) $x - \frac{1}{2}, -y + 1, z + \frac{1}{2}$; (vi) $x, y + 1, z$; (vii) $x + 1, y + 1, z$; (viii) $x - \frac{1}{2}, -y + 2, z - \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT-Plus (Bruker, 2001); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXS97 (Sheldrick, 1997); program(s) used to refine structure: SHELXL97 (Sheldrick, 1997); molecular graphics: SHELXTL (Bruker, 2001); software used to prepare material for publication: SHELXTL.

The authors thank the Education Department of Shandong Province for research and development projects (No. J06A55).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS2256).

References

- Bruker (2001). *SADABS*, *SAINT-Plus* and *SHELXTL*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2004). *APEX2*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chung, L., Rajan, K. S., Merdinger, E. & Crez, N. (1971). *Biophys. J.* **469**, 469–472.
- Church, B. S. & Halvorson, H. (1959). *Nature (London)*, **183**, 124–125.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Okabe, N. & Oya, N. (2000). *Acta Cryst.* **C56**, 1416–1417.
- Pocker, Y. & Fong, C. T. O. (1980). *Biochemistry*, **16**, 2045–2049.
- Poowell, J. F. (1953). *Biochemistry*, **54**, 205–207.
- Scapin, G., Reddy, S. G., Zheng, R. & Blanchard, J. S. (1997). *Biochemistry*, **36**, 15081–15092.
- Serre, C., Marrot, J. & Ferey, G. (2005). *Inorg. Chem.* **44**, 654–657.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.

Article retracted

supporting information

Acta Cryst. (2008). E64, m170–m171 [https://doi.org/10.1107/S1600536807065361]

Oxalatobis(propane-1,3-diamine)manganese(II) chloride monohydrate**Qing-Guo Meng, Lin-Tong Wang, Yan-Zhen Liu and Yan Pang****S1. Comment**

During the past decades, dicarboxylic acid has been widely used as one poly-dentate ligand involved in various metal chelation reactions to form transition or rare earth metal complexes that own thermal resistance of certain bacteria spores (Poowell, 1953; Church & Halvorson, 1959; Chung *et al.*, 1971; Okabe & Oya, 2000) and the activation (Serre *et al.*, 2005) or inhabitation (Pocker & Fong, 1980; Scapin *et al.*, 1997) in some metallo-enzymes. In this paper, we report the structure of the title compound, (I).


In the title compound, the Mn atom is hexa-coordinated by four nitrogen atoms from two chelating propane-1,3-diamine ligands and two oxygen atoms from one oxalic acid, resulting in a slightly distorted octahedral MnO_2N_4 geometry for the metal (Fig. 1, Table 1). Mn—O and Mn—N bond lengths are in the range of 1.969 (2) - 2.020 (3) Å and 2.068 (3) - 2.113 (4) Å, respectively. Moreover, there exist weak intermolecular hydrogen bonds with the distance range of 2.830–3.423 Å (Table 2), forming a three-dimensional structure (Fig. 2).

S2. Experimental


A mixed water and ethanol solution of manganese(III) acetate (1 mmol) and oxalic acid (1 mmol) was neutralized by propane-1,3-diamine. The resulted solution was saturated with 1 g sodium hydrochloride and evaporated at room temperature for one week. yellow block crystals were obtained with a yield of 21%. Anal. Calc. for $\text{C}_8\text{H}_{22}\text{ClMnN}_4\text{O}_5$: C 27.83, H 6.38, N 16.23%; Found: C 27.80, H 6.42, N 16.18%.

S3. Refinement

The H atoms of the water molecule were located in a difference Fourier map and were refined with distance restraints of $\text{H}\cdots\text{H} = 1.38$ (2) Å and $\text{O}-\text{H} = 0.82$ (2) Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{O})$. Other H atoms were placed in calculated positions ($\text{C}-\text{H} = 0.93$ Å and $\text{N}-\text{H} = 0.90$ Å) and treated as riding, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$.

**Figure 1**

The asymmetric unit of the title compound, drawn with 30% probability displacement ellipsoids for the non-hydrogen atoms.

**Figure 2**

A packing diagram of the title compound. Hydrogen bonds are indicated by dashed lines.

Oxalatobis(propane-1,3-diamine)manganese(II) chloride monohydrate

Crystal data

[Mn(C₂O₄)(C₃H₁₀N₂)₂]Cl·H₂O $M_r = 344.69$ Monoclinic, Pn Hall symbol: $P -2yc$ $a = 9.1286 (17) \text{ \AA}$ $b = 11.807 (2) \text{ \AA}$ $c = 13.912 (3) \text{ \AA}$ $\beta = 100.037 (14)^\circ$ $V = 1476.6 (5) \text{ \AA}^3$ $Z = 4$ $F(000) = 720$ $D_x = 1.551 \text{ Mg m}^{-3}$ Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 3060 reflections

 $\theta = 2.9\text{--}25.0^\circ$ $\mu = 1.10 \text{ mm}^{-1}$ $T = 293 \text{ K}$

Block, yellow

 $0.43 \times 0.28 \times 0.23 \text{ mm}$

Data collection

Bruker APEXII CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 φ and ω scansAbsorption correction: multi-scan
(*SADABS*; Bruker, 2001) $T_{\min} = 0.650$, $T_{\max} = 0.787$

3373 measured reflections

3060 independent reflections

3032 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.023$ $\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 2.3^\circ$ $h = -1 \rightarrow 10$ $k = -14 \rightarrow 1$ $l = -16 \rightarrow 16$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.032$ $wR(F^2) = 0.085$ $S = 1.00$

3060 reflections

356 parameters

8 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sitesH atoms treated by a mixture of independent
and constrained refinement $w = 1/[\sigma^2(F_o^2) + (0.072P)^2 + 0.2126P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.53 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -0.43 \text{ e \AA}^{-3}$ Extinction correction: *SHELXL*, $F_c^* = kF_c[1 + 0.001 \times F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0164 (13)

Absolute structure: Flack (1983), 521 Friedel
pairs

Absolute structure parameter: 0.040 (15)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Mn1	0.11279 (5)	0.45504 (4)	0.50737 (3)	0.02681 (15)

Mn2	0.91703 (6)	0.99201 (4)	0.36505 (4)	0.02996 (16)
C1	0.3604 (4)	0.3805 (3)	0.3916 (3)	0.0360 (8)
H1A	0.3263	0.4355	0.3409	0.043*
H1B	0.4034	0.3172	0.3619	0.043*
C2	0.4805 (5)	0.4350 (4)	0.4694 (3)	0.0404 (9)
H2A	0.5053	0.3832	0.5239	0.048*
H2B	0.5697	0.4471	0.4417	0.048*
C3	0.4322 (5)	0.5455 (3)	0.5062 (3)	0.0378 (9)
H3A	0.3975	0.5942	0.4508	0.045*
H3B	0.5178	0.5818	0.5453	0.045*
C4	-0.1361 (5)	0.5164 (3)	0.6257 (3)	0.0373 (8)
H4A	-0.2080	0.4832	0.5738	0.045*
H4B	-0.1848	0.5778	0.6541	0.045*
C5	-0.0860 (5)	0.4271 (3)	0.7035 (3)	0.0401 (9)
H5A	-0.0135	0.4604	0.7550	0.048*
H5B	-0.1709	0.4035	0.7319	0.048*
C6	-0.0192 (5)	0.3260 (3)	0.6642 (3)	0.0351 (8)
H6A	-0.0886	0.2967	0.6090	0.042*
H6B	-0.0038	0.2677	0.7140	0.042*
C7	-0.1264 (4)	0.4394 (3)	0.3465 (3)	0.0289 (7)
C8	-0.0207 (4)	0.5353 (3)	0.3226 (3)	0.0308 (8)
C9	0.5848 (5)	0.9640 (4)	0.3832 (3)	0.0417 (9)
H9A	0.4953	0.9199	0.3620	0.050*
H9B	0.6156	0.9518	0.4527	0.050*
C10	0.5504 (5)	1.0872 (4)	0.3649 (4)	0.0486 (10)
H10A	0.4667	1.1079	0.3957	0.058*
H10B	0.5214	1.0994	0.2952	0.058*
C11	0.6810 (5)	1.1632 (3)	0.4035 (3)	0.0431 (9)
H11A	0.6490	1.2416	0.3972	0.052*
H11B	0.7139	1.1477	0.4723	0.052*
C12	1.0022 (5)	1.0344 (3)	0.5651 (3)	0.0377 (9)
C13	1.1333 (5)	1.0806 (3)	0.5152 (3)	0.0360 (8)
C14	1.1008 (6)	0.9676 (4)	0.1977 (3)	0.0478 (11)
H14A	1.1031	0.9793	0.1290	0.057*
H14B	1.1750	1.0167	0.2349	0.057*
C15	1.1384 (6)	0.8483 (4)	0.2233 (4)	0.0516 (11)
H15A	1.2274	0.8282	0.1979	0.062*
H15B	1.0582	0.8004	0.1913	0.062*
C16	1.1650 (5)	0.8233 (4)	0.3342 (3)	0.0428 (9)
H16A	1.2009	0.7463	0.3452	0.051*
H16B	1.2415	0.8738	0.3670	0.051*
N1	1.0262 (4)	0.8377 (2)	0.3781 (2)	0.0324 (6)
H1C	1.0502	0.8218	0.4422	0.039*
H1D	0.9606	0.7847	0.3515	0.039*
N2	0.9487 (5)	0.9993 (3)	0.2183 (2)	0.0375 (8)
H2C	0.8814	0.9534	0.1829	0.045*
H2D	0.9284	1.0704	0.1965	0.045*
N3	0.7056 (4)	0.9230 (2)	0.3304 (2)	0.0327 (7)

H3C	0.6738	0.9341	0.2662	0.039*
H3D	0.7143	0.8477	0.3398	0.039*
N4	0.8095 (4)	1.1468 (3)	0.3503 (2)	0.0340 (7)
H4C	0.8773	1.2009	0.3706	0.041*
H4D	0.7762	1.1586	0.2863	0.041*
N5	0.3116 (4)	0.5354 (2)	0.5663 (2)	0.0291 (6)
H5C	0.2889	0.6059	0.5835	0.035*
H5D	0.3501	0.4987	0.6217	0.035*
N6	0.2302 (4)	0.3396 (2)	0.4346 (2)	0.0297 (6)
H6C	0.2630	0.2837	0.4769	0.036*
H6D	0.1648	0.3083	0.3860	0.036*
N7	0.1268 (4)	0.3507 (2)	0.6321 (2)	0.0313 (6)
H7A	0.1688	0.2845	0.6197	0.038*
H7B	0.1877	0.3841	0.6818	0.038*
N8	-0.0078 (4)	0.5630 (2)	0.5835 (2)	0.0319 (6)
H8A	0.0568	0.5937	0.6330	0.038*
H8B	-0.0430	0.6199	0.5430	0.038*
O1	0.0942 (3)	0.55365 (19)	0.39175 (18)	0.0302 (5)
O2	-0.0783 (3)	0.3917 (2)	0.43141 (18)	0.0301 (5)
O3	-0.2391 (3)	0.4152 (2)	0.2880 (2)	0.0402 (6)
O4	-0.0474 (4)	0.5855 (3)	0.24339 (19)	0.0440 (7)
O5	1.0023 (5)	1.0508 (3)	0.6534 (2)	0.0583 (9)
O6	0.8993 (3)	0.9836 (2)	0.50604 (19)	0.0362 (6)
O7	1.1132 (3)	1.0658 (2)	0.42064 (19)	0.0365 (6)
O8	1.2408 (4)	1.1305 (2)	0.5639 (2)	0.0457 (7)
O9	0.0571 (4)	0.2740 (3)	0.2343 (2)	0.0600 (9)
H1W	0.114 (5)	0.246 (5)	0.201 (4)	0.072*
H2W	-0.031 (2)	0.256 (6)	0.214 (4)	0.072*
O10	0.7709 (7)	0.7953 (3)	0.6200 (3)	0.0795 (13)
H3W	0.828 (7)	0.772 (6)	0.586 (4)	0.095*
H4W	0.728 (8)	0.854 (4)	0.603 (5)	0.095*
Cl1	0.72741 (14)	0.19856 (8)	0.10470 (7)	0.0495 (3)
Cl2	0.72971 (12)	0.67594 (8)	0.41008 (8)	0.0427 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Mn1	0.0300 (3)	0.0243 (2)	0.0252 (2)	-0.0010 (2)	0.00198 (19)	0.00041 (19)
Mn2	0.0339 (3)	0.0262 (3)	0.0277 (3)	-0.0024 (2)	-0.0005 (2)	0.00098 (18)
C1	0.0375 (19)	0.0338 (18)	0.0389 (18)	-0.0004 (16)	0.0132 (17)	-0.0015 (14)
C2	0.034 (2)	0.0431 (19)	0.044 (2)	0.0002 (18)	0.0074 (18)	0.0010 (17)
C3	0.038 (2)	0.036 (2)	0.039 (2)	-0.0106 (16)	0.0072 (17)	-0.0029 (15)
C4	0.039 (2)	0.0368 (18)	0.0375 (19)	0.0047 (16)	0.0108 (17)	-0.0016 (15)
C5	0.044 (2)	0.040 (2)	0.038 (2)	0.0020 (19)	0.0125 (18)	0.0046 (16)
C6	0.042 (2)	0.0303 (17)	0.0341 (18)	-0.0043 (16)	0.0088 (17)	0.0045 (14)
C7	0.0311 (19)	0.0277 (15)	0.0270 (16)	-0.0012 (14)	0.0025 (15)	-0.0032 (12)
C8	0.037 (2)	0.0277 (17)	0.0276 (17)	0.0020 (15)	0.0042 (16)	0.0006 (13)
C9	0.037 (2)	0.044 (2)	0.045 (2)	-0.0050 (18)	0.0078 (18)	0.0003 (17)

C10	0.037 (2)	0.045 (2)	0.062 (3)	0.0071 (19)	0.005 (2)	-0.002 (2)
C11	0.049 (2)	0.0307 (18)	0.051 (2)	0.0075 (18)	0.0103 (19)	-0.0051 (16)
C12	0.048 (2)	0.0302 (17)	0.032 (2)	0.0042 (17)	-0.0012 (18)	0.0017 (14)
C13	0.036 (2)	0.0230 (15)	0.043 (2)	0.0026 (16)	-0.0082 (17)	0.0000 (15)
C14	0.061 (3)	0.044 (2)	0.044 (2)	0.001 (2)	0.022 (2)	0.0083 (18)
C15	0.068 (3)	0.040 (2)	0.052 (2)	0.013 (2)	0.025 (2)	0.0015 (19)
C16	0.039 (2)	0.0382 (19)	0.052 (2)	0.0076 (17)	0.0103 (18)	0.0076 (17)
N1	0.0350 (15)	0.0268 (13)	0.0335 (15)	-0.0003 (13)	0.0005 (13)	0.0046 (12)
N2	0.052 (2)	0.0294 (16)	0.0305 (16)	0.0041 (14)	0.0041 (15)	0.0026 (11)
N3	0.0333 (16)	0.0261 (14)	0.0348 (15)	-0.0006 (13)	-0.0049 (13)	0.0040 (12)
N4	0.0404 (17)	0.0233 (13)	0.0361 (16)	-0.0014 (13)	0.0003 (14)	0.0002 (11)
N5	0.0293 (16)	0.0256 (14)	0.0298 (15)	-0.0026 (12)	-0.0021 (13)	-0.0019 (11)
N6	0.0374 (16)	0.0235 (12)	0.0284 (14)	-0.0014 (13)	0.0063 (12)	0.0001 (11)
N7	0.0385 (16)	0.0265 (13)	0.0276 (14)	-0.0007 (13)	0.0023 (13)	0.0000 (11)
N8	0.0369 (17)	0.0257 (12)	0.0322 (15)	0.0019 (13)	0.0033 (14)	-0.0001 (11)
O1	0.0371 (14)	0.0263 (11)	0.0244 (11)	-0.0045 (10)	-0.0022 (10)	0.0027 (9)
O2	0.0336 (12)	0.0258 (10)	0.0291 (11)	-0.0037 (10)	0.0003 (10)	-0.0006 (9)
O3	0.0378 (16)	0.0410 (14)	0.0380 (13)	-0.0050 (13)	-0.0041 (13)	-0.0037 (12)
O4	0.0555 (18)	0.0412 (14)	0.0300 (13)	-0.0042 (14)	-0.0072 (13)	0.0097 (11)
O5	0.073 (2)	0.069 (2)	0.0287 (15)	-0.0011 (19)	-0.0039 (16)	-0.0109 (13)
O6	0.0444 (17)	0.0346 (12)	0.0281 (13)	-0.0032 (12)	0.0018 (12)	0.0045 (10)
O7	0.0375 (15)	0.0334 (13)	0.0356 (14)	-0.0069 (12)	-0.0019 (12)	0.0007 (11)
O8	0.0472 (17)	0.0290 (13)	0.0522 (16)	-0.0039 (12)	-0.0155 (14)	-0.0002 (12)
O9	0.059 (2)	0.074 (2)	0.0471 (17)	0.0031 (19)	0.0081 (16)	-0.0112 (16)
O10	0.122 (4)	0.048 (2)	0.081 (3)	-0.007 (2)	0.052 (3)	-0.0004 (19)
C11	0.0658 (7)	0.0332 (4)	0.0420 (5)	-0.0013 (5)	-0.0118 (5)	0.0025 (4)
C12	0.0448 (5)	0.0307 (4)	0.0529 (5)	0.0008 (4)	0.0090 (4)	0.0068 (4)

Geometric parameters (Å, °)

Mn1—O1	1.969 (2)	C10—C11	1.514 (7)
Mn1—O2	2.019 (3)	C10—H10A	0.9700
Mn1—N5	2.085 (3)	C10—H10B	0.9700
Mn1—N8	2.089 (3)	C11—N4	1.504 (5)
Mn1—N6	2.100 (3)	C11—H11A	0.9700
Mn1—N7	2.113 (3)	C11—H11B	0.9700
Mn2—O6	1.999 (3)	C12—O5	1.243 (5)
Mn2—O7	2.020 (3)	C12—O6	1.284 (5)
Mn2—N4	2.068 (3)	C12—C13	1.582 (6)
Mn2—N1	2.069 (3)	C13—O8	1.241 (5)
Mn2—N3	2.073 (3)	C13—O7	1.307 (5)
Mn2—N2	2.113 (4)	C14—C15	1.478 (6)
C1—N6	1.500 (5)	C14—N2	1.513 (7)
C1—C2	1.541 (6)	C14—H14A	0.9700
C1—H1A	0.9700	C14—H14B	0.9700
C1—H1B	0.9700	C15—C16	1.548 (6)
C2—C3	1.495 (6)	C15—H15A	0.9700
C2—H2A	0.9700	C15—H15B	0.9700

C2—H2B	0.9700	C16—N1	1.509 (5)
C3—N5	1.499 (5)	C16—H16A	0.9700
C3—H3A	0.9700	C16—H16B	0.9700
C3—H3B	0.9700	N1—H1C	0.9000
C4—N8	1.503 (5)	N1—H1D	0.9000
C4—C5	1.523 (6)	N2—H2C	0.9000
C4—H4A	0.9700	N2—H2D	0.9000
C4—H4B	0.9700	N3—H3C	0.9000
C5—C6	1.488 (6)	N3—H3D	0.9000
C5—H5A	0.9700	N4—H4C	0.9000
C5—H5B	0.9700	N4—H4D	0.9000
C6—N7	1.506 (5)	N5—H5C	0.9000
C6—H6A	0.9700	N5—H5D	0.9000
C6—H6B	0.9700	N6—H6C	0.9000
C7—O3	1.229 (5)	N6—H6D	0.9000
C7—O2	1.314 (4)	N7—H7A	0.9000
C7—C8	1.560 (5)	N7—H7B	0.9000
C8—O4	1.238 (5)	N8—H8A	0.9000
C8—O1	1.311 (5)	N8—H8B	0.9000
C9—C10	1.501 (6)	O9—H1W	0.82 (5)
C9—N3	1.507 (6)	O9—H2W	0.83 (3)
C9—H9A	0.9700	O10—H3W	0.81 (6)
C9—H9B	0.9700	O10—H4W	0.81 (5)
O1—Mn1—O2	81.23 (11)	N4—C11—C10	112.5 (3)
O1—Mn1—N5	89.86 (12)	N4—C11—H11A	109.1
O2—Mn1—N5	171.03 (11)	C10—C11—H11A	109.1
O1—Mn1—N8	94.33 (11)	N4—C11—H11B	109.1
O2—Mn1—N8	90.45 (12)	C10—C11—H11B	109.1
N5—Mn1—N8	91.28 (13)	H11A—C11—H11B	107.8
O1—Mn1—N6	88.21 (11)	O5—C12—O6	125.4 (4)
O2—Mn1—N6	88.53 (11)	O5—C12—C13	120.6 (4)
N5—Mn1—N6	90.15 (12)	O6—C12—C13	113.9 (3)
N8—Mn1—N6	177.09 (12)	O8—C13—O7	124.9 (4)
O1—Mn1—N7	178.39 (13)	O8—C13—C12	120.7 (4)
O2—Mn1—N7	97.88 (11)	O7—C13—C12	114.3 (3)
N5—Mn1—N7	91.05 (12)	C15—C14—N2	111.9 (4)
N8—Mn1—N7	84.33 (12)	C15—C14—H14A	109.2
N6—Mn1—N7	93.11 (12)	N2—C14—H14A	109.2
O6—Mn2—O7	82.17 (12)	C15—C14—H14B	109.2
O6—Mn2—N4	91.31 (12)	N2—C14—H14B	109.2
O7—Mn2—N4	91.72 (12)	H14A—C14—H14B	107.9
O6—Mn2—N1	89.44 (12)	C14—C15—C16	114.3 (4)
O7—Mn2—N1	88.01 (12)	C14—C15—H15A	108.7
N4—Mn2—N1	179.16 (14)	C16—C15—H15A	108.7
O6—Mn2—N3	88.68 (13)	C14—C15—H15B	108.7
O7—Mn2—N3	170.31 (13)	C16—C15—H15B	108.7
N4—Mn2—N3	85.28 (13)	H15A—C15—H15B	107.6

N1—Mn2—N3	95.11 (13)	N1—C16—C15	112.7 (4)
O6—Mn2—N2	176.79 (16)	N1—C16—H16A	109.1
O7—Mn2—N2	95.20 (14)	C15—C16—H16A	109.1
N4—Mn2—N2	90.61 (13)	N1—C16—H16B	109.1
N1—Mn2—N2	88.62 (13)	C15—C16—H16B	109.1
N3—Mn2—N2	94.04 (15)	H16A—C16—H16B	107.8
N6—C1—C2	111.7 (3)	C16—N1—Mn2	119.3 (2)
N6—C1—H1A	109.3	C16—N1—H1C	107.5
C2—C1—H1A	109.3	Mn2—N1—H1C	107.5
N6—C1—H1B	109.3	C16—N1—H1D	107.5
C2—C1—H1B	109.3	Mn2—N1—H1D	107.5
H1A—C1—H1B	107.9	H1C—N1—H1D	107.0
C3—C2—C1	112.9 (3)	C14—N2—Mn2	117.2 (3)
C3—C2—H2A	109.0	C14—N2—H2C	108.0
C1—C2—H2A	109.0	Mn2—N2—H2C	108.0
C3—C2—H2B	109.0	C14—N2—H2D	108.0
C1—C2—H2B	109.0	Mn2—N2—H2D	108.0
H2A—C2—H2B	107.8	H2C—N2—H2D	107.2
C2—C3—N5	114.3 (3)	C9—N3—Mn2	119.5 (2)
C2—C3—H3A	108.7	C9—N3—H3C	107.5
N5—C3—H3A	108.7	Mn2—N3—H3C	107.5
C2—C3—H3B	108.7	C9—N3—H3D	107.5
N5—C3—H3B	108.7	Mn2—N3—H3D	107.5
H3A—C3—H3B	107.6	H3C—N3—H3D	107.0
N8—C4—C5	111.9 (3)	C11—N4—Mn2	117.5 (2)
N8—C4—H4A	109.2	C11—N4—H4C	107.9
C5—C4—H4A	109.2	Mn2—N4—H4C	107.9
N8—C4—H4B	109.2	C11—N4—H4D	107.9
C5—C4—H4B	109.2	Mn2—N4—H4D	107.9
H4A—C4—H4B	107.9	H4C—N4—H4D	107.2
C6—C5—C4	112.7 (3)	C3—N5—Mn1	119.4 (2)
C6—C5—H5A	109.1	C3—N5—H5C	107.5
C4—C5—H5A	109.1	Mn1—N5—H5C	107.5
C6—C5—H5B	109.1	C3—N5—H5D	107.5
C4—C5—H5B	109.1	Mn1—N5—H5D	107.5
H5A—C5—H5B	107.8	H5C—N5—H5D	107.0
C5—C6—N7	113.2 (3)	C1—N6—Mn1	119.3 (2)
C5—C6—H6A	108.9	C1—N6—H6C	107.5
N7—C6—H6A	108.9	Mn1—N6—H6C	107.5
C5—C6—H6B	108.9	C1—N6—H6D	107.5
N7—C6—H6B	108.9	Mn1—N6—H6D	107.5
H6A—C6—H6B	107.7	H6C—N6—H6D	107.0
O3—C7—O2	127.3 (4)	C6—N7—Mn1	115.2 (2)
O3—C7—C8	120.4 (3)	C6—N7—H7A	108.5
O2—C7—C8	112.3 (3)	Mn1—N7—H7A	108.5
O4—C8—O1	125.6 (4)	C6—N7—H7B	108.5
O4—C8—C7	120.2 (3)	Mn1—N7—H7B	108.5
O1—C8—C7	114.2 (3)	H7A—N7—H7B	107.5

C10—C9—N3	112.2 (4)	C4—N8—Mn1	119.4 (2)
C10—C9—H9A	109.2	C4—N8—H8A	107.5
N3—C9—H9A	109.2	Mn1—N8—H8A	107.5
C10—C9—H9B	109.2	C4—N8—H8B	107.5
N3—C9—H9B	109.2	Mn1—N8—H8B	107.5
H9A—C9—H9B	107.9	H8A—N8—H8B	107.0
C9—C10—C11	112.7 (4)	C8—O1—Mn1	116.5 (2)
C9—C10—H10A	109.1	C7—O2—Mn1	115.8 (2)
C11—C10—H10A	109.1	C12—O6—Mn2	115.4 (3)
C9—C10—H10B	109.1	C13—O7—Mn2	113.7 (3)
C11—C10—H10B	109.1	H1W—O9—H2W	112 (5)
H10A—C10—H10B	107.8	H3W—O10—H4W	117 (7)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O10—H3W \cdots C12	0.81 (6)	2.70 (7)	3.205 (5)	122 (7)
O10—H4W \cdots O6	0.81 (5)	2.71 (7)	3.078 (5)	110 (5)
O9—H2W \cdots C11 ⁱ	0.83 (3)	2.55 (2)	3.348 (4)	163 (5)
O9—H1W \cdots O10 ⁱⁱ	0.82 (5)	2.03 (5)	2.845 (6)	170 (6)
N8—H8B \cdots C12 ⁱ	0.90	2.61	3.366 (3)	142
N8—H8A \cdots O3 ⁱⁱⁱ	0.90	2.59	3.426 (4)	154
N7—H7B \cdots O4 ⁱⁱⁱ	0.90	2.45	3.195 (4)	141
N7—H7A \cdots O8 ^{iv}	0.90	2.13	3.014 (4)	169
N6—H6D \cdots O9	0.90	2.20	3.054 (4)	157
N6—H6C \cdots O8 ^{iv}	0.90	2.21	3.046 (4)	155
N5—H5D \cdots O4 ⁱⁱⁱ	0.90	2.04	2.942 (4)	176
N5—H5C \cdots C11 ^v	0.90	2.41	3.299 (3)	172
N4—H4D \cdots C11 ^{vi}	0.90	2.53	3.423 (3)	170
N4—H4C \cdots O2 ^{vii}	0.90	2.42	3.206 (4)	147
N3—H3D \cdots C12	0.90	2.25	3.115 (3)	162
N3—H3C \cdots O5 ^{viii}	0.90	2.02	2.831 (4)	149
N2—H2D \cdots C11 ^{vi}	0.90	2.54	3.317 (4)	144
N2—H2C \cdots O8 ^{viii}	0.90	2.15	3.024 (5)	163
N1—H1D \cdots C12	0.90	2.71	3.404 (3)	135
N1—H1C \cdots C11 ⁱⁱⁱ	0.90	2.55	3.388 (3)	155

Symmetry codes: (i) $x-1, y, z$; (ii) $x-1/2, -y+1, z-1/2$; (iii) $x+1/2, -y+1, z+1/2$; (iv) $x-1, y-1, z$; (v) $x-1/2, -y+1, z+1/2$; (vi) $x, y+1, z$; (vii) $x+1, y+1, z$; (viii) $x-1/2, -y+2, z-1/2$.