

2-(4,5-Dihydro-1,3-oxazol-2-yl)quinoline

Pablo Machado,^a Fernanda A. Rosa,^a Rubia M. S. da Silva,^b
Robert A. Burrow^{b*} and Marcos A. P. Martins^{a*}

^aNúcleo de Química de Heterociclos (NUQUIMHE), Departamento de Química, Universidade Federal de Santa Maria, 97105-900, Santa Maria, RS, Brazil, and

^bLaboratório de Materiais Inorgânicos, Departamento de Química, Universidade Federal de Santa Maria, 97105-900, Santa Maria, RS, Brazil

Correspondence e-mail: rburrow@ewald.base.ufsm.br, mmartins@base.ufsm.br


Received 27 August 2007; accepted 13 September 2007

Key indicators: single-crystal X-ray study; $T = 273\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; disorder in main residue; R factor = 0.048; wR factor = 0.142; data-to-parameter ratio = 15.9.

The title compound, $C_{12}H_{10}N_2O$, is approximately planar. The angle between the quinoline and 4,5-dihydrooxazole ring systems is $11.91(12)^\circ$. The molecules pack into a herringbone array with no significant $\pi-\pi$ interactions. The dihydrooxazole N and O atoms are disordered over two positions, with almost equal site occupancy factors.

Related literature

For related 2-substituted quinoline compounds, see: Mague *et al.* (1997); Yang *et al.* (2001); Qi *et al.* (2003); Xu *et al.* (2006). For the synthesis, see: Ishihara & Togo (2007). For related literature, see: Allen (2002); Cunico *et al.* (2006); Hartline *et al.* (2005).

**Experimental***Crystal data*

$C_{12}H_{10}N_2O$

$M_r = 198.22$

Monoclinic, $P2_1/c$

$a = 6.2240(3)\text{ \AA}$

$b = 13.6649(6)\text{ \AA}$

$c = 11.8186(6)\text{ \AA}$

$\beta = 102.097(3)^\circ$

$V = 982.86(8)\text{ \AA}^3$

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 0.09\text{ mm}^{-1}$

$T = 273(2)\text{ K}$

$0.24 \times 0.21 \times 0.13\text{ mm}$

Data collection

Bruker APEXII CCD area-detector diffractometer

Absorption correction: multi-scan (*SADABS*; Bruker, 2006)

$T_{\min} = 0.705$, $T_{\max} = 1$

(expected range = 0.697–0.989)

9874 measured reflections

2183 independent reflections

1128 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.039$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$

$wR(F^2) = 0.143$

$S = 1.00$

2183 reflections

137 parameters

H-atom parameters constrained

$\Delta\rho_{\max} = 0.14\text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.15\text{ e \AA}^{-3}$

Data collection: *APEX2/COSMO/BIS* (Bruker, 2006); cell refinement: *SAINT* (Bruker, 2006); data reduction: *SAINT* and *SADABS* (Bruker, 2006); program(s) used to solve structure: *SHELXTL* (Bruker, 2001); program(s) used to refine structure: *SHELXTL*; molecular graphics: *DIAMOND* (Brandenburg, 2007); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors thank the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq/PRONEX) and Fundação de Amparo à Pesquisa do Estado do Rio Grande do Sul (FAPERGS) for financial support. Fellowships from CNPq and CAPES are also acknowledged. The diffractometer was funded by a CT-INFRA grant from the Financiadora de Estudos e Projetos (FINEP), Brazil.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2491).

References

- Allen, F. H. (2002). *Acta Cryst. B* **58**, 380–388.
- Brandenburg, K. (2007). *DIAMOND*. Version 3.1e. Crystal Impact Gbr, Bonn, Germany.
- Bruker (2001). *SHELXTL*. Version 6.10. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2006). *APEX2* (Version 2.1), *COSMO* (Version 1.56), *BIS* (Version 2.0.1.9), *SAINT* (Version 7.34A) and *SADABS* (Version 2004/1). Bruker AXS Inc., Madison, Wisconsin, USA.
- Cunico, W., Cechinel, C. A., Bonacorso, H. G., Martins, M. A. P., Zanatta, N., de Souza, M. V. N., Freitas, I. O., Soares, R. P. P. & Krettli, A. U. (2006). *Bioorg. Med. Chem. Lett.* **16**, 649–653.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Hartline, C. B., Harden, E. A., Williams-Aziz, S. L., Kushner, N. L., Brideau, R. J. & Kern, E. R. (2005). *Antiviral Res.* **65**, 97–105.
- Ishihara, M. & Togo, H. (2007). *Tetrahedron*, **63**, 1474–1480.
- Mague, J. T., Vang, S., Berge, D. G. & Wacholtz, W. F. (1997). *Acta Cryst. C* **53**, 973–979.
- Qi, J. Y., Qiu, L. Q., Yang, Q. Y., Zhou, Z. Y. & Chan, A. S. C. (2003). *Acta Cryst. E* **59**, o104–o105.
- Xu, R.-H., Zhou, J., Xu, Y., Qi, L., Shen, X. & Zhu, D.-R. (2006). *Acta Cryst. E* **62**, o5234–o5235.
- Yang, Q.-Y., Zhou, Z.-Y. & Qi, J.-Y. (2001). *Acta Cryst. E* **57**, o971–o972.

supporting information

Acta Cryst. (2008). E64, o154 [https://doi.org/10.1107/S1600536807045023]

2-(4,5-Dihydro-1,3-oxazol-2-yl)quinoline

Pablo Machado, Fernanda A. Rosa, Rubia M. S. da Silva, Robert A. Burrow and Marcos A. P. Martins

S1. Comment


Quinoline derivatives are known to possess a variety of biological properties such as antimalarial and antiviral activity (Cunico *et al.*, 2006; Hartline *et al.*, 2005). In addition, oxazoline-derived complexes of Pd(II) and other metals have attracted a great deal of attention due to their high efficiency in enantioselective catalysis. In this context, we were interested in detailed knowledge of the molecular structure of the above derivatives. In this communication we report the crystal structure of the title compound, 2-(4,5-dihydrooxazol-2-yl)quinoline. The analysis was focused on the planarity of the molecule. Both the quinoline and the 4,5-dihydrooxazole rings are planar, with r.m.s. deviations from planarity of 0.0136 Å for the first and 0.0176 ° for the last. The entire molecule is almost planar; the angle between the two rings is 11.91 (12) °. Quinoline and dihydrooxazole rings are essentially planar, with an average r.m.s. deviation from planarity of 0.06 (3) Å for 552 observations for the first and 0.05 (3) Å for 31 observations for the last in the Cambridge Structural Database [CSD, Version 5.28, update of May 2007; Allen, 2002)]. The molecules pack into a herringbone array with no significant π - π interactions.

S2. Experimental

The title compound was synthesized from quinoline-2-carbaldehyde and aminoethanol according to the general procedure of Ishihara & Togo (2007). The crystal used for the data collection was obtained by recrystallization from hexane followed by slow evaporation at room temperature.

S3. Refinement

All H atoms were refined using a riding model, with C—H = 0.97%Å and $U_{\text{iso}}(\text{H}) = 1.2\text{U}_{\text{eq}}(\text{C})$ for the methylene C atoms and C—H = 0.93 Å and $U_{\text{iso}} = 1.2\text{U}_{\text{eq}}(\text{C})$ for the quinoline C atoms.

**Figure 1**

View of the asymmetric unit of the title compound, showing the atom labelling scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are represented by circles of arbitrary radii.

2-(4,5-Dihydro-1,3-oxazol-2-yl)quinoline

Crystal data

$C_{12}H_{10}N_2O$
 $M_r = 198.22$
Monoclinic, $P2_1/c$
Hall symbol: -P 2ybc
 $a = 6.2240 (3)$ Å
 $b = 13.6649 (6)$ Å
 $c = 11.8186 (6)$ Å
 $\beta = 102.097 (3)^\circ$
 $V = 982.86 (8)$ Å³
 $Z = 4$

$F(000) = 416$
 $D_x = 1.34 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 1185 reflections
 $\theta = 6.9\text{--}40.4^\circ$
 $\mu = 0.09 \text{ mm}^{-1}$
 $T = 273 \text{ K}$
Block, colourless
 $0.24 \times 0.21 \times 0.13 \text{ mm}$

Data collection

Bruker APEXII CCD area-detector
diffractometer
Graphite monochromator
phi and ω scans
Absorption correction: multi-scan
(SADABS; Bruker, 2006) was used to perform
the multi-scan semi-empirical (using intensity
measurements) absorption correction and to
scale the data.

$T_{\min} = 0.705, T_{\max} = 1$
9874 measured reflections
2183 independent reflections
1128 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.039$
 $\theta_{\max} = 27.5^\circ, \theta_{\min} = 3.4^\circ$
 $h = -7 \rightarrow 8$
 $k = -16 \rightarrow 16$
 $l = -15 \rightarrow 15$

*Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.048$ $wR(F^2) = 0.143$ $S = 1.00$

2183 reflections

137 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.069P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\text{max}} < 0.001$ $\Delta\rho_{\text{max}} = 0.14 \text{ e } \text{\AA}^{-3}$ $\Delta\rho_{\text{min}} = -0.15 \text{ e } \text{\AA}^{-3}$ *Special details*

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C1	0.0592 (3)	0.64386 (12)	0.61057 (14)	0.0526 (5)	
C2	-0.0108 (3)	0.68102 (15)	0.49740 (15)	0.0664 (5)	
H2	-0.1414	0.6592	0.4508	0.080*	
C3	0.1126 (3)	0.74813 (14)	0.45761 (15)	0.0671 (6)	
H3	0.0666	0.7738	0.3836	0.080*	
C4	0.3119 (3)	0.77950 (12)	0.52818 (13)	0.0531 (5)	
C5	0.4498 (3)	0.85086 (13)	0.49471 (15)	0.0644 (5)	
H5	0.4114	0.8792	0.4217	0.077*	
C6	0.6375 (4)	0.87843 (14)	0.56788 (18)	0.0701 (6)	
H6	0.7272	0.9256	0.5448	0.084*	
C7	0.6978 (3)	0.83631 (15)	0.67849 (17)	0.0729 (6)	
H7	0.8267	0.8561	0.7283	0.087*	
C8	0.5700 (3)	0.76724 (14)	0.71306 (15)	0.0644 (5)	
H8	0.6128	0.7392	0.7860	0.077*	
C9	0.3723 (3)	0.73727 (12)	0.63946 (13)	0.0514 (5)	
C11	-0.0822 (3)	0.57463 (13)	0.65666 (15)	0.0591 (5)	
C12	-0.2299 (4)	0.48788 (16)	0.77775 (19)	0.0812 (7)	
H12A	-0.3146	0.5189	0.8280	0.097*	
H12B	-0.1788	0.4247	0.8099	0.097*	
C13	-0.3652 (4)	0.47695 (17)	0.65699 (18)	0.0815 (7)	
H13A	-0.3657	0.4096	0.6312	0.098*	
H13B	-0.5156	0.4977	0.6533	0.098*	
N1	0.2441 (2)	0.67018 (10)	0.67998 (11)	0.0538 (4)	
N2	-0.2574 (3)	0.54003 (12)	0.58733 (13)	0.0854 (7)	0.473 (17)
O1	-0.0448 (2)	0.54973 (11)	0.76424 (12)	0.0751 (7)	0.473 (17)

O1A	-0.2574 (3)	0.54003 (12)	0.58733 (13)	0.0854 (7)	0.527 (17)
N2A	-0.0448 (2)	0.54973 (11)	0.76424 (12)	0.0751 (7)	0.527 (17)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0561 (11)	0.0528 (10)	0.0477 (10)	0.0063 (9)	0.0085 (9)	-0.0031 (8)
C2	0.0650 (13)	0.0777 (13)	0.0509 (11)	-0.0018 (11)	-0.0007 (9)	0.0018 (9)
C3	0.0771 (14)	0.0744 (14)	0.0450 (10)	0.0102 (11)	0.0019 (10)	0.0063 (8)
C4	0.0640 (12)	0.0516 (10)	0.0448 (9)	0.0099 (9)	0.0141 (9)	0.0004 (7)
C5	0.0796 (14)	0.0627 (12)	0.0534 (11)	0.0080 (11)	0.0194 (11)	0.0074 (9)
C6	0.0747 (14)	0.0647 (13)	0.0751 (13)	-0.0037 (11)	0.0253 (11)	0.0090 (10)
C7	0.0706 (14)	0.0771 (14)	0.0676 (13)	-0.0116 (11)	0.0071 (11)	0.0052 (10)
C8	0.0699 (14)	0.0682 (13)	0.0515 (10)	-0.0032 (10)	0.0047 (10)	0.0087 (8)
C9	0.0583 (11)	0.0506 (10)	0.0439 (9)	0.0083 (9)	0.0075 (8)	-0.0007 (7)
C11	0.0600 (12)	0.0562 (11)	0.0604 (12)	0.0043 (9)	0.0112 (10)	-0.0050 (9)
C12	0.0734 (15)	0.0889 (15)	0.0850 (15)	-0.0119 (12)	0.0249 (12)	0.0050 (11)
C13	0.0698 (14)	0.0826 (15)	0.0923 (16)	-0.0137 (11)	0.0178 (13)	0.0012 (11)
N1	0.0580 (9)	0.0536 (9)	0.0481 (8)	0.0009 (7)	0.0075 (7)	0.0019 (6)
N2	0.0765 (12)	0.1034 (13)	0.0690 (11)	-0.0272 (9)	-0.0013 (9)	0.0046 (8)
O1	0.0772 (12)	0.0848 (12)	0.0607 (10)	-0.0168 (8)	0.0083 (8)	0.0091 (7)
O1A	0.0765 (12)	0.1034 (13)	0.0690 (11)	-0.0272 (9)	-0.0013 (9)	0.0046 (8)
N2A	0.0772 (12)	0.0848 (12)	0.0607 (10)	-0.0168 (8)	0.0083 (8)	0.0091 (7)

Geometric parameters (\AA , ^\circ)

C1—N1	1.315 (2)	C7—H7	0.9300
C1—C2	1.412 (2)	C8—C9	1.411 (2)
C1—C11	1.472 (3)	C8—H8	0.9300
C2—C3	1.343 (3)	C9—N1	1.366 (2)
C2—H2	0.9300	C11—O1	1.289 (2)
C3—C4	1.409 (2)	C11—N2	1.307 (2)
C3—H3	0.9300	C12—O1	1.464 (2)
C4—C5	1.409 (2)	C12—C13	1.505 (3)
C4—C9	1.413 (2)	C12—H12A	0.9700
C5—C6	1.353 (2)	C12—H12B	0.9700
C5—H5	0.9300	C13—N2	1.450 (2)
C6—C7	1.405 (2)	C13—H13A	0.9700
C6—H6	0.9300	C13—H13B	0.9700
C7—C8	1.352 (3)		
N1—C1—C2	123.27 (18)	C9—C8—H8	119.7
N1—C1—C11	117.23 (15)	N1—C9—C8	118.48 (15)
C2—C1—C11	119.46 (16)	N1—C9—C4	122.63 (16)
C3—C2—C1	119.45 (17)	C8—C9—C4	118.87 (18)
C3—C2—H2	120.3	O1—C11—N2	118.51 (18)
C1—C2—H2	120.3	O1—C11—C1	122.40 (16)
C2—C3—C4	119.85 (16)	N2—C11—C1	119.03 (16)

C2—C3—H3	120.1	O1—C12—C13	104.53 (16)
C4—C3—H3	120.1	O1—C12—H12A	110.8
C3—C4—C5	123.84 (16)	C13—C12—H12A	110.8
C3—C4—C9	117.14 (18)	O1—C12—H12B	110.8
C5—C4—C9	119.01 (17)	C13—C12—H12B	110.8
C6—C5—C4	120.54 (17)	H12A—C12—H12B	108.9
C6—C5—H5	119.7	N2—C13—C12	104.22 (15)
C4—C5—H5	119.7	N2—C13—H13A	110.9
C5—C6—C7	120.47 (19)	C12—C13—H13A	110.9
C5—C6—H6	119.8	N2—C13—H13B	110.9
C7—C6—H6	119.8	C12—C13—H13B	110.9
C8—C7—C6	120.50 (18)	H13A—C13—H13B	108.9
C8—C7—H7	119.7	C1—N1—C9	117.63 (14)
C6—C7—H7	119.7	C11—N2—C13	106.47 (15)
C7—C8—C9	120.60 (17)	C11—O1—C12	106.02 (16)
C7—C8—H8	119.7		
N1—C1—C2—C3	-1.4 (3)	N1—C1—C11—O1	8.9 (3)
C11—C1—C2—C3	176.45 (16)	C2—C1—C11—O1	-169.11 (17)
C1—C2—C3—C4	1.1 (3)	N1—C1—C11—N2	-173.99 (16)
C2—C3—C4—C5	-178.70 (17)	C2—C1—C11—N2	8.0 (3)
C2—C3—C4—C9	0.2 (3)	O1—C12—C13—N2	-4.9 (2)
C3—C4—C5—C6	179.00 (17)	C2—C1—N1—C9	0.3 (2)
C9—C4—C5—C6	0.2 (3)	C11—C1—N1—C9	-177.55 (14)
C4—C5—C6—C7	0.0 (3)	C8—C9—N1—C1	179.31 (15)
C5—C6—C7—C8	0.4 (3)	C4—C9—N1—C1	1.0 (2)
C6—C7—C8—C9	-0.9 (3)	O1—C11—N2—C13	-1.2 (2)
C7—C8—C9—N1	-177.34 (17)	C1—C11—N2—C13	-178.42 (15)
C7—C8—C9—C4	1.1 (3)	C12—C13—N2—C11	3.8 (2)
C3—C4—C9—N1	-1.3 (2)	N2—C11—O1—C12	-2.2 (2)
C5—C4—C9—N1	177.67 (14)	C1—C11—O1—C12	174.98 (16)
C3—C4—C9—C8	-179.57 (15)	C13—C12—O1—C11	4.4 (2)
C5—C4—C9—C8	-0.6 (2)		