

# Bis(4-aminopyridinium) bis(oxalato- $\kappa^2O,O'$ )cuprate(II) dihydrate

 Zi-Cai Pan,<sup>a</sup> Kou-Lin Zhang<sup>a</sup> and Seik Weng Ng<sup>b\*</sup>
<sup>a</sup>College of Chemistry and Chemical Engineering, Yangzhou University, Yangzhou 225002, People's Republic of China, and <sup>b</sup>Department of Chemistry, University of Malaya, 50603, Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 19 November 2007; accepted 20 November 2007

 Key indicators: single-crystal X-ray study;  $T = 295$  K; mean  $\sigma(C-C) = 0.004$  Å;  $R$  factor = 0.037;  $wR$  factor = 0.109; data-to-parameter ratio = 10.4.

The Cu<sup>II</sup> atom in the title salt, (C<sub>5</sub>H<sub>7</sub>N<sub>2</sub>)<sub>2</sub>[Cu(C<sub>2</sub>O<sub>4</sub>)<sub>2</sub>]·2H<sub>2</sub>O, is located on a center of inversion and is chelated by two oxalate groups in a square-planar coordination geometry. The cation, anion and water molecules interact through hydrogen bonds, forming a three-dimensional hydrogen-bonded network.

## Related literature

See Geiser *et al.* (1987) for the square-planar pyridinium dioxalato cuprate(II) oxalic acid co-crystal. See Sun *et al.* (2004) for 2,6-bis(4'-pyridyl-1'-pyridinium)pyrazine bis(bis(oxalato)cuprate(II)), which is also square planar. In bis(2-aminoanilinium) bis(oxalato)cuprate(II), the amino groups coordinate to the metal atom, which exhibits octahedral coordination (Keene *et al.*, 2003).



## Experimental

### Crystal data

 (C<sub>5</sub>H<sub>7</sub>N<sub>2</sub>)<sub>2</sub>[Cu(C<sub>2</sub>O<sub>4</sub>)<sub>2</sub>]·2H<sub>2</sub>O

 $M_r = 465.86$ 

 Monoclinic,  $P2_1/c$ 
 $a = 3.7105$  (3) Å

 $b = 20.311$  (1) Å

 $c = 11.9261$  (9) Å

 $\beta = 90.450$  (1)°

 $V = 898.8$  (1) Å<sup>3</sup>
 $Z = 2$ 

 Mo  $K\alpha$  radiation

 $\mu = 1.28$  mm<sup>-1</sup>
 $T = 295$  (2) K

 $0.14 \times 0.10 \times 0.08$  mm

### Data collection

Bruker SMART area-detector

diffractometer

Absorption correction: multi-scan

(SADABS; Sheldrick, 1996)

 $T_{\min} = 0.764$ ,  $T_{\max} = 0.905$ 

2623 measured reflections

1590 independent reflections

 1498 reflections with  $I > 2\sigma(I)$ 
 $R_{\text{int}} = 0.016$ 

### Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.037$ 
 $wR(F^2) = 0.109$ 
 $S = 1.12$ 

1590 reflections

153 parameters

5 restraints

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{\text{max}} = 0.23$  e Å<sup>-3</sup>
 $\Delta\rho_{\text{min}} = -0.49$  e Å<sup>-3</sup>
**Table 1**

Selected geometric parameters (Å, °).

Cu1—O1	1.932 (2)	Cu1—O3	1.927 (2)
O1—Cu1—O3	85.4 (1)		

**Table 2**

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1w—H11 $\cdots$ O1	0.85 (3)	2.03 (2)	2.852 (3)	164 (5)
O1w—H12 $\cdots$ O3 <sup>i</sup>	0.85 (3)	2.12 (5)	2.931 (3)	160 (5)
N1—H1 $\cdots$ O2	0.85 (3)	2.12 (2)	2.858 (4)	146 (4)
N2—H21 $\cdots$ O4 <sup>ii</sup>	0.85 (1)	2.07 (1)	2.906 (4)	168 (4)
N2—H22 $\cdots$ O1w <sup>iii</sup>	0.85 (3)	2.02 (3)	2.867 (4)	176 (4)

 Symmetry codes: (i)  $-x + 2, -y + 1, -z + 1$ ; (ii)  $x + 1, -y + \frac{1}{2}, z - \frac{1}{2}$ ; (iii)  $-x + 3, -y + 1, -z$ .

Data collection: SMART (Bruker, 2000); cell refinement: SAINT (Bruker, 2000); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 1997); program(s) used to refine structure: SHELXL97 (Sheldrick, 1997); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2008).

We thank the Foundation of Jiangsu Provincial Key Program of Physical Chemistry in Yangzhou University, China, and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2384).

## References

- Barbour, L. J. (2001). *J. Supramol. Chem.*, **1**, 189–191.  
 Bruker (2000). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Geiser, U., Ramakrishna, B. L., Willett, R. D., Hulsberg, F. B. & Reedijk, J. (1987). *Inorg. Chem.* **26**, 3750–3756.  
 Keene, T. D., Hursthouse, M. B. & Price, D. J. (2003). *Acta Cryst.* **E59**, m1131–m1133.  
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.  
 Sheldrick, G. M. (1997). SHELXS97 and SHELXL97. University of Göttingen, Germany.  
 Sun, Y.-Q., Zhang, J., Chen, J.-L. & Yang, G.-Y. (2004). *Eur. J. Inorg. Chem.* pp. 3837–3841.  
 Westrip, S. P. (2008). publCIF. In preparation.

**supplementary materials**

*Acta Cryst.* (2008). E64, m221 [ doi:10.1107/S160053680706117X ]

## Bis(4-aminopyridinium) bis(oxalato- $\kappa^2O,O'$ )cuprate(II) dihydrate

Z.-C. Pan, K.-L. Zhang and S. W. Ng

### Comment

There are many crystallographic studies of coordination compounds of oxalic acid (Cambridge Structural Database, Version 5.28, Nov. 2006). The copper(II) center in the title compound shows square-planar coordination (Table 1); the cations, anions and lattice water molecules interact through hydrogen bonds (Table 2) to give rise to a three-dimensional network motif.

### Experimental

Potassium oxalate monohydrate (0.036 g, 0.2 mmol) dissolved in water (5 ml) was reacted with copper nitrate trihydrate (0.048 g, 0.2 mmol) in water (5 ml). To this solution was added 4-C<sub>5</sub>H<sub>4</sub>N-NH-C(O)-C(O)-NH-4-C<sub>5</sub>H<sub>4</sub>N (0.048 g, 0.2 mmol) dissolved in methanol (15 ml). Blue crystals separated after a few days in 60% yield. CH&N elemental analysis. Calc. for C<sub>14</sub>H<sub>18</sub>CuN<sub>4</sub>O<sub>10</sub>: C 36.09, H 3.89, N 12.02%. Found: C 36.43, H 3.74, N 12.18%.

### Refinement

The carbon-bound H atoms were placed in calculated positions and were allowed to ride on the parent atoms. The oxygen- and nitrogen-bound H atoms were refined with a distance restraint O-H = N-H = 0.85±0.01 Å. Their temperature factors were freely refined.

### Figures




Fig. 1. Thermal ellipsoid plot of 2[C<sub>5</sub>H<sub>7</sub>N<sub>2</sub>]<sup>+</sup>[Cu(C<sub>2</sub>O<sub>4</sub>)<sub>2</sub>]<sup>2-</sup>·2H<sub>2</sub>O; Displacement ellipsoids are drawn at the 50% probability level, and H atoms as spheres of arbitrary radii.

## Bis(4-aminopyridinium) bis(oxalato- $\kappa^2O,O'$ )cuprate(II) dihydrate

### Crystal data

(C<sub>5</sub>H<sub>7</sub>N<sub>2</sub>)<sub>2</sub>[Cu(C<sub>2</sub>O<sub>4</sub>)<sub>2</sub>]·2H<sub>2</sub>O

*M<sub>r</sub>* = 465.86

Monoclinic, *P*2<sub>1</sub>/*c*

Hall symbol: -*P* 2ybc

*a* = 3.7105 (3) Å

*b* = 20.311 (1) Å

*F*<sub>000</sub> = 478

*D<sub>x</sub>* = 1.721 Mg m<sup>-3</sup>

Mo *K*α radiation

λ = 0.71073 Å

Cell parameters from 2198 reflections

θ = 2.0–25.1°

μ = 1.28 mm<sup>-1</sup>

# supplementary materials

---

$c = 11.9261(9) \text{ \AA}$   
 $\beta = 90.450(1)^\circ$   
 $V = 898.8(1) \text{ \AA}^3$   
 $Z = 2$

$T = 295(2) \text{ K}$   
Block, blue  
 $0.14 \times 0.10 \times 0.08 \text{ mm}$

## Data collection

Bruker SMART area-detector diffractometer	1590 independent reflections
Radiation source: fine-focus sealed tube	1498 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.016$
$T = 295(2) \text{ K}$	$\theta_{\text{max}} = 25.1^\circ$
$\varphi$ and $\omega$ scans	$\theta_{\text{min}} = 2.0^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -4 \rightarrow 4$
$T_{\text{min}} = 0.764, T_{\text{max}} = 0.905$	$k = -21 \rightarrow 24$
2623 measured reflections	$l = -8 \rightarrow 14$

## Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.037$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.109$	$w = 1/[\sigma^2(F_o^2) + (0.0536P)^2 + 1.2237P]$
$S = 1.12$	where $P = (F_o^2 + 2F_c^2)/3$
1590 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
153 parameters	$\Delta\rho_{\text{max}} = 0.23 \text{ e \AA}^{-3}$
5 restraints	$\Delta\rho_{\text{min}} = -0.49 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

## Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.5000	0.5000	0.5000	0.0283 (2)
O1	0.8063 (6)	0.52464 (11)	0.37661 (18)	0.0352 (5)
O2	1.0084 (7)	0.48225 (12)	0.21648 (19)	0.0399 (6)
O3	0.4985 (6)	0.41370 (10)	0.43287 (17)	0.0313 (5)
O4	0.7190 (7)	0.36328 (11)	0.28166 (19)	0.0386 (5)
O1w	1.1418 (8)	0.65099 (13)	0.3772 (2)	0.0477 (6)
N1	1.1417 (8)	0.37249 (17)	0.0727 (2)	0.0439 (7)
N2	1.5342 (8)	0.27528 (14)	-0.1995 (2)	0.0365 (6)
C1	0.8457 (8)	0.47791 (15)	0.3044 (2)	0.0275 (6)
C2	0.6748 (8)	0.41168 (14)	0.3400 (2)	0.0275 (6)
C3	1.1500 (8)	0.3067 (2)	0.0721 (3)	0.0411 (8)

H3	1.0674	0.2837	0.1343	0.049*
C4	1.2757 (8)	0.27283 (17)	-0.0171 (3)	0.0354 (7)
H4	1.2776	0.2270	-0.0160	0.043*
C5	1.4040 (7)	0.30699 (15)	-0.1115 (2)	0.0266 (6)
C6	1.3848 (8)	0.37643 (16)	-0.1090 (3)	0.0344 (7)
H6	1.4609	0.4011	-0.1702	0.041*
C7	1.2545 (9)	0.40679 (17)	-0.0167 (3)	0.0428 (8)
H7	1.2429	0.4525	-0.0151	0.051*
H11	1.045 (13)	0.6140 (13)	0.363 (4)	0.084 (17)*
H12	1.268 (11)	0.642 (3)	0.434 (3)	0.077 (17)*
H1	1.068 (10)	0.3915 (18)	0.132 (2)	0.046 (11)*
H21	1.563 (10)	0.2337 (6)	-0.199 (3)	0.042 (10)*
H22	1.621 (11)	0.2979 (18)	-0.253 (2)	0.055 (12)*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cu1	0.0380 (3)	0.0234 (3)	0.0237 (3)	-0.00362 (19)	0.0082 (2)	-0.00456 (18)
O1	0.0493 (13)	0.0257 (11)	0.0309 (11)	-0.0074 (10)	0.0139 (10)	-0.0061 (9)
O2	0.0526 (14)	0.0358 (12)	0.0314 (12)	-0.0023 (11)	0.0165 (11)	-0.0029 (10)
O3	0.0416 (12)	0.0258 (11)	0.0265 (11)	-0.0047 (9)	0.0058 (9)	-0.0030 (8)
O4	0.0550 (14)	0.0256 (11)	0.0354 (12)	-0.0043 (10)	0.0117 (10)	-0.0088 (9)
O1w	0.0663 (17)	0.0358 (14)	0.0411 (14)	-0.0106 (12)	-0.0018 (13)	0.0083 (11)
N1	0.0379 (15)	0.064 (2)	0.0295 (15)	0.0085 (14)	-0.0009 (12)	-0.0171 (14)
N2	0.0476 (16)	0.0320 (15)	0.0300 (14)	0.0043 (12)	0.0065 (12)	-0.0016 (12)
C1	0.0325 (15)	0.0265 (15)	0.0236 (15)	0.0023 (12)	0.0016 (12)	-0.0007 (11)
C2	0.0310 (14)	0.0277 (15)	0.0240 (14)	0.0019 (11)	0.0004 (12)	-0.0020 (12)
C3	0.0321 (16)	0.065 (2)	0.0265 (16)	0.0059 (15)	-0.0001 (13)	0.0060 (16)
C4	0.0328 (15)	0.0393 (17)	0.0342 (16)	0.0024 (13)	-0.0023 (13)	0.0077 (14)
C5	0.0248 (13)	0.0307 (15)	0.0244 (14)	0.0030 (11)	-0.0051 (11)	-0.0015 (12)
C6	0.0353 (16)	0.0330 (16)	0.0348 (17)	0.0010 (13)	0.0012 (13)	0.0021 (13)
C7	0.0436 (18)	0.0353 (18)	0.049 (2)	0.0056 (14)	-0.0018 (16)	-0.0133 (16)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

Cu1—O1	1.932 (2)	N2—C5	1.325 (4)
Cu1—O3	1.927 (2)	N2—H21	0.85 (1)
Cu1—O3 <sup>i</sup>	1.927 (2)	N2—H22	0.85 (3)
Cu1—O1 <sup>i</sup>	1.932 (2)	C1—C2	1.548 (4)
O1—C1	1.290 (4)	C3—C4	1.352 (5)
O2—C1	1.217 (4)	C3—H3	0.9300
O3—C2	1.291 (4)	C4—C5	1.409 (4)
O4—C2	1.216 (4)	C4—H4	0.9300
O1w—H11	0.85 (3)	C5—C6	1.413 (4)
O1w—H12	0.85 (3)	C6—C7	1.355 (5)
N1—C3	1.337 (5)	C6—H6	0.9300
N1—C7	1.343 (5)	C7—H7	0.9300
N1—H1	0.85 (3)		

## supplementary materials

O3—Cu1—O3 <sup>i</sup>	180	O4—C2—O3	126.0 (3)
O3—Cu1—O1 <sup>i</sup>	94.7 (1)	O4—C2—C1	119.2 (3)
O3 <sup>i</sup> —Cu1—O1 <sup>i</sup>	85.4 (1)	O3—C2—C1	114.8 (2)
O1—Cu1—O3	85.4 (1)	N1—C3—C4	121.4 (3)
O3 <sup>i</sup> —Cu1—O1	94.7 (1)	N1—C3—H3	119.3
O1 <sup>i</sup> —Cu1—O1	180	C4—C3—H3	119.3
C1—O1—Cu1	112.83 (19)	C3—C4—C5	119.9 (3)
C2—O3—Cu1	112.66 (18)	C3—C4—H4	120.0
H11—O1w—H12	101 (5)	C5—C4—H4	120.0
C3—N1—C7	120.4 (3)	N2—C5—C4	121.4 (3)
C3—N1—H1	118 (3)	N2—C5—C6	121.4 (3)
C7—N1—H1	122 (3)	C4—C5—C6	117.2 (3)
C5—N2—H21	122 (3)	C7—C6—C5	119.3 (3)
C5—N2—H22	118 (3)	C7—C6—H6	120.3
H21—N2—H22	120 (4)	C5—C6—H6	120.3
O2—C1—O1	125.5 (3)	N1—C7—C6	121.7 (3)
O2—C1—C2	120.4 (3)	N1—C7—H7	119.2
O1—C1—C2	114.0 (2)	C6—C7—H7	119.2
O3—Cu1—O1—C1	5.3 (2)	O2—C1—C2—O3	-177.7 (3)
O3 <sup>i</sup> —Cu1—O1—C1	-174.7 (2)	O1—C1—C2—O3	4.2 (4)
O1 <sup>i</sup> —Cu1—O3—C2	177.2 (2)	C7—N1—C3—C4	1.0 (5)
O1—Cu1—O3—C2	-2.8 (2)	N1—C3—C4—C5	0.5 (5)
Cu1—O1—C1—O2	175.7 (3)	C3—C4—C5—N2	178.9 (3)
Cu1—O1—C1—C2	-6.3 (3)	C3—C4—C5—C6	-1.7 (4)
Cu1—O3—C2—O4	179.1 (3)	N2—C5—C6—C7	-179.1 (3)
Cu1—O3—C2—C1	0.2 (3)	C4—C5—C6—C7	1.6 (4)
O2—C1—C2—O4	3.3 (4)	C3—N1—C7—C6	-1.1 (5)
O1—C1—C2—O4	-174.8 (3)	C5—C6—C7—N1	-0.2 (5)

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ .

### Hydrogen-bond geometry ( $\text{\AA}, ^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1w—H11 $\cdots$ O1	0.85 (3)	2.03 (2)	2.852 (3)	164 (5)
O1w—H12 $\cdots$ O3 <sup>ii</sup>	0.85 (3)	2.12 (5)	2.931 (3)	160 (5)
N1—H1 $\cdots$ O2	0.85 (3)	2.12 (2)	2.858 (4)	146 (4)
N2—H21 $\cdots$ O4 <sup>iii</sup>	0.85 (1)	2.07 (1)	2.906 (4)	168 (4)
N2—H22 $\cdots$ O1w <sup>iv</sup>	0.85 (3)	2.02 (3)	2.867 (4)	176 (4)

Symmetry codes: (ii)  $-x+2, -y+1, -z+1$ ; (iii)  $x+1, -y+1/2, z-1/2$ ; (iv)  $-x+3, -y+1, -z$ .

Fig. 1

